

Arithmetic cutoff bounds and sharp tail capacity for local Nyman–Beurling certificates

Scott Schubr
Luminary Oddities

October 27, 2026

Abstract

We give complete arithmetic receivers for local Nyman–Beurling projection gain in the whole-line Poisson norm. Finite mixed-order Bernoulli covariances retain all complex phases and diagonalize in common divisor coordinates. Their height-scale limit is a positive cotangent form with a declared resonance boundary. A complete Farey-operator estimate, including variation of the primitive amplitudes and the exact original weight, extends its uniform support regime to $N = o(T)$ at every fixed wall gap $K \sim \gamma NT$, $\gamma > 1/(2\pi)$. Combining this norm comparison with the compensated mean gives explicit bounds for the actual optimized double-null gain in terms of a corrected finite model, including a uniform additive error. We also prove a limitation of possible constructions: on every fixed finite geometric module, optimized over arbitrary degrees and coefficients, true gain falls below each positive threshold on a positive-lower-density set of actual heights. A quantitative evaluation defect describes the necessary escape from such modules. The sharp fixed-trial gcd capacity, variance-priced comparison, evaluation dual, necessary support growth and robust constrained certificate are retained. Complete full-head examples and exact Hardy continuation return successful finite certificates to zero-free disks and covered height intervals. The resulting finite-model criterion retains the complete shared head, exact coefficient constraints and all approximation errors.

1 The local arithmetic problem

We study the complete finite arithmetic recipient of local Nyman–Beurling projection gain. The mixed-order receiver of Theorem 3.1 retains every Bernoulli covariance and pays its entire infinite remainder. Common divisor coordinates give positive explicit weights, whose height-scale resummation is a cotangent form. The variance-priced comparison in Section 5 remains a complete quantitative option. The Farey-operator return in Section 6 strengthens the uniform support regime to $N = o(T)$ throughout every fixed-gap pole-free wall.

The norm and numerator are then combined before optimizing. Section 15 approximates the actual optimized double-null gain by its corrected finite full model with explicit additive error. Its target implication prices both mean and norm errors in the same coefficient space. Section 17 rules out one plausible construction family: arbitrary geometric depth within fixed finite modules, even with phase-adapted coefficients and a finite menu of preconditioners. Its paid evaluation defect quantifies the required escape.

The exact support-cost dual, null adaptation, sharp fixed-parameter tail capacity, compensated mean and robust certificate remain complementary parts of the same method. The applications retain the original shared arithmetic head and complete tail. Local Hardy projection returns successful gains to zero-free disks and propagates each seed through a certified height interval.

The local Hardy method is classical [8]; so are fractional-part gcd covariance [10], Jordan/divisor factorization [11], Bernoulli summation [5], uniform high-parameter truncation [13, 14], absolute Farey block estimates [17, 18], robust ellipsoidal response optimization [6, 7], and zeta universality [20]. Bernoulli-2 Farey covariance occurs in [4]; the unshifted wavelet-tail large-sieve sandwich and prefix deflation in [19] are close predecessors. The source-dependent conclusions here concern the complete height-dependent mixed receiver, its paid return to optimized native gain, and the degree-uniform limitation for fixed arithmetic modules.

Fix throughout

$$0 < a < \frac{1}{2}, \quad \alpha = \frac{1}{2} + a, \quad \sigma = \frac{1}{2} - a, \quad \lambda = \alpha + iT, \quad q = \sigma + iT, \quad T \in \mathbb{R}. \quad (1)$$

Let $\mathcal{A}_N$ be the Dirichlet polynomials

$$A(s) = \sum_{d \leq N} c_d d^{-s}, \quad A(1) = \sum_{d \leq N} \frac{c_d}{d} = 0.$$

The coefficients may be complex and may depend on T . The integer N is a support cutoff, not the number of nonzero terms. Put

$$d_n = \sum_{d|n} c_d, \quad B_n = \sum_{j \leq n} d_j j^{-q}, \quad p_n = n^{-2a} - (n+1)^{-2a}, \quad (2)$$

where $c_d = 0$ beyond the support. Define

$$m(A) = \sum_{n \geq 1} p_n B_n, \quad Q(A) = \sum_{n \geq 1} p_n |B_n|^2, \quad g(A) = \frac{|m(A)|^2}{Q(A)} \quad (A \neq 0),$$

and $g_N(\lambda) = \sup_{A \in \mathcal{A}_N \setminus \{0\}} g(A)$. For a trivial coefficient space the supremum is defined to be zero. For every integer $M \geq 1$, put $K = M + 1 \geq 2$ and define the finite heads

$$m_M = \sum_{n \leq M} p_n B_n, \quad Q_M = \sum_{n \leq M} p_n |B_n|^2.$$

Proposition 1.1 (Complete recipient). *For every anchored finite polynomial, all these moments exist and*

$$m(A) = \zeta(\lambda) A(\lambda), \quad (3)$$

$$Q(A) = \frac{a}{\pi} \int_{\mathbb{R}} \frac{|\zeta(1/2 + it) A(1/2 + it)|^2}{a^2 + (t - T)^2} dt. \quad (4)$$

For $A \neq 0$, $Q(A) > 0$ and $0 \leq g(A) \leq 1$.

Proof. The summatory divisor sequence satisfies

$$D(x) = \sum_{j \leq x} d_j = \sum_d c_d \lfloor x/d \rfloor = - \sum_d c_d \{x/d\}, \quad |D(x)| \leq S_0 := \sum_d |c_d|.$$

Partial summation shows that $\sum d_n n^{-s}$ converges on $\Re s > 0$, agrees there by continuation with $\zeta(s) A(s)$, and that B_n is bounded at fixed q . Thus both moments converge. Summation by parts against p_n gives

$$\sum_{n \leq M} p_n B_n = \sum_{j \leq M} d_j j^{-(q+2a)} - (M+1)^{-2a} B_M.$$

The last term tends to zero, proving (3) since $q + 2a = \lambda$.

Initially on $\Re s > 1$, the Mellin identity is

$$\int_1^\infty B_{[x]} x^{q-s} \frac{dx}{x} = \frac{\zeta(s)A(s)}{s-q}.$$

Its bounded summatory representation continues to $\Re s > 1/2$. Mellin Plancherel on the boundary gives

$$\frac{1}{2\pi} \int_{\mathbb{R}} \left| \frac{\zeta(1/2 + it)A(1/2 + it)}{1/2 + it - q} \right|^2 dt = \int_1^\infty |B_{[x]}|^2 x^{-2a} \frac{dx}{x} = \frac{Q(A)}{2a},$$

which is (4). If $Q = 0$, positivity of p_n forces every B_n , hence every d_n and c_n , to vanish. Finally $\sum p_n = 1$, so Cauchy-Schwarz proves $|m|^2 \leq Q$. $\square$

Minimizing $\sum p_n |1 - zB_n|^2$ over $z \in \mathbb{C}$ gives $1 - g(A)$. This identifies g_N as an optimized projection gain, without an independence assumption on a chosen trial dictionary.

1.1 Periodic profiles and divisor coordinates

Write $B_j(x)$ for the Bernoulli polynomials, with $B_1(x) = x - 1/2$, and set

$$H_{r,d}(x) = -\frac{d^r}{(r+1)!} B_{r+1}(\{x/d\}), \quad r \geq 0, \quad H_{r,c} = \sum_d c_d H_{r,d}. \quad (5)$$

In particular $H_{0,d}(n) = 1/2 - \{n/d\}$ uses the inclusive arithmetic value at an integer seam. For $r \geq 1$ the profiles are continuous and $H'_{r,d} = H_{r-1,d}$ almost everywhere. Put

$$t_k = \sum_{d|k} \frac{c_d}{d}, \quad J_s(k) = \sum_{d|k} \mu(k/d) d^s.$$

Finite Möbius inversion gives $c_d = d \sum_{d|k} \mu(k/d) t_k$ and

$$A(s) = \sum_{k \leq N} t_k J_{1-s}(k), \quad t_1 = A(1) = 0. \quad (6)$$

The divisor coordinates in (6) will also diagonalize the complete mixed-order source. The exact classical variance and its full fixed-trial remainder are proved in Theorem 11.1.

2 Complete periodic expansions and uniform walls

Put $(q)_0 = 1$, $(q)_r = q(q+1) \cdots (q+r-1)$. The Bernoulli identities and Fourier bounds used here are classical [5].

Lemma 2.1 (Exact finite primitive formula). *If $A(1) = A(q) = 0$, then for every integer $R \geq 1$ and $n \geq 1$,*

$$B_n = n^{-q} \sum_{r=0}^R (q)_r n^{-r} H_{r,c}(n) - (q)_{R+1} \int_n^\infty H_{R,c}(x) x^{-q-R-1} dx. \quad (7)$$

For $r \geq 1$,

$$|H_{r,c}(x)| \leq \frac{4N^r S_0}{(2\pi)^{r+1}}, \quad |H_{0,c}(n)| \leq S_0/2.$$

Proof. The limiting Dirichlet sum is $\zeta(q)A(q) = 0$. Exact partial summation yields

$$B_n = D(n)n^{-q} - q \int_n^\infty D(x)x^{-q-1} dx = H_{0,c}(n)n^{-q} - q \int_n^\infty H_{0,c}(x)x^{-q-1} dx.$$

The constant $D - H_{0,c} = -\sum c_d/2$ cancels. Repeated integration by parts using $H'_{r,c} = H_{r-1,c}$ gives a plus sign at every endpoint and the final negative remainder in (7). All primitive boundaries at infinity vanish since $\sigma + r > 0$. Bernoulli's complete Fourier series gives $\|B_{r+1}\|_\infty/(r+1)! \leq 2\zeta(r+1)/(2\pi)^{r+1} < 4/(2\pi)^{r+1}$, proving the bounds. The endpoint $H_{0,c}(n)$ is not replaced by a Fourier midpoint. $\square$

Lemma 2.2 (A declared pole-free wall). *Choose $0 < \theta < 1$ and $R \geq 1$, put $L = |q| + R + 1$, and suppose*

$$K \geq \frac{NL}{2\pi\theta}. \quad (8)$$

Define

$$b_\theta = \frac{1}{2} + \frac{2}{\pi} \frac{\theta}{1-\theta}, \quad e_\theta = \frac{2}{\pi} (|q| + 2)\theta^R. \quad (9)$$

For every $n \geq K$, the finite profile sum in (7) has modulus at most $b_\theta S_0$, and its entire remainder has modulus at most $e_\theta S_0 n^{-\sigma}$. Consequently

$$Q - Q_M \leq (b_\theta + e_\theta)^2 S_0^2 / K, \quad |m - m_M| \leq (b_\theta + e_\theta) S_0 K^{-\alpha}.$$

Proof. Since $|(q)_r| \leq L^r$ for $r \leq R+1$, the endpoint sum is bounded by $S_0[1/2 + (2/\pi) \sum_{r=1}^R \theta^r] \leq b_\theta S_0$. The complete remainder, after division by $n^{-\sigma}$, is at most

$$\frac{2}{\pi} S_0 \frac{L}{\sigma + R} \left(\frac{LN}{2\pi n} \right)^R \leq e_\theta S_0,$$

using $L/(\sigma + R) \leq |q| + 2$. Sum against the complete p_n weights and use $\sum_{n \geq K} p_n = K^{-2a}$, $2\sigma + 2a = 1$, $\sigma + 2a = \alpha$. $\square$

For example $\theta = 1/2$ and $2^R \geq 8(|q| + 2)$ give the clean scalar receiver $|B_n| < (5/4)S_0 n^{-\sigma}$ after $n \geq N(|q| + R + 1)/\pi$. A shorter alternative paired-wave argument gives $1/2 + 2/\pi$ already at $n \geq N|q|/\pi$; it is stated with proof in Appendix A. Qualitative height-uniform truncation at a near-linear wall is classical [13, 14]. Here its role is the complete double-null arithmetic norm; the higher-order expansion also supplies the mixed covariance and its sharp height-scale law.

3 A complete mixed-order covariance receiver

For fixed R, q, K define the positive quadratic form

$$U_R(c) = \frac{2a}{K} \int_0^1 \text{av}_n \left| \sum_{r=0}^R (q)_r K^{-r} x^r H_{r,c}(n) \right|^2 dx. \quad (10)$$

The average may be taken over any common integer period. The computation below avoids enumeration of $\text{lcm}(1, \dots, N)$.

Theorem 3.1 (Complete mixed-order tail). *Under the exact double null and wall (8), put $b = b_\theta$, $e = e_\theta$ and*

$$\delta_R = \frac{[4aN^2 + 4a(a+1)]b^2}{K^2} + \frac{2be + e^2}{K}. \quad (11)$$

Then the whole infinite tail satisfies

$$|Q - Q_M - U_R(c)| \leq \delta_R S_0^2. \quad (12)$$

In particular $G_M + U_R + N\delta_R I$ is a complete norm upper on $\{c : A(1) = A(q) = 0\}$, where $c^* G_M c = Q_M$.

Proof. For fixed orders and support indices, the centered periodic product $H_{r,d}(n)\overline{H_{s,e}(n)} - \text{av } H_{r,d}\overline{H_{s,e}}$ has period $\text{lcm}(d, e) \leq N^2$. Its partial sums over any interval are bounded by twice that period times the product of the profile suprema. Abel summation against n^{-2-r-s} therefore bounds its tail by the same constant times K^{-2-r-s} . Multiplying by $2a$ and summing all orders, all support pairs and their absolute coefficient products pays at most $4aN^2b^2S_0^2/K^2$.

The exact weight obeys $|p_n - 2an^{-2a-1}| \leq a(2a+1)n^{-2a-2}$. Since $\sum_{n \geq K} n^{-3-r-s} \leq 2K^{-2-r-s}$, replacing this weight costs at most $2a(2a+1)b^2S_0^2/K^2$. Replacing the remaining power sum by $\int_K^\infty x^{-2-r-s} dx = K^{-1-r-s}/(r+s+1)$ costs at most $2ab^2S_0^2/K^2$. The two costs combine to $4a(a+1)b^2S_0^2/K^2$. The resulting main term is precisely (10), including every complex mixed-order contribution.

Finally Lemma 2.2 bounds the difference between the complete square and the finite-profile square by $(2be + e^2)S_0^2n^{-2\sigma}$. Its full p_n sum is at most $(2be + e^2)S_0^2/K$. This proves the two-sided enclosure. The matrix upper follows from $S_0^2 \leq N\|c\|_2^2$. $\square$

3.1 An exact divisor diagonalization

For $k \geq 2$, $1 \leq v < k$, $(v, k) = 1$, set

$$u_r(k, v) = -\frac{k^r}{(r+1)!} \sum_{j=0}^{k-1} B_{r+1}(j/k) e^{-2\pi i v j/k}.$$

Theorem 3.2 (Full mixed-order divisor weights). *On the anchored space,*

$$U_R(c) = \frac{2a}{K} \sum_{k=2}^N W_{R,k}(q, K) |t_k|^2, \quad (13)$$

where the strictly positive weights are

$$W_{R,k} = \int_0^1 \sum_{\substack{1 \leq v < k \\ (v,k)=1}} \left| \sum_{r=0}^R (q)_r K^{-r} x^r u_r(k, v) \right|^2 dx > 0. \quad (14)$$

For $R = 0$ they reduce to $J_2(k)/12$.

Proof. If $k \mid d$, Bernoulli multiplication gives the conditional mean

$$\mathbb{E}[H_{r,d}(n) \mid n \bmod k = j] = -\frac{k^{r+1}}{d(r+1)!} B_{r+1}(j/k).$$

The normalized Fourier coefficient at the reduced frequency v/k is thus $u_r(k, v)/d$; if $k \nmid d$ it is zero. This includes the exact $r = 0$ jump convention. Consequently the coefficient of $H_{r,c}$ is $u_r(k, v)t_k$. Orthogonality of distinct reduced rational frequencies on a common period proves (13). Every zero-frequency coefficient is proportional to t_1 and vanishes by anchoring. Since $u_0(k, v) = (1 - e^{-2\pi i v/k})^{-1} \neq 0$, the polynomial in x cannot vanish identically, proving strict positivity. Finally the classical cosecant-square sum gives $\sum_{(v,k)=1} |u_0(k, v)|^2 = J_2(k)/12$. $\square$

This is an all-order use of established Bernoulli/Farey covariance and divisor grouping. The quadratic Bernoulli wave and its divisor energy already appear in [4, Theorem 1]; the order-zero gcd form and the first correction are analyzed in Theorem 11.1 and Corollary 13.2. The mixed q -dependent receiver retains their complete interaction.

For exact real arithmetic, define the Ramanujan sum $c_k(j) = \sum_{h|\gcd(k,j)} h\mu(k/h)$ and $b_r(j) = -k^r B_{r+1}(j/k)/(r+1)!$. Then

$$F_{rs}(k) = \sum_{j,l=0}^{k-1} b_r(j)b_s(l)c_k(l-j), \quad W_{R,k} = \sum_{r,s=0}^R \frac{(q)_r \overline{(q)}_s}{K^{r+s}(r+s+1)} F_{rs}(k).$$

Thus the entire positive weight can be assembled from rational covariance data and enclosed complex rising factorials.

4 Height-proportional tail capacity and resonance

The finite covariance weights have a closed generating function. For reduced v/k ,

$$\sum_{r \geq 0} u_r(k, v) z^r = \frac{1}{1 - e^{z-2\pi i v/k}}, \quad |z| < 2\pi \min(v/k, 1 - v/k). \quad (15)$$

To prove this, sum the Bernoulli generating function $te^{xt}/(e^t - 1)$ over $x = j/k$ with the character $e^{-2\pi i v j/k}$, and put $t = kz$. The constant term vanishes because the character is nontrivial; the finite geometric sum cancels $e^{kz} - 1$. This also recovers the discrete value $u_0 = (1 - e^{-2\pi i v/k})^{-1}$.

Theorem 4.1 (Complete height-scale law). *Fix N and a . Suppose $T \rightarrow +\infty$, $K/T \rightarrow \kappa > N/(2\pi)$. Choose a fixed θ with $N/(2\pi\kappa) < \theta < 1$ and integers $R = O(\log(T+2))$ satisfying $\theta^R(|q|+2)^2 \leq 1/8$. Uniformly on the height-dependent double-null spaces with $\|c\|_2 \leq 1$,*

$$\frac{K}{2a}(Q - Q_M) - \sum_{k=2}^N W_k(\kappa) |t_k|^2 \longrightarrow 0, \quad (16)$$

where

$$\begin{aligned} W_k(\kappa) &= \int_0^1 \sum_{\substack{1 \leq v < k \\ (v,k)=1}} |1 - e^{i(x/\kappa - 2\pi v/k)}|^{-2} dx \\ &= -\frac{\kappa}{2} \sum_{d|k} \mu(k/d) d \cot \frac{d}{2\kappa}. \end{aligned} \quad (17)$$

All arithmetic indices in the complete tail are included in this limit.

Proof. The wall (8) holds eventually. For fixed N , Theorem 3.1 gives $KN\delta_R \rightarrow 0$: its periodic part is $O(N^3/K)$, while $e_\theta = O(1/T)$. For each fixed order, $(q)_r/K^r \rightarrow (i/\kappa)^r$. The complete geometric bound from Lemma 2.2 is uniform, because $LN/(2\pi K)$ is eventually below the fixed $\theta < 1$. Dominated convergence in the finite-dimensional source and (15) prove (16), uniformly in its whole coefficient unit ball.

Now $|1 - e^{iy}|^{-2} = \frac{1}{4} \csc^2(y/2)$. Integrate each term to obtain a difference of cotangents. The starting cotangents sum to zero by opposite-residue pairing. Finally Möbius inversion of $\sum_{j=0}^{d-1} \cot(y - \pi j/d) = d \cot(dy)$ gives (17). The zero frequency $k = 1$ has already been removed by the anchor. $\square$

The theorem fixes N while height varies. For arbitrary moving support the quantitative recipient is the finite enclosure (12), with its displayed masses, wall and error; it is available before passing to any limit.

Corollary 4.2 (Higher-Jordan inflation). *For $k > 1$ and $\kappa > k/(2\pi)$,*

$$W_k(\kappa) = \sum_{\ell \geq 1} \frac{\zeta(2\ell) J_{2\ell}(k)}{2^{2\ell-1} \pi^{2\ell} \kappa^{2\ell-2}} = \frac{J_2(k)}{12} + \frac{J_4(k)}{720\kappa^2} + \frac{J_6(k)}{30240\kappa^4} + \cdots. \quad (18)$$

In particular $W_k(\kappa) > J_2(k)/12$. If also $\kappa > k/4$,

$$W_k(\kappa) \leq \left(1 - \frac{k}{4\kappa}\right)^{-2} \frac{J_2(k)}{12}.$$

At the first resonance,

$$W_k(\kappa) \sim \frac{k^3}{8\pi^3(\kappa - k/(2\pi))} \quad \text{as } \kappa \downarrow k/(2\pi). \quad (19)$$

Proof. Use the convergent expansion $\cot y = y^{-1} - 2 \sum_{\ell \geq 1} \zeta(2\ell) y^{2\ell-1} / \pi^{2\ell}$ in (17). The inverse-power term cancels because $\sum_{d|k} \mu(k/d) = 0$; the remaining divisor sums are exactly $J_{2\ell}(k) > 0$. For the upper bound, the triangle inequality gives

$$|1 - e^{i(x/\kappa - 2\pi v/k)}| \geq |1 - e^{-2\pi i v/k}| - 1/\kappa,$$

while $|1 - e^{-2\pi i v/k}| \geq 2 \sin(\pi/k) \geq 4/k$. Compare termwise with the order-zero sum. At the boundary in (19), only the $d = k$ cotangent has a pole. Expanding its argument at $\kappa = k/(2\pi)$ gives the stated coefficient. $\square$

The old fixed-parameter gcd coefficient is thus strictly smaller than the height-scale leading coefficient at every finite pole-free κ . Its original fixed-parameter theorem retains its remainder; the mixed receiver pays this additional contribution when height varies. Cotangent/Nyman–Beurling connections and Vasyunin sums have classical antecedents [15]; the present law is for the complete native tail with simultaneous height and wall variation.

4.1 Capacity in the resummed metric

For the finite weights $W_{R,k} > 0$, put $j_z(k) = J_{1-z}(k)$ and

$$\mathcal{D}_R = \sum_{k=2}^N \frac{|j_\lambda(k)|^2}{W_{R,k}} - \frac{\left| \sum_{k=2}^N j_\lambda(k) \overline{j_q(k)} / W_{R,k} \right|^2}{\sum_{k=2}^N |j_q(k)|^2 / W_{R,k}}.$$

When the constraints are feasible and $\mathcal{D}_R > 0$, exact weighted projection gives

$$\inf_{\substack{A(1)=A(q)=0 \\ A(\lambda)=1}} U_R(c) = \frac{2a}{K\mathcal{D}_R}.$$

Indeed (6) expresses the two rows in the diagonal W metric; project the λ row off the q row and apply Cauchy–Schwarz with equality at its representer. The q row is nonzero for $N \geq 2$, since $J_{1-q}(2) = 2^{1-q} - 1 \neq 0$.

Define $U_\infty(c) = (2a/K) \sum_{k=2}^N W_k(\kappa) |t_k|^2$. For $\kappa \geq N$, Corollary 4.2 gives

$$\frac{2a}{K} V(c) \leq U_\infty(c) \leq \frac{16}{9} \frac{2a}{K} V(c).$$

The corresponding constrained resummed capacities lie between the old variance capacity and 16/9 times it. Near resonance, the explicitly growing weights instead identify the expensive divisor coordinates. The complete gain receiver retains its arithmetic head and error as described next.

5 The variance-priced growing-support comparison

Throughout this section $N \geq 2$ is an integer, and coefficients have the native support $1, \dots, N$. The preceding height-scale theorem fixes the support. We now give a quantitative comparison on varying support and height, paying the inverse divisor-coordinate cost explicitly. Write $\sigma_1(k) = \sum_{d|k} d$ and

$$w_0(k) = \frac{J_2(k)}{12}, \quad V(c) = \sum_{k=2}^N w_0(k) |t_k|^2, \quad C_N = 12 \sum_{k=2}^N \frac{\sigma_1(k)^2}{J_2(k)}.$$

These quantities use the same t_k and original coefficients as the full native recipient.

Lemma 5.1 (Complete coefficient price). *For every anchored vector,*

$$S_0^2 \leq C_N V(c), \quad C_N \leq \frac{897}{896} \frac{\pi^6}{18} N. \quad (20)$$

Proof. Möbius inversion gives $c_d = d \sum_{d|k} \mu(k/d) t_k$. Since $t_1 = 0$, this implies $S_0 \leq \sum_{k=2}^N \sigma_1(k) |t_k|$; weighted Cauchy–Schwarz proves the first inequality, including the coefficient c_1 .

For the second, put $f(k) = \sigma_1(k)^2 / J_2(k)$, $f(1) = 1$, and $g = f * \mu$. Both are multiplicative. If $x = 1/p$ and $j \geq 1$, then

$$f(p^j) = \frac{(1 - x^{j+1})^2}{(1 - x)^2(1 - x^2)}.$$

These values increase, including $f(p) > f(1)$, so $g \geq 0$. The local sum is

$$\begin{aligned} B_p &= \sum_{j \geq 0} g(p^j) p^{-j} = (1 - x) \sum_{j \geq 0} f(p^j) x^j \\ &= \frac{1 + x^2}{(1 - x)^3(1 + x)^2(1 + x + x^2)} - \frac{x^2}{1 + x}. \end{aligned}$$

Direct algebra gives

$$B_p(1 - x^2)^3 = 1 + \frac{x^3}{1 + x + x^2} - x^2(1 - x)(1 - x^2)^2.$$

For $p \geq 3$, $x \leq 1/3$ and

$$\frac{x}{1+x+x^2} \leq \frac{1}{3} < \frac{128}{243} \leq (1-x)(1-x^2)^2,$$

so $B_p \leq (1-p^{-2})^{-3}$. At $p = 2$, $B_2 = 299/126 = (897/896)(1-1/4)^{-3}$. Nonnegative partial Euler products are therefore bounded by $(897/896)\zeta(2)^3$. Consequently

$$\sum_{k \leq N} f(k) = \sum_{d \leq N} g(d) \lfloor N/d \rfloor \leq N \sum_{d \geq 1} \frac{g(d)}{d} \leq \frac{897}{896} \zeta(2)^3 N.$$

Multiplying by 12 proves (20). Positivity also justifies the series and product limits used here. $\square$

Lemma 5.2 (Comparison throughout the pole-free range). *For every $k > 1$ and $\kappa > k/(2\pi)$,*

$$w_0(k) \leq W_k(\kappa) \leq \left(1 - \frac{k}{2\pi\kappa}\right)^{-2} w_0(k).$$

Proof. The lower bound is the positive Jordan series. For the upper, put $y = \pi v/k$ and $h = x/(2\kappa)$. Then $0 \leq h < \pi/k \leq y < \pi$. Concavity of sine on $[0, \pi]$ gives

$$\sin(y-h) \geq (1-h/y) \sin y \geq \left(1 - \frac{k}{2\pi\kappa}\right) \sin y.$$

Compare the squared reciprocal denominators in (17), integrate in x , and sum over the reduced residues. $\square$

Theorem 5.3 (Finite relative cotangent comparison). *Let $T \geq 1$, $R \geq 1$ and $0 < \theta < 1$, and require the declared wall (8). Use δ_R from (11) with this same θ . Put*

$$\begin{aligned} L &= |q| + R + 1, & \kappa &= K/T, & Z &= \left(1 - \frac{NT}{2\pi K}\right)^{-1}, \\ \eta_R &= 2\sqrt{2} \left(\frac{\theta}{L(1-\theta)^3} + \frac{\theta^{R+1}}{1-\theta} \right), & \mathcal{H} &= \eta_R(2Z + \eta_R) + \frac{K\delta_R C_N}{2a}. \end{aligned}$$

Then, uniformly over the actual exact double-null coefficient space,

$$\left| \frac{K}{2a} (Q - Q_M) - \sum_{k=2}^N W_k(K/T) |t_k|^2 \right| \leq \mathcal{H} V(c). \quad (21)$$

Every infinite arithmetic tail and mixed primitive order in this comparison is paid.

Proof. The wall implies $NT/(2\pi K) < 1$, so every cotangent weight and Z are finite. For $r \geq 1$, discrete Parseval applied to the unnormalized Fourier transform defining u_r gives

$$\left(\sum_{(v,k)=1} |u_r(k, v)|^2 \right)^{1/2} \leq \frac{4k^{r+1}}{(2\pi)^{r+1}} \leq 2\sqrt{2} \left(\frac{k}{2\pi} \right)^r \sqrt{w_0(k)}.$$

Indeed the full Fourier sum is k times the sum of squares of the k samples, each bounded by $4k^r/(2\pi)^{r+1}$; restriction to reduced residues decreases it. The second inequality uses $J_2(k) \geq 6k^2/\pi^2$.

For $1 \leq r \leq R$, telescope the products $(q)_r$ and $(iT)^r$. Every individual factor has modulus at most L , while its difference from iT is $\sigma + j$. Consequently

$$|(q)_r - (iT)^r| \leq L^{r-1} \sum_{j=0}^{r-1} (\sigma + j) \leq \frac{r(r+1)}{2} L^{r-1}.$$

In the Hilbert space $L^2([0, 1])$ with reduced-residue coordinates, compare the finite profile $\sum_{r=0}^R (q)_r K^{-r} x^r u_r(k, v)$ with its exact resummation $(1 - e^{iT x/K - 2\pi i v/k})^{-1}$. The $r = 0$ terms agree, including the original jump. Using $kL/(2\pi K) \leq \theta$ and the complete identity

$$\sum_{r \geq 1} \frac{r(r+1)}{2} \theta^r = \frac{\theta}{(1-\theta)^3},$$

the finite coefficient discrepancy has norm at most $2\sqrt{2}\theta\sqrt{w_0(k)}/[L(1-\theta)^3]$. All omitted $r > R$ terms together cost at most $2\sqrt{2}\theta^{R+1}\sqrt{w_0(k)}/(1-\theta)$. The generating function (15) is used strictly inside its pole-free disk, so the total norm distance is at most $\eta_R\sqrt{w_0(k)}$.

Lemma 5.2 bounds the resummed norm by $Z\sqrt{w_0(k)}$. The difference of the squared norms is therefore at most $\eta_R(2Z + \eta_R)w_0(k)$. Sum against $|t_k|^2$ and add the complete scaled error $K\delta_R S_0^2/(2a)$ from Theorem 3.1. Lemma 5.1 proves (21). $\square$

Corollary 5.4 (A growing-support height law). *Fix $a \in (0, 1/2)$, $\gamma > 1/(2\pi)$ and $1/(2\pi\gamma) < \theta < 1$. As $T \rightarrow \infty$, choose*

$$K = \lceil \gamma NT \rceil, \quad R = \left\lceil \frac{3 \log T}{\log(1/\theta)} \right\rceil.$$

For all sufficiently large T , $R \geq 1$, the wall is admitted uniformly in N , and

$$\mathcal{H} = O_{a,\gamma,\theta} \left(\frac{N^2}{T} + \frac{1}{T} \right).$$

In particular $N = o(\sqrt{T})$ gives (21) with error tending to zero uniformly after normalization $V(c) = 1$. Both null spaces and their coefficient vectors may vary with height.

Proof. Here $R = O_\theta(\log T)$ and $|q|/T \rightarrow 1$, so the fixed strict inequality $2\pi\theta\gamma > 1$ admits the wall eventually, independently of N . Also $\theta^R \leq T^{-3}$, hence $e_\theta = O(T^{-2})$, $C_N = O(N)$, $\eta_R = O_\theta(T^{-1})$, and $Z \leq (1 - 1/(2\pi\gamma))^{-1}$. The displayed complete δ_R gives $K\delta_R C_N/(2a) = O_{a,\gamma,\theta}(N^2/T + N/T^2)$. Since $N \geq 2$, these estimates prove the assertion. $\square$

5.1 The variance-priced matrix option

Let C be the inverse divisor map $c = Ct$, restricted to $t_1 = 0$. Another complete positive upper for the original norm is

$$C^* G_M C + \frac{2a}{K} \text{diag}(W_{R,k}) + \delta_R C_N \text{diag}(w_0(k)).$$

The same exterior-null row remains imposed. This follows directly from $S_0^2 \leq C_N V(c)$ and is an alternative whole quadratic envelope to the earlier coefficient-metric price. The corrected mean is still evaluated on the actual $c = Ct$; its error is not transferred to an unpaid identity metric. The full-head applications below use their stated complete envelope.

Since $W_k \geq w_0(k)$, an error $\mathcal{H} < 1$ also gives a relative $(1 \pm \mathcal{H})$ enclosure around the cotangent tail. Adding the positive unchanged head preserves the same multiplicative enclosure for the full

norm. On any nonempty family with additional exact linear constraints, taking infima preserves these bounds; no conditioning assumption on its evaluation rows is needed for that comparison. This supplies uniform moving-support norm control; a favorable corrected numerator and full-head gain still require their own certificates.

6 A complete Farey-operator return

The variance-priced comparison above bounds each periodic product separately. Applying the classical absolute Farey block inequality before squaring the varying primitive profile improves the moving-support regime. The absolute Hilbert and large-sieve input is Montgomery–Vaughan [17, Section 2]; the same sine-quotient argument is explained in [18, Section 17.3]. An unshifted Nyman–Beurling wavelet tail sandwich using primitive Farey frequencies, the absolute block estimate and Abel summation appears in [19, Theorem 4]. The return here retains the height-dependent rising factors, their varying amplitudes, the exact Poisson weight and every mixed primitive order of the original double-null recipient.

Theorem 6.1 (Complete fixed-gap pole-free Farey receiver). *Let $0 < a < 1/2$, $T \geq 1$, $N \geq 2$, $R \geq 1$ and $K = M + 1 \geq 2$. Require both exact nulls $A(1) = A(q) = 0$, fix $0 < \theta < 1$, and impose the declared wall $K \geq NL/(2\pi\theta)$, where $L = |q| + R + 1$. Use the same b_θ, e_θ as in (9), and put*

$$\begin{aligned} B_0 &= 1 + \frac{2\sqrt{2}\theta}{1-\theta}, & B_1 &= \frac{2\sqrt{2}\theta}{(1-\theta)^2}, & A_\theta &= B_0^2 + B_0 B_1, \\ \Delta_\theta &= \frac{2a(N^2 + 1)A_\theta + a(2a + 1)B_0^2}{K^2} + \frac{(2b_\theta e_\theta + e_\theta^2)C_N}{K}, \\ \eta_\theta &= 2\sqrt{2} \left(\frac{\theta}{L(1-\theta)^3} + \frac{\theta^{R+1}}{1-\theta} \right), & Z &= \left(1 - \frac{NT}{2\pi K} \right)^{-1}, \\ H_\theta &= \eta_\theta(2Z + \eta_\theta) + \frac{K\Delta_\theta}{2a}. \end{aligned}$$

For the full mixed form U_R of (10) and the actual cotangent form $W(c) = \sum_{k=2}^N W_k(K/T)|t_k|^2$, one has

$$|Q - Q_M - U_R(c)| \leq \Delta_\theta V(c), \quad (22)$$

$$\left| \frac{K}{2a}(Q - Q_M) - W(c) \right| \leq H_\theta V(c). \quad (23)$$

These finite complete-tail statements do not require $H_\theta < 1$.

Proof. Index the distinct nonzero reduced Farey fractions of denominators at most N by $\omega = v/k$. Write A_r for the vector with entries $u_r(k, v)t_k$. The exact discrete Fourier representation of the finite primitive profile is

$$f(n) = \sum_{\omega} b_{\omega}(n) e^{2\pi i \omega n}, \quad b(x) = \sum_{r=0}^R (q)_r x^{-r} A_r.$$

Every order has zero constant Fourier component because $t_1 = 0$. The right-jump value at order zero is retained in u_0 . Discrete Parseval, as in the proof of Theorem 5.3, gives

$$\|A_0\| = \sqrt{V(c)}, \quad \|A_r\| \leq 2\sqrt{2} \left(\frac{N}{2\pi} \right)^r \sqrt{V(c)} \quad (r \geq 1).$$

Since $|(q)_r| \leq L^r$, the declared wall gives, for every $x \geq K$,

$$\|b(x)\| \leq B_0\sqrt{V}, \quad \|b'(x)\| \leq B_1\sqrt{V}/x. \quad (24)$$

Distinct reduced fractions have circular separation at least N^{-2} . Let E_n have entries $e^{2\pi i(\omega_j - \omega_i)n}$ and put $D_{K,J} = \sum_{n=K}^J (E_n - I)$. The absolute cosecant Hilbert inequality gives

$$\|D_{K,J}\|_{\text{op}} \leq N^2 \quad (J \geq K). \quad (25)$$

For completeness, each off-diagonal geometric sum is a diagonal unitary conjugate of a sine quotient with denominator $\sin \pi(\omega_j - \omega_i)$. Splitting its sine numerator into two exponentials expresses the matrix as the difference of two unitary conjugates of the cosecant kernel, divided by $2i$. Apply the absolute Hilbert bound to both terms. This proves both signs of the Hermitian operator estimate; a one-sided upper large sieve alone is not used as a two-sided estimate.

Apply matrix Abel summation to the exact original weight

$$w(x) = x^{-2\sigma} [x^{-2a} - (x+1)^{-2a}] = 2ax^{-2\sigma} \int_0^1 (x+t)^{-2a-1} dt.$$

It satisfies $0 < w(x) \leq 2ax^{-2}$ and $|w'(x)| \leq 4ax^{-3}$; the derivative uses $2\sigma + 2a = 1$. For $F(x) = w(x)b(x)b(x)^*$, (24) gives, whenever $\|D\| \leq N^2$,

$$|\text{Tr}(DF'(x))| \leq N^2(|w'|\|b\|^2 + 2w\|b'\|\|b\|) \leq 4aN^2A_\theta Vx^{-3}.$$

The Abel sum starts with $D_{K,K-1} = 0$; its boundary at infinity vanishes by these bounds. Integration over the entire half-line gives

$$\left| \sum_{n \geq K} w(n)(|f(n)|^2 - \|b(n)\|^2) \right| \leq \frac{2aN^2A_\theta V}{K^2}.$$

This pays variation of every amplitude, rather than freezing it at K .

Replacing the diagonal weight by $2an^{-2}$ costs at most $a(2a+1)B_0^2V/K^2$: use $|w(n) - 2an^{-2}| \leq a(2a+1)n^{-3}$ and $\sum_{n \geq K} n^{-3} \leq K^{-2}$ for $K \geq 2$. The complete sum/integral replacement for $h(x) = 2ax^{-2}\|b(x)\|^2$ costs at most $2aA_\theta V/K^2$, because $|h'(x)| \leq 4aA_\theta Vx^{-3}$. Its integral is exactly $U_R(c)$.

Finally the full primitive remainder and its cross term are bounded by $(2b_\theta e_\theta + e_\theta^2)S_0^2/K$. Here Lemma 2.2 uses the same general θ , and $\sum_{n \geq K} p_n n^{-2\sigma} \leq K^{-1}$. Apply $S_0^2 \leq C_N V$ and add all payments to prove (22). The constants from a half-ratio wall cannot be retained unchanged when the declared ratio is enlarged.

The finite-profile/cotangent comparison in Theorem 5.3 gives the squared-norm error $\eta_\theta(2Z + \eta_\theta)V$. It retains all omitted orders and uses Lemma 5.2 throughout the pole-free range. Add the scaled error $K\Delta_\theta V/(2a)$ to obtain (23). $\square$

Corollary 6.2 (Near-linear growing support). *Fix $a \in (0, 1/2)$ and $\gamma > 1/(2\pi)$. Set*

$$\theta = \frac{1 + 1/(2\pi\gamma)}{2}, \quad K = \lceil \gamma NT \rceil, \quad R = \max \left(1, \left\lceil \frac{3 \log T}{\log(1/\theta)} \right\rceil \right).$$

Eventually the wall is admitted uniformly in N and

$$H_\theta = O_{a,\gamma}(N/T + 1/T).$$

Thus $N = o(T)$ gives a uniform complete cotangent tail law on the actual varying double-null spaces after normalization $V = 1$. The constants need not remain bounded as the fixed gap $\gamma - 1/(2\pi)$ shrinks.

Proof. Here $L/T \rightarrow 1$, $\theta^R \leq T^{-3}$, $e_\theta = O(T^{-2})$, $\eta_\theta = O(T^{-1})$ and $Z = O_\gamma(1)$. Since $C_N = O(N)$ and $K \asymp_\gamma NT$, $K\Delta_\theta = O_{a,\gamma}(N/T + N/T^2)$. This proves the asserted bound and its sufficient regime. $\square$

Since $W \geq V$, whenever $H_\theta < 1$ the model $\hat{Q} = Q_M + (2a/K)W$ satisfies

$$(1 - H_\theta)\hat{Q} \leq Q \leq (1 + H_\theta)\hat{Q}.$$

The same relative bounds hold after taking infima over any nonempty exact-null affine coefficient family, even if its infimum is zero. A complete finite mixed upper in inverse divisor coordinates $c = Ct$ is

$$C^*G_M C + \frac{2a}{K} \text{diag}(W_{R,k}) + \Delta_\theta \text{diag}(J_2(k)/12),$$

with $t_1 = 0$ and the original exterior-null row imposed. The complete corrected mean and its error use the same C and actual Ct . This is a full receiving matrix, not merely a tail capacity. It does not itself assert favorable numerator size or positive all-height gain.

7 Exact evaluation duals

Write μ for the Möbius function and

$$M_x(z) = \sum_{m \leq x} \mu(m) m^{-z}.$$

For $1 \leq n \leq N$ set

$$v_n = n^{-2a} M_{N/n}(\lambda), \quad v_{N+1} = 0, \quad \ell_n = v_n - v_{n+1}, \quad \mathcal{D}_N = \sum_{n \leq N} \frac{|\ell_n|^2}{p_n}. \quad (26)$$

Theorem 7.1 (Evaluation-first dual). *For every coefficient vector supported at most N ,*

$$A(\lambda) = \sum_{n \leq N} \ell_n B_n.$$

The squared norm of this functional on the unrestricted first-head space with norm $\sum_{n \leq N} p_n |B_n|^2$ is exactly $\mathcal{D}_N$. For the anchored complete recipient,

$$g_N(\lambda) \leq \min\{1, |\zeta(\lambda)|^2 \mathcal{D}_N\}. \quad (27)$$

Proof. Finite inversion gives

$$c_k = \sum_{d|k} \mu(k/d) d^q (B_d - B_{d-1}), \quad B_0 = 0.$$

Substitution in $\sum c_k k^{-\lambda}$ yields $\sum_{d \leq N} d^{q-\lambda} M_{N/d}(\lambda) (B_d - B_{d-1})$. Since $q - \lambda = -2a$, summation by parts gives the stated row. Weighted Cauchy–Schwarz proves the norm bound, with equality in the unrestricted head at $B_n = \overline{\ell_n}/p_n$. The full Q is at least its positive first head, so (3) proves (27). $\square$

Anchoring can be retained exactly, rather than discarded in this upper bound. Put

$$w_n = n^{q-1} M_{N/n}(1), \quad w_{N+1} = 0, \quad h_n = w_n - w_{n+1}.$$

The same inversion gives $A(1) = \sum h_n B_n$. In particular $h \neq 0$, since $h_N = N^{q-1}$. The exact squared head norm restricted to anchoring is

$$\mathcal{D}_N^{\text{anch}} = \mathcal{D}_N - \frac{\left| \sum_{n \leq N} \ell_n \overline{h_n} / p_n \right|^2}{\sum_{n \leq N} |h_n|^2 / p_n}. \quad (28)$$

This follows by orthogonal projection after setting $x_n = \sqrt{p_n} B_n$. Thus $\mathcal{D}_N$ in (27) may be replaced by $\mathcal{D}_N^{\text{anch}}$. Neither head norm is asserted equal to the optimized gain in the complete arithmetic recipient.

8 A log-free power law and a necessary cutoff cost

Let $\delta = 1 - \alpha = \sigma$ and $b = 1 - 2a = 2\delta$. In the following lemma K, L are dummy integer cutoffs. For $K \geq 1$ define a finite sequence $H_K(n) = n^{-2a} \mathbf{1}_{n \leq K}$ and its forward difference $\Delta H_K(n) = H_K(n) - H_K(n+1)$.

Lemma 8.1 (Rank-one plus diagonal Gram). *In the difference norm with weights $1/p_n$,*

$$\langle \Delta H_K, \Delta H_L \rangle_{1/p} = 1 \quad (K \neq L), \quad d_K := \|\Delta H_K\|_{1/p}^2 = 1 - K^{-2a} + \frac{K^{-4a}}{p_K}.$$

Writing

$$u_K = \sum_{\lfloor N/m \rfloor = K} \mu(m) m^{-\lambda},$$

one consequently has the exact identity

$$\mathcal{D}_N = |M_N(\lambda)|^2 + \sum_{K \leq N} (d_K - 1) |u_K|^2. \quad (29)$$

Proof. The difference ΔH_K equals p_n for $n < K$, equals K^{-2a} at $n = K$, and vanishes after that. For $K < L$ the cross product is $\sum_{n < K} p_n + K^{-2a} = 1$. The diagonal formula follows similarly. Now $v = \sum_{m \leq N} \mu(m) m^{-\lambda} H_{\lfloor N/m \rfloor}$. Expand its difference norm and group equal floor values. The all-ones part is $|\sum u_K|^2 = |M_N(\lambda)|^2$. $\square$

Theorem 8.2 (Uniform power law). *For every $T \in \mathbb{R}$ and integer $N \geq 1$,*

$$N^b \left(\frac{1}{2a} - \frac{1}{N} \right) \leq \mathcal{D}_N \leq K_a N^b, \quad (30)$$

where one admissible explicit constant is

$$K_a = \delta^{-2} + \frac{2^{2a}}{a} (2^{2\alpha+2} + 1) \zeta(2).$$

In particular $g_N(\lambda) \leq K_a N^{2-2\alpha} |\zeta(\lambda)|^2$. The exponent in (30) is sharp for $\mathcal{D}_N$.

Proof. The integral formula for p_K gives

$$a 2^{-2a} K^{-2a-1} \leq p_K \leq 2a K^{-2a-1}, \quad 0 < d_K - 1 \leq \frac{2^{2a}}{a} K^b.$$

Also $|M_N(\lambda)| \leq \sum_{m \leq N} m^{-\alpha} \leq N^\delta / \delta$. If $K \leq \sqrt{N}$, the interval $N/(K+1) < m \leq N/K$ contains at most $2N/K^2$ integers. Hence

$$|u_K| \leq 2^{\alpha+1} N^\delta K^{\alpha-2}, \quad K^b |u_K|^2 \leq 2^{2\alpha+2} N^b / K^2.$$

If $K > \sqrt{N}$, that interval has length less than one. Each nonempty group contains just one m , and its contribution is at most N^b/m^2 . Summing both regimes proves the upper bound via (29). This is the weighted quotient-increment split underlying the related harmonic estimates of [1].

All terms in (29) are nonnegative. The group $K = N$ contains only $m = 1$, so $u_N = 1$ and

$$\mathcal{D}_N \geq d_N - 1 = \frac{N^{-4a}}{p_N} - N^{-2a} \geq N^b \left(\frac{1}{2a} - \frac{1}{N} \right).$$

This sharpness statement concerns the unrestricted head functional; anchoring and the full tail remain present in the gain inequality. $\square$

Corollary 8.3 (All-height gain requires growing support). *Fix $\alpha \in (1/2, 1)$ and $\eta > 0$. If an arbitrary integer schedule $N(t) \geq 2$ satisfies $g_{N(t)}(\alpha + it) \geq \eta$ for all sufficiently large t , then along an unbounded sequence t_j ,*

$$\log N(t_j) \geq \frac{\sqrt{\alpha - 1/2}}{20(1 - \alpha)} \frac{(\log t_j)^{1-\alpha}}{(\log \log t_j)^\alpha} + O_{\alpha, \eta}(1). \quad (31)$$

Thus every cutoff with logarithm of smaller order than the displayed height scale fails a fixed positive gain requirement infinitely often.

Proof. Montgomery's unconditional Theorem 1 [9], at angle π , gives for every sufficiently large X a height $X^{(\alpha-1/2)/3} \leq t \leq X$ such that

$$|\zeta(\alpha + it)| \leq \exp \left[-\frac{\sqrt{\alpha - 1/2}}{20} \frac{(\log X)^{1-\alpha}}{(\log \log X)^\alpha} \right].$$

The scale is eventually increasing, so it can be replaced by its value at t . These heights are unbounded. Apply Theorem 8.2 and take logarithms of $\eta \leq K_a N(t)^b |\zeta(\alpha + it)|^2$. Since $b = 2(1 - \alpha)$, this is (31). $\square$

The source's sufficiently-large onset is not replaced by a numerical height here. The statement fixes α , permits all complex coefficient choices at every height, and concerns the largest permitted integer index rather than coefficient cardinality.

9 Null adaptation without loss of the closed trial space

The support lower cost does not construct a successful trial. We next give a complete way to certify any proposed growing trial.

Theorem 9.1 (Costed exterior-null adaptation). *For nonzero $A \in \mathcal{A}_N$ and integer $k \geq 1$, set*

$$\varepsilon = 2^{-ka}, \quad A_k(s) = A(s)(1 - 2^{k(q-s)}).$$

Then $A_k(1) = A_k(q) = 0$ and its support is at most $2^k N$. Moreover

$$Q(A_k - A) = \varepsilon^2 Q(A), \quad (32)$$

$$(1 - \varepsilon)^2 Q(A) \leq Q(A_k) \leq (1 + \varepsilon)^2 Q(A), \quad (33)$$

$$m(A_k) = (1 - \varepsilon^2)m(A), \quad (34)$$

$$(1 - \varepsilon)^2 g(A) \leq g(A_k) \leq (1 + \varepsilon)^2 g(A). \quad (35)$$

The arithmetic images satisfying both nulls have the same closed span as all anchored arithmetic images, at every fixed a, T .

Proof. On $\Re s = 1/2$, the multiplier $2^{k(q-s)}$ has constant modulus ε . Equation (4) proves (32); the triangle inequality proves the two norm bounds. At λ its value is $2^{-2ka} = \varepsilon^2$, proving the mean identity and the gain bounds. The two nulls and support claim are immediate, with any coincident integer coefficients added. As $k \rightarrow \infty$, the full-norm error tends to zero, proving equality of closures. The reverse inclusion holds because the doubly null polynomials are already anchored. $\square$

The transformed coefficient mass obeys

$$S_0(A_k) \leq (1 + 2^{k\sigma})S_0(A). \quad (36)$$

Thus small gain loss is accompanied by an explicit cost both in support and in coefficient mass; the construction is not cost-free as $a \rightarrow 0$.

10 Complete finite arithmetic lower certificates

Theorem 10.1 (Tail-paid lower gain). *Suppose $A \neq 0$ and $A(1) = A(q) = 0$, and put*

$$\mathcal{K} = \left(1 + \frac{|q|}{\sigma}\right) S_0(A), \quad m_M = \sum_{n \leq M} p_n B_n, \quad Q_M = \sum_{n \leq M} p_n |B_n|^2.$$

Then for every integer $M \geq 1$,

$$0 \leq Q - Q_M \leq \frac{\mathcal{K}^2}{M+1}, \quad (37)$$

$$|m - m_M| \leq \frac{\mathcal{K}}{(M+1)^\alpha}. \quad (38)$$

Consequently the finite arithmetic quantity

$$L_M(A) = \frac{(\max\{0, |m_M| - \mathcal{K}/(M+1)^\alpha\})^2}{Q_M + \mathcal{K}^2/(M+1)} \quad (39)$$

satisfies $0 \leq L_M(A) \leq g(A)$ and tends to $g(A)$ as $M \rightarrow \infty$. No zeta value is needed to evaluate (39).

Proof. Partial summation and boundedness of D give

$$\lim_{n \rightarrow \infty} B_n = q \int_1^\infty D(x) x^{-q-1} dx = \zeta(q) A(q) = 0.$$

Subtracting this limit yields

$$B_n = D(n) n^{-q} - q \int_n^\infty D(x) x^{-q-1} dx, \quad |B_n| \leq \mathcal{K} n^{-\sigma}.$$

Since $\sum_{n>M} p_n = (M+1)^{-2a}$, positivity and $2a + 2\sigma = 1$, $2a + \sigma = \alpha$ prove both tail bounds. The triangle inequality for m and upper bound for Q prove the certificate; convergence follows from the vanishing tails and $Q > 0$. $\square$

Combining Theorems 9.1 and 10.1 gives a complete finite hierarchy at a fixed center. To approach the limiting optimized gain, first choose a finite anchored A near its supremum, then k to retain its gain, then M to approximate $g(A_k)$. For a monotone sequence of lower bounds, retain the largest certificate obtained so far. Completeness is an exhaustion statement; a successful uniform schedule must additionally bound its support, mass and tail parameters over the claimed height range.

As a literal example,

$$A(s) = (1 - 2^{1-s})(1 - 2^{q-s})$$

uses only 1, 2, 4 and satisfies both nulls. Its center value is $(1 - 2^{1-\lambda})(1 - 2^{-2a}) \neq 0$ because $|2^{1-\lambda}| = 2^{1-\alpha} > 1$. It demonstrates the finite construction; its fixed cutoff remains subject to Corollary 8.3.

11 The sharp complete tail and a period-free remainder

The elementary tail upper in (37) can be refined to its leading coefficient. For a finite anchored coefficient vector put

$$V(c) = \frac{1}{12} \sum_{d,e} c_d \bar{c}_e \frac{\gcd(d,e)^2}{de}, \quad S_1 = \sum_d d |c_d|, \quad H = \frac{S_0}{2}. \quad (40)$$

Let $J_2(k) = k^2 \prod_{p|k} (1 - p^{-2})$ be the second Jordan function.

Theorem 11.1 (Sharp null-adapted tail). *For every nonzero finite A with $A(1) = A(q) = 0$, $V(c) > 0$. Set*

$$E = \frac{|q|S_1}{12} \left(1 + \frac{|q+1|}{\sigma+1} \right), \quad C_0 = 2a(V(c) + S_1^2) + \frac{a(2a+1)H^2}{2}.$$

Then, for every integer $M \geq 1$,

$$\left| \sum_{n>M} p_n |B_n|^2 - \frac{2aV(c)}{M} \right| \leq \frac{C_0}{M^2} + \frac{2HE}{(M+1)^2} + \frac{E^2}{(M+1)^3}. \quad (41)$$

In particular the M^{-1} order is exact for every fixed nonzero trial.

Proof. The continuous period mean of $D(x)$ is $\bar{D} = -\frac{1}{2} \sum c_d$. Define

$$h(x) = D(x) - \bar{D} = \sum_d c_d (1/2 - \{x/d\}), \quad P(x) = -\frac{1}{2} \sum_d c_d dB_2(\{x/d\}),$$

where $B_2(u) = u^2 - u + 1/6$. The periodic continuous primitive P satisfies $P' = h$ almost everywhere, $|h| \leq H$ and $|P| \leq S_1/12$. The zero limit of B_n and integration by parts give

$$B_n = h(n)n^{-q} + qP(n)n^{-q-1} - q(q+1) \int_n^\infty P(x)x^{-q-2} dx.$$

Thus $B_n = h(n)n^{-q} + R_n$ with

$$|R_n| \leq En^{-\sigma-1}. \quad (42)$$

We require the discrete rather than continuous fractional-part covariance. Write $f_d(n) = 1/2 - \{n/d\}$ and $r = \gcd(d, e)$. Conditional on $n \bmod r = z$, the residues modulo d and e are independent on their admissible classes by the Chinese remainder theorem. Their conditional means are $(r/d)f_r(z)$ and $(r/e)f_r(z)$. Since $r^{-1} \sum_{z=0}^{r-1} f_r(z)^2 = 1/12 + 1/(6r^2)$, the exact discrete period mean is

$$\text{av}_n f_d(n)f_e(n) = \frac{\gcd(d, e)^2}{12de} + \frac{1}{6de}. \quad (43)$$

The extra term cancels in the coefficient quadratic form by anchoring. Hence $\text{av} |h(n)|^2 = V(c)$. The familiar continuous gcd covariance [10] is thus used here with its necessary discrete correction. The divisor identity for J_2 gives

$$12V(c) = \sum_k J_2(k) \left| \sum_{k|d} \frac{c_d}{d} \right|^2. \quad (44)$$

For a largest index with $c_d \neq 0$, the corresponding term is positive, which proves strict positivity.

For each pair (d, e) , the centered sequence in (43) has mean zero, period $\text{lcm}(d, e)$, and terms of modulus at most $1/2$. Therefore, writing $z_n = |h(n)|^2 - V(c)$,

$$\left| \sum_{n \leq L} z_n \right| \leq \frac{1}{2} \sum_{d, e} |c_d c_e| \text{lcm}(d, e) \leq \frac{S_1^2}{2} \quad (L \geq 1).$$

Abel summation gives $|\sum_{n > M} z_n/n^2| \leq S_1^2/(M+1)^2$. Also

$$|p_n - 2an^{-2a-1}| \leq a(2a+1)n^{-2a-2}, \quad \frac{1}{M+1} \leq \sum_{n > M} n^{-2} \leq \frac{1}{M}.$$

It follows that

$$\left| \sum_{n > M} p_n |h(n)|^2 n^{-2\sigma} - \frac{2aV(c)}{M} \right| \leq \frac{C_0}{M^2}.$$

Finally (42) bounds the difference of the squared terms by $2HEN^{-2\sigma-1} + E^2n^{-2\sigma-2}$. Summing against the full p_n and using their telescoping tail proves (41). No common-period length appears in its constants. $\square$

The denominator of (39) may use the smaller of the proved upper tails $\mathcal{K}^2/(M+1)$ and $2aV/M$ plus the right-hand side of (41). Subtracting a known leading term can improve an enclosure, but additional zeros at q alone do not change the nonzero M^{-1} coefficient.

12 Exact all-support tail capacity

The coefficient in Theorem 11.1 itself has a sharp optimization independent of the full head norm. For complex z define $J_z(k) = \sum_{d|k} \mu(k/d)d^z$.

Theorem 12.1 (Sharp tail-variance capacity). *At every fixed center (1), put*

$$S_\infty = \frac{\zeta(2)\zeta(2\alpha)}{|\zeta(1+\lambda)|^2} - 1 > 0.$$

Then

$$\inf_{\substack{A \text{ finite, } A(1)=A(q)=0 \\ A(\lambda)=1}} V(c) = \frac{1}{12S_\infty}. \quad (45)$$

Every zeta factor displayed in S_∞ is in its absolutely convergent half-plane. The infimum is approached by explicit finite polynomials.

Proof. Put $x_d = c_d/d$ and $t_k = \sum_{k|d} x_d$. On support at most N the inverse is $x_d = \sum_{m \leq N/d} \mu(m) t_{dm}$, and

$$A(s) = \sum_{k \leq N} t_k J_{1-s}(k), \quad A(1) = t_1, \quad 12V(c) = \sum_{k \leq N} J_2(k) |t_k|^2. \quad (46)$$

Thus anchoring is $t_1 = 0$. On coordinates $2 \leq k \leq N$ define

$$u_k = \frac{J_{1-\lambda}(k)}{\sqrt{J_2(k)}}, \quad v_k = \frac{J_{1-q}(k)}{\sqrt{J_2(k)}}.$$

The exact finite capacity is

$$S_N = \sum_{k=2}^N |u_k|^2 - \frac{|\sum_{k=2}^N u_k \overline{v_k}|^2}{\sum_{k=2}^N |v_k|^2}. \quad (47)$$

The denominator is nonzero for $N \geq 2$ since $2^{1-q} \neq 1$. When $S_N > 0$, the exact minimum of V under both nulls and $A(\lambda) = 1$ is $1/(12S_N)$, by projecting the evaluation row off the null row in the coordinates $y_k = \sqrt{J_2(k)} t_k$. Writing $C_N = \sum u_k \overline{v_k}$, an explicit minimizer is

$$y_k = \frac{\overline{u_k} - \overline{v_k} \overline{C_N} / \sum |v_j|^2}{S_N}.$$

It satisfies $\sum v_k y_k = 0$, $\sum u_k y_k = 1$, and $\sum |y_k|^2 = 1/S_N$. Formula (46) then constructs the coefficients. If $S_N = 0$, these normalized constraints are infeasible rather than an occasion to invert a singular matrix.

For $z = 1 - \lambda = \sigma - iT$, the prime-power formulas give

$$\begin{aligned} 1 + \sum_{j \geq 1} \frac{|J_z(p^j)|^2}{J_2(p^j)} &= 1 + \frac{|p^z - 1|^2}{(p^2 - 1)(1 - p^{2\sigma-2})} \\ &= \frac{|1 - p^{z-2}|^2}{(1 - p^{-2})(1 - p^{2\sigma-2})}. \end{aligned}$$

All factors converge absolutely since $2 - 2\sigma = 2\alpha > 1$ and $2 - \sigma > 1$. Multiplicativity and the positive series imply

$$\sum_{k \geq 1} \frac{|J_{1-\lambda}(k)|^2}{J_2(k)} = \frac{\zeta(2)\zeta(2\alpha)}{|\zeta(1+\lambda)|^2}. \quad (48)$$

In particular $u \in \ell^2$ with squared norm $S_\infty > 0$ after omitting coordinate 1. Cauchy-Schwarz in (46) already proves $V \geq |A(\lambda)|^2/(12S_\infty)$ for every finite anchored polynomial, even without the q -null condition.

In contrast, $v \notin \ell^2$. At primes,

$$\frac{|J_{1-q}(p)|^2}{J_2(p)} \geq (1 - 2^{-\alpha})^2 p^{2\alpha-2} \geq \frac{(1 - 2^{-\alpha})^2}{p},$$

and the reciprocal primes have divergent sum. Thus $\|v_{2:N}\| \rightarrow \infty$. The normalized rows converge weakly to zero: each fixed finite head divided by this norm tends to zero, and Cauchy–Schwarz bounds its pairing with the remaining part of u by that part’s ℓ^2 norm. Consequently

$$\frac{|\sum_{k=2}^N u_k \overline{v_k}|}{\|v_{2:N}\|} \rightarrow 0, \quad S_N \rightarrow S_\infty.$$

For all sufficiently large N , $S_N > 0$ and the finite minimizers therefore approach the lower bound, proving (45). $\square$

Corollary 12.2 (Uniform finite-cutoff capacity). *Let*

$$L_\alpha = \frac{2 - 2^{-\alpha}}{(1 - 2^{-\alpha})^2}, \quad \mathfrak{C}_\alpha = \exp(L_\alpha/\alpha).$$

For every real T and every integer $N \geq 2$ with $\log N \geq 1/(2a)$,

$$0 \leq S_\infty - S_N \leq N^{-2a} \left[e\zeta(2)(1 + \log N)^4 + 4\mathfrak{C}_\alpha \zeta(2)^2 (1 + \log N)^8 \right]. \quad (49)$$

Moreover $S_\infty \geq (2^\sigma - 1)^2/3$ uniformly in T . Thus a cutoff making the right side smaller than this positive constant guarantees feasibility of the normalized finite constraints for every T .

Proof. Write $\tau(n)$ for the divisor function and $H_N = \sum_{n \leq N} 1/n$. The elementary estimates $|J_z(n)| \leq n^{\Re z} \tau(n)$ and $J_2(n) \geq n^2/\zeta(2)$ imply

$$\left| \sum_{k=2}^N u_k \overline{v_k} \right| \leq \zeta(2) \sum_{k \leq N} \frac{\tau(k)^2}{k} \leq \zeta(2) H_N^4.$$

Here $\tau(n)^2 \leq d_4(n)$ prime-power by prime-power, and the fourfold divisor convolution has harmonic sum at most H_N^4 .

For a lower bound on $\sum |v_k|^2$, put $F(n) = \prod_{p|n} (1 - p^{-\alpha})^2$. Then

$$|v_n|^2 \geq n^{-1+2a} F(n).$$

The nonnegative squarefree-supported multiplicative function r with $r(p) = (1 - p^{-\alpha})^{-2} - 1$ satisfies $1/F = r * 1$ and

$$\sum_{n \leq N} \frac{1}{F(n)} \leq N \prod_p \left(1 + \frac{r(p)}{p} \right) \leq N \mathfrak{C}_\alpha.$$

Indeed $r(p) \leq L_\alpha p^{-\alpha}$, and $\sum_p p^{-1-\alpha} \leq \zeta(1 + \alpha) - 1 \leq 1/\alpha$. Cauchy–Schwarz on $N/2 < n \leq N$, which contains at least $N/2$ integers and omits coordinate 1, now gives

$$\sum_{n=2}^N |v_n|^2 \geq N^{-1+2a} \sum_{N/2 < n \leq N} F(n) \geq \frac{N^{2a}}{4\mathfrak{C}_\alpha}.$$

This proves the second term of (49) for the projection penalty in (47).

For the omitted evaluation norm set $\varepsilon = 1/\log N \leq 2a$. Rankin’s elementary comparison and the divisor Dirichlet series give

$$\sum_{n > N} |u_n|^2 \leq \zeta(2) N^{-2a+\varepsilon} \zeta(1 + \varepsilon)^4 \leq e\zeta(2) N^{-2a} (1 + \log N)^4.$$

Adding this tail and the projection penalty proves the bound. Finally the $n = 2$ term of (48), with $J_2(2) = 3$, is at least $(2^\sigma - 1)^2/3$, proving the last assertion. $\square$

The extra null has no penalty in this limiting variance optimum. It still has its exact finite cost in (47). This theorem optimizes the leading tail variance, rather than the complete Poisson norm or the full gain: the minimizing coefficients may have large head norm, coefficient mass and remainder constants. Combining those quantities is the further arithmetic optimization required for a uniformly successful certificate.

13 Compensated mean tails and actual alignment

The elementary mean bound (38) pays magnitudes separately. The two nulls permit a matched kernel that keeps their cancellation. Write $B_j(u)$ for the Bernoulli polynomials and $(z)_j$ for rising factorials. For integers $r \geq 1$ define

$$S_r = \sum_d d^r |c_d|, \quad C_r = \frac{\max_{0 \leq u \leq 1} |B_{r+1}(u)|}{(r+1)!}, \quad \Delta_r = (\lambda)_r - (q)_r.$$

In particular $C_1 = 1/12$, $C_2 = \sqrt{3}/216$ and $C_3 = 1/720$. Bernoulli differentiation and multiplication used below are classical [5, Section 24.4].

Theorem 13.1 (Matched complete mean correction). *Suppose $A(1) = A(q) = 0$. Put $K = M + 1$, where $M \geq 1$ is an integer. For each finite integer $R \geq 1$,*

$$m - m_M = t_{R,K}c + \mathcal{E}_{R,K}(c), \tag{50}$$

$$t_{R,K}c = \sum_{r=1}^R \frac{\Delta_r K^{-\lambda-r}}{(r+1)!} \sum_d c_d d^r B_{r+1}(\{K/d\}), \tag{51}$$

$$|\mathcal{E}_{R,K}(c)| \leq C_R S_R K^{-\alpha-R} \mathcal{A}_R, \tag{52}$$

$$\mathcal{A}_R = \frac{|\Delta_{R+1}|}{\alpha + R} + \frac{2a|(q)_{R+1}|}{(\sigma + R)(\alpha + R)}. \tag{53}$$

Every statement is for the complete infinite tail.

Proof. Set $F_K(x) = x^{-\lambda} - K^{-2a}x^{-q}$ for $x \geq K$. The summation identity in Proposition 1.1 and the zero limit of B_M under $A(q) = 0$ give

$$m - m_M = \sum_{j \geq K} d_j F_K(j) = - \int_K^\infty D(x) F'_K(x) dx.$$

The Dirichlet tails converge since D is bounded and $\Re q > 0$. Crucially $F_K(K) = 0$, so there is no endpoint jump contribution, including when K is divisible by a support index. Also $F_K(\infty) = 0$, so the constant part $\overline{D}$ contributes zero. For the centered $h = D - \overline{D}$ used in the sharp norm-tail proof put

$$P_r(x) = - \sum_d \frac{c_d d^r}{(r+1)!} B_{r+1}(\{x/d\}).$$

These are continuous periodic functions with $P'_1 = h$ almost everywhere, $P'_r = P_{r-1}$ for $r \geq 2$, and $|P_r| \leq C_r S_r$. Repeated ordinary integration by parts yields

$$m - m_M = \sum_{r=1}^R (-1)^{r+1} P_r(K) F_K^{(r)}(K) + (-1)^{R+1} \int_K^\infty P_R(x) F_K^{(R+1)}(x) dx.$$

Here $F_K^{(r)}(K) = (-1)^r \Delta_r K^{-\lambda-r}$, which proves the correction and its signs. To bound the remainder, scale $x = Ky$ and use the exact split

$$(\lambda)_{R+1} y^{-2a} - (q)_{R+1} = \Delta_{R+1} y^{-2a} + (q)_{R+1} (y^{-2a} - 1).$$

Against $y^{-\sigma-R-1} dy$, its absolute upper bound integrates to

$$\frac{|\Delta_{R+1}|}{\alpha + R} + |(q)_{R+1}| \left(\frac{1}{\sigma + R} - \frac{1}{\alpha + R} \right) = \mathcal{A}_R.$$

The remaining scaling factor is $K^{-\alpha-R}$, proving (52) for every complex phase. $\square$

Corollary 13.2 (Sharp fixed-trial alignment). *For a fixed nonzero anchored, q -null trial let*

$$\mathcal{B}_c(K) = \sum_d c_d dB_2(\{K/d\}), \quad W(c) = \text{av}_K |\mathcal{B}_c(K)|^2.$$

The average is over a common integer period. Then

$$m - m_M = a\mathcal{B}_c(K)K^{-\lambda-1} + O_{a,q,c}(K^{-\alpha-2}), \quad (54)$$

$$W(c) = \frac{1}{180} \sum_k (J_4(k) + 10J_2(k)) \left| \sum_{k|d} \frac{c_d}{d} \right|^2 > 0, \quad (55)$$

$$\limsup_{M \rightarrow \infty} K^{\alpha+1} |m - m_M| = a \max_K |\mathcal{B}_c(K)| > 0. \quad (56)$$

Writing $Q_{\text{tail}} = Q - Q_M$ and $\pi_M = K^{-2a}$, one also has

$$\frac{K^2 |m - m_M|^2}{\pi_M Q_{\text{tail}}} = \frac{a}{2V(c)} |\mathcal{B}_c(K)|^2 + O_{a,q,c}(K^{-1}). \quad (57)$$

Thus the actual tail alignment is $O(K^{-2})$, with the Cesàro mean of the scaled ratio tending to $aW(c)/(2V(c))$.

Proof. Use $R = 2$ in Theorem 13.1. Since $\Delta_1 = 2a$, the leading term is as stated; one explicit error constant in (54) is $C_2 S_2(|\Delta_2| + \mathcal{A}_2)$. For $g = \gcd(d, e)$ the exact discrete identity is

$$\text{av}_n B_2(\{n/d\}) B_2(\{n/e\}) = \frac{g^4}{180d^2e^2} + \frac{g^2}{18d^2e^2} - \frac{1}{30d^2e^2}.$$

Indeed, conditional on $n \bmod g = z$, multiplication of Bernoulli polynomials gives means $(g/d)^2 B_2(z/g)$ and $(g/e)^2 B_2(z/g)$. Their conditional product is valid by the Chinese remainder theorem, and direct quartic summation gives $g^{-1} \sum_{z=0}^{g-1} B_2(z/g)^2 = 1/180 + 1/(18g^2) - 1/(30g^4)$. Anchoring cancels the last rank-one term and gives $\text{av} \mathcal{B}_c = \sum c_d/(6d) = 0$. The divisor identities for J_4, J_2 prove (55). The largest occupied divisor coordinate proves its strict positivity.

This covariance is credited to [4, Theorem 1]: its wave is $b_d(k) = dB_2(\{k/d\})/2 - 1/(12d)$ and its correlation is $(g^2 - 1)(g^2 + 11)/(720de)$. On the anchored space $\mathcal{B}_c = 2 \sum c_d b_d$, so (55) is four times that correlation form. The calculation here fixes its normalization for the matched Poisson mean.

Periodicity and the uniform-in-residue error in (54) prove (56); the corresponding Cesàro squared limit is $a^2 W(c)$. Finally Theorem 11.1 gives $Q_{\text{tail}} = 2aV(c)K^{-1}(1 + O(K^{-1}))$, proving (57). All these asymptotics hold with the trial, height and parameters fixed; their constants need not be uniform when support or coefficient masses grow. $\square$

14 A corrected complete arithmetic receiver

For coefficient selection, the original common arithmetic head must be kept with the tail. Fix $N \geq 4$ and write

$$\mathcal{H}_N = \{c \in \mathbb{C}^N : A_c(1) = A_c(q) = 0\}, \quad b_M c = m_M, \quad c^* G_M c = Q_M.$$

The entries of G_M are computed from the original states $B_n(e_d)$; these individual coordinate states are used only in finite sums. Complete norms below are asserted on $\mathcal{H}_N$.

Proposition 14.1 (Paid head and norm envelope). *Let V_N have entries $\gcd(d, e)^2/(12de)$ and put*

$$\begin{aligned} D_N &= \sum_{d \leq N} d^2, \quad \gamma = \frac{|q|}{12} \left(1 + \frac{|q+1|}{\sigma+1}\right), \\ r_{N,M} &= \frac{2aD_N + a(2a+1)N/8}{M^2} + \frac{\gamma\sqrt{ND_N}}{(M+1)^2} + \frac{\gamma^2 D_N}{(M+1)^3}, \\ T_+ &= \left(\frac{2a}{M} + \frac{2a}{M^2}\right) V_N + r_{N,M} I, \quad R_M = G_M + T_+. \end{aligned}$$

These matrices are positive definite where needed, and for $c \in \mathcal{H}_N$,

$$Q_{\text{tail}}(c) \leq c^* T_+ c, \quad Q(c) \leq c^* R_M c.$$

In particular the matrix includes the complete norm, not only its head.

Proof. In (41) use $S_0 \leq \sqrt{N} \|c\|$, $S_1 \leq \sqrt{D_N} \|c\|$, $H = S_0/2$, $E = \gamma S_1$. Retain the $2aV/M^2$ term as a multiple of V_N . The other terms are bounded by $r_{N,M} \|c\|^2$ exactly as displayed. The gcd matrix V_N is positive definite by its invertible divisor factorization; also $r_{N,M} > 0$. Adding $G_M \succeq 0$ proves the claims. $\square$

14.1 Choice of a complete norm envelope

The following corrected-gain proposition uses only a positive definite upper for the complete norm on the exact coefficient subspace. Besides the gcd envelope above, Theorem 3.1 supplies

$$R_M^{\text{mix}} = G_M + U_R + N\delta_R I, \quad Q(c) \leq c^* R_M^{\text{mix}} c \quad (c \in \mathcal{H}_N). \quad (58)$$

Here U_R is the positive matrix of (10); a positive extension using its divisor formula is equally valid on the anchored subspace. Every convex combination of complete upper matrices is again an upper. One may select the best certified scalar upper for a fixed trial, while retaining each matrix's actual congruence in a matrix solve.

In exact divisor coordinates $c = Ct$, $C_{dk} = d\mu(k/d)\mathbf{1}_{d|k}$, the full mixed upper on $t_1 = 0$ is

$$C^* G_M C + \frac{2a}{K} \text{diag}_{k \geq 2}(W_{R,k}) + N\delta_R C^* C.$$

The compensated mean row transforms by the same C ; its Euclidean error is charged against $\|Ct\|$. More generally, on any exact double-null basis $c = Jx$, the raw error $N\delta_R I$ becomes $N\delta_R J^* J$. Thus diagonalizing the tail does not replace the common head or the coefficient error by an unrelated identity metric.

For the rest of this section, R_M may denote any such proved positive definite complete envelope, including (58). The displayed gcd choice is its fixed-parameter specialization.

Proposition 14.2 (Corrected gain and constrained scalar test). *For any fixed finite $R \geq 1$ define the computable row and scalar*

$$b = b_M + t_{R,K}, \quad \delta = C_R \left(\sum_{d \leq N} d^{2R} \right)^{1/2} K^{-\alpha-R} \mathcal{A}_R, \quad D = \delta^2 I.$$

For every nonzero $c \in \mathcal{H}_N$,

$$\frac{(\max\{0, |bc| - \delta \|c\|\})^2}{c^* R_M c} \leq g(A_c). \quad (59)$$

Fix a desired gain $\eta > 0$ and any injective basis matrix J for $\mathcal{H}_N$. A trial makes the left side of (59) strictly exceed η if and only if

$$\sup_{0 < \theta < 1} F(\theta) > 1, \quad F(\theta) = bJ \left[J^* \left(\frac{D}{\theta} + \frac{\eta R_M}{1 - \theta} \right) J \right]^{-1} J^* b^*. \quad (60)$$

If $bJ \neq 0$, F is strictly concave and has a unique maximizer, the unique zero of its strictly decreasing derivative. If $bJ = 0$, then $F \equiv 0$ and no trial passes this certificate.

Proof. Cauchy–Schwarz gives $S_R \leq (\sum d^{2R})^{1/2} \|c\|$, so Theorem 13.1 gives $|m - bc| \leq \delta \|c\|$. Proposition 14.1 proves (59). The target inequality is $|bc| > \sqrt{c^* D c} + \sqrt{\eta c^* R_M c}$. For positive x, y ,

$$(\sqrt{x} + \sqrt{y})^2 = \min_{0 < \theta < 1} \left(\frac{x}{\theta} + \frac{y}{1 - \theta} \right).$$

For each θ , the constrained Rayleigh quotient of bc is the displayed inverse expression. The two existential quantifiers over c and θ commute, proving (60).

For completeness whiten $D_H = J^* D J$ and set $w = bJ D_H^{-1/2}$, $S = \eta D_H^{-1/2} J^* R_M J D_H^{-1/2} > 0$. Then

$$F(\theta) = \theta(1 - \theta)w[(1 - \theta)I + \theta S]^{-1}w^*.$$

Spectral decomposition reduces its second derivative to positive weights times $-2s/[1 + (s - 1)\theta]^3$, $s > 0$. Thus $F'' < 0$ if $w \neq 0$, while $F'(0) = ww^* > 0$ and $F'(1) = -wS^{-1}w^* < 0$. This proves uniqueness and permits certified derivative bisection; no secondary extrema need be excluded numerically. $\square$

The worst-response norm and scalar optimization above are standard robust quadratic optimization, as in [6, 7]. The new input is the actual corrected arithmetic row and its complete error. Equivalence in (60) is to the paid certificate, not to the true finite- M gain. No numerical inverse is certified merely by writing its exact formula.

An alternative selection retains the center normalization explicitly. Let C have the three rows $A(1), A(q), A(\lambda)$, and $e = (0, 0, 1)^t$. The columns 1, 2, 4 form a Vandermonde matrix with nodes $2^{-1}, 2^{-q}, 2^{-\lambda}$ of distinct moduli, so C has full row rank. Standard constrained least squares gives

$$\min_{C^* c = e} c^* R_M c = e^* [C R_M^{-1} C^*]^{-1} e.$$

This optimizes a complete upper model and therefore supplies an upper bound for the actual norm of the resulting polynomial. It does not replace the retained full head by its variance capacity.

For perspective, a mean-tail estimate using only the moment information $Q_{\text{tail}} \leq u = c^* T_+ c$ and tail mass π_M gives

$$L_{\text{mom}}(c) = \frac{(\max\{0, |b_M c| - \sqrt{\pi_M u}\})^2}{c^* G_M c + u}.$$

This is sharp for abstract tails with those moment constraints: a constant tail state of opposing phase attains the adverse mean, or reduces it to zero when allowed. Such a state need not be realizable by arithmetic coefficients. Corollary 13.2 proves that its extremal alignment is asymptotically impossible for each fixed nonzero trial. This is the cancellation the corrected row preserves.

Reproducible bounded examples. The accompanying `verify_compensated_means.py` uses exact rational identities and 256-bit Arb/Acb enclosures. It checks 256 discrete pair covariances, seven complex whole-period energy identities, and orders $R = 1, 2, 3$ for five actual arithmetic cases, using $A(s) = (1 - 2^{1-s})(1 - 2^{q-s})$. For these chosen coefficients one may use the sharper scalar error $C_R S_R K^{-\alpha-R} \mathcal{A}_R$ and the scalar norm upper from (41). The same complete denominator is used in the old and corrected certificates. Rounded displays of their order-two values are:

a	T	M	moment certificate	corrected certificate
1/8	0	512	.03133023	.04032653
1/4	3	512	.19172325	.20066055
2/5	0	512	.12157883	.12442350
2/5	3	2048	.28624639	.28717439
2/5	1000	8192	.08507373	.09517727

For the first case the paid mean radius falls from about .01185507 to $2.05425 \cdot 10^{-8}$; at $T = 1000$ it falls from .03760239 to .0002024034. All fifteen finite corrected cases strictly improve their moment comparison, and eight wrong-sign controls are rejected. The verification uses $\zeta(\lambda)A(\lambda)$ only as an independent truth check; the actual corrected certificate requires no zeta value. The explicit rising-factorial and coefficient-mass costs remain essential when parameters or support grow.

15 A complete finite model for optimized gain

The full moving-support norm estimate and compensated mean can be combined before optimization. This controls approximation of the actual optimal gain, with a target implication that prices the complete approximation error.

Fix $0 < a < 1/2$, $\sigma = 1/2 - a$, $\alpha = 1/2 + a$, $q = \sigma + iT$, $\lambda = \alpha + iT$, and let

$$\mathcal{H}_N = \{c \in \mathbb{C}^N : A_c(1) = A_c(q) = 0\}.$$

All suprema below use this same exact double-null space, assumed nontrivial. They need not equal an optimum on the larger space with only $A(1) = 0$. Retain

$$V(c) = \sum_{k=2}^N \frac{J_2(k)}{12} |t_k|^2, \quad C_N = 12 \sum_{k=2}^N \frac{\sigma_1(k)^2}{J_2(k)}, \quad S_0^2 \leq C_N V(c).$$

Suppose the complete cotangent model is

$$\hat{Q}(c) = Q_M(c) + \frac{2a}{K} W(c), \quad W(c) \geq V(c), \quad K = M + 1,$$

and its proved finite error satisfies

$$|Q(c) - \hat{Q}(c)| \leq \frac{2a}{K} H V(c) \leq H \hat{Q}(c), \quad 0 \leq H < 1. \quad (61)$$

In particular the complete common head Q_M is retained.

Theorem 15.1 (Corrected optimized gain). *Let $R \geq 1$, $L_q = |q| + R + 1$, and suppose $0 < \theta < 1$ and $K \geq NL_q/(2\pi\theta)$. Use the actual corrected row $b = b_M + t_{R,K}$ from the compensated mean formula, and define*

$$e = \frac{2\sqrt{2a}}{\pi} \sqrt{C_N} K^{-a} \theta^R \left(2 + \frac{L_q}{R^2}\right),$$

$$G_b = \sup_{c \in \mathcal{H}_N \setminus \{0\}} \frac{|bc|^2}{\widehat{Q}(c)}, \quad G_{\text{true}} = \sup_{c \in \mathcal{H}_N \setminus \{0\}} \frac{|m(c)|^2}{Q(c)}.$$

Then

$$\frac{[\sqrt{G_b} - e]_+^2}{1 + H} \leq G_{\text{true}} \leq \frac{[\sqrt{G_b} + e]^2}{1 - H}. \quad (62)$$

There is also the explicit additive bound

$$|G_{\text{true}} - G_b| \leq H + 2e\sqrt{1 + H} + e^2. \quad (63)$$

In particular

$$G_b > [\sqrt{\eta(1 + H)} + e]^2 \implies G_{\text{true}} > \eta. \quad (64)$$

Proof. The complete compensated-mean estimate is

$$|m - bc| \leq C_R S_R K^{-\alpha - R} \mathcal{A}_R,$$

$$\mathcal{A}_R = \frac{|(\lambda)_{R+1} - (q)_{R+1}|}{\alpha + R} + \frac{2a|(q)_{R+1}|}{(\sigma + R)(\alpha + R)}.$$

The Bernoulli Fourier estimate gives $C_R \leq 4/(2\pi)^{R+1}$, and $S_R \leq N^R \sqrt{C_N V(c)}$. Along the segment from q to λ , each factor in the rising product of length $R + 1$ has magnitude at most L_q , since $2a < 1$. Differentiate that product and integrate over the segment to obtain

$$|(\lambda)_{R+1} - (q)_{R+1}| \leq 2a(R + 1)L_q^R, \quad |(q)_{R+1}| \leq L_q^{R+1}.$$

Because $(R + 1)/(R + \alpha) \leq 2$ and $(R + \sigma)(R + \alpha) \geq R^2$, it follows that

$$\mathcal{A}_R \leq 2aL_q^R(2 + L_q/R^2).$$

Use $NL_q/(2\pi K) \leq \theta$ and $\widehat{Q} \geq 2aV/K$ to conclude

$$|m - bc| \leq e\sqrt{\widehat{Q}(c)}. \quad (65)$$

The divisor transform is invertible on the anchored finite space, so $V(c) > 0$ for nonzero $c \in \mathcal{H}_N$. Thus $\widehat{Q}$ is a norm square there. Let

$$E = \sup_{c \in \mathcal{H}_N \setminus \{0\}} |m(c)|^2 / \widehat{Q}(c).$$

The triangle inequality for dual norms in (65) gives $|\sqrt{E} - \sqrt{G_b}| \leq e$, including vanishing rows. The complete norm comparison (61) gives

$$E/(1 + H) \leq G_{\text{true}} \leq E/(1 - H),$$

proving (62). Also $|E - G_{\text{true}}| \leq HG_{\text{true}} \leq H$ and $E \leq (1 + H)G_{\text{true}} \leq 1 + H$, since true native gain is at most 1. Consequently

$$|G_b - E| \leq 2e\sqrt{E} + e^2 \leq 2e\sqrt{1 + H} + e^2,$$

which proves (63); (64) follows from the lower bound. All coefficient constraints, head terms and errors are on the same space before taking suprema. $\square$

For an injective exact-null basis $c = Jx$,

$$G_b = bJ(J^*\hat{Q}J)^{-1}J^*b^*.$$

Every matrix and error uses this same congruence. This exact formula does not by itself certify a numerical inverse.

Fix a and $\gamma > 1/(2\pi)$, and choose

$$\theta = \frac{1 + 1/(2\pi\gamma)}{2}, \quad K = \lceil \gamma NT \rceil, \quad R = \max\left(1, \left\lceil \frac{3 \log T}{\log(1/\theta)} \right\rceil\right).$$

The wall is eventually admitted uniformly in N . The complete Farey norm comparison has $H = O_{a,\gamma}(N/T + 1/T)$, while $C_N = O(N)$ and the preceding theorem give, for $2 \leq N \leq T$,

$$e = O_{a,\gamma}(N^{1/2-a}T^{-a-2}) = O_{a,\gamma}(T^{-3/2-2a}).$$

Hence along $N = o(T)$ the corrected finite model approximates the actual optimized double-null gain uniformly in an additive sense. The finite target condition (64) converts a certified model gain into a true gain at each admitted height. Its error is additive, so its quantitative payment remains explicit when the model gain is small.

16 Projection disks and arithmetic searches

For completeness, let $\Omega = \{\Re s > 1/2\}$ and $k_\lambda(s) = (s + \bar{\lambda} - 1)^{-1}$ be its Hardy kernel. The image of A is $f_A(s) = a\zeta(s)A(s)/(s - q)$, with $\|f_A\|_{H^2}^2 = aQ(A)/2$. The normalized squared projection of ak_λ onto the span of these images is g_N . If $\zeta(\rho) = 0$, $\rho \in \Omega$, every image vanishes at ρ . Projecting the kernel off k_ρ therefore gives

$$g_N(\lambda) \leq \left| \frac{\rho - \lambda}{\rho + \bar{\lambda} - 1} \right|^2.$$

This is the classical local projection mechanism of [8]. A proved lower certificate $L > 0$ thus excludes zeros from the disk

$$\left| \frac{s - \lambda}{s + \bar{\lambda} - 1} \right| < \sqrt{L}.$$

An implementation must use enclosures for the finite sums in (39) or (59), and pay the coefficient growth in (36) and the explicit correction costs. The upper dual in (28) and the lower certificate measure opposite sides of the same complete arithmetic problem. A family whose certified lower gains exceed a prescribed geometric threshold throughout a height range supplies zero-free disks throughout that range. The compensated numerator retains the exact finite-trial cancellation.

16.1 Continuous height coverage from one seed

Fix the same anchored A , a seed T_0 , and $0 \leq L_0 \leq g_{T_0}$. Set

$$x = |T - T_0|/a, \quad C(x) = \frac{\sqrt{4 + x^2} + x}{2}.$$

Theorem 16.1 (Complete-norm height continuation). *For every real T ,*

$$g_T \geq \left[\frac{\max\{0, \sqrt{L_0} - (x/2)\sqrt{1 - L_0}\}}{C(x)} \right]^2. \quad (66)$$

For a target $\eta < L_0$, the same trial supplies $g_T \geq \eta$ throughout

$$|T - T_0| \leq a \frac{\sqrt{L_0} - \sqrt{\eta}}{\frac{1}{2}\sqrt{1 - L_0} + \sqrt{\eta}}. \quad (67)$$

Proof. Let $F_0(s) = a\zeta(s)A(s)/(s - q_{T_0})$. Proposition 1.1 gives $\|F_0\|^2 = aQ_0/2$ and $F_0(\lambda_{T_0}) = m_0/2$. Its orthogonal decomposition along the Hardy kernel is

$$F_0 = am_0k_{\lambda_{T_0}} + R_0, \quad \|R_0\|^2 = (aQ_0/2)(1 - g_0), \quad R_0(\lambda_{T_0}) = 0.$$

Writing $\Delta = T - T_0$, direct kernel algebra gives

$$\|k_{\lambda_T} - P_{k_{\lambda_{T_0}}}k_{\lambda_T}\|^2 = \frac{\Delta^2}{2a(4a^2 + \Delta^2)}.$$

Since $m_T = (2a + i\Delta)F_0(\lambda_T)/a$, its projected part is exactly m_0 . Thus $|m_T - m_0| \leq \sqrt{Q_0(1 - g_0)}x/2$, preserving its complex phase. The boundary multiplier from F_0 to F_T has squared supremum

$$\sup_{u \in \mathbb{R}} \frac{a^2 + (u + \Delta)^2}{a^2 + u^2} = C(x)^2,$$

so $Q_T \leq C(x)^2Q_0$. The function $\sqrt{g} - (x/2)\sqrt{1 - g}$ is increasing on $[0, 1]$; replace g_0 by its lower L_0 to prove (66). Finally $C(x) \leq 1 + x$ gives (67). $\square$

The exterior null is imposed at the seed to certify its arithmetic tails. Height continuation uses the same anchored trial and its complete Hardy norm; it does not impose that extra null again at neighboring heights. A finite union of the certified intervals gives continuous compact-range coverage.

16.2 A band-adapted sufficient gain

Suppose an outer boundary $1/2 < B < B_0 < 1$ is available. Put $b = B - 1/2$, $c = B_0 - 1/2$. The center minimizing the largest own-height pseudohyperbolic distance from $[B, B_0]$ is

$$\alpha_* = 1/2 + \sqrt{bc}, \quad r_* = (\sqrt{c} - \sqrt{b})/(\sqrt{c} + \sqrt{b}).$$

Indeed the endpoint distances $(v - b)/(v + b)$ and $(c - v)/(c + v)$ balance at $v^2 = bc$, with $v = \alpha - 1/2$. For any radius $r > r_*$, a complete gain family at that center above r^2 excludes the whole band at every height where the family is certified.

Taking the outer boundary $B_0 = 174797/200000$ recorded in [16], an exact rational choice used in the applications is

$$B_0 = \frac{174797}{200000}, \quad B = \frac{377}{432}, \quad \alpha = \frac{43667}{50000}, \quad r = \frac{9}{10000}, \quad \eta = r^2 = \frac{81}{10^8}.$$

The disk diameter endpoints are

$$L = \frac{436726997}{500450000} < B < B_0 < U = \frac{436613003}{499550000}.$$

Accordingly, given that outer boundary, complete gains at least η at every real height imply zeta nonvanishing on $\Re s > B$. This is the same local Hardy disk principle with an explicitly matched band and strict endpoint margins.

17 A degree-uniform obstruction for geometric modules

The complete receiver permits growing arithmetic support. Increasing the degree within a fixed geometric dictionary, however, cannot by itself produce a positive all-height gain. The obstruction below uses the original complete norm, rather than a particular finite certificate.

Keep $0 < a < 1/2$, $\alpha = 1/2 + a$, $\lambda = \alpha + iT$ and $q = 1/2 - a + iT$. For every nonzero finite anchored polynomial $A(1) = 0$, put

$$F(s) = \frac{a\zeta(s)A(s)}{s - q}.$$

The complete recipient gives $F \in H^2(\Re s > 1/2)$, $\|F\|^2 = aQ(A)/2$ and $F(\lambda) = m(A)/2$. The Hardy norm uses $(2\pi)^{-1} \int_{\mathbb{R}}$ on the boundary. The additional null $A(q) = 0$ is not required in this section; every conclusion remains valid on that smaller coefficient space.

Theorem 17.1 (Single geometric lattice). *Fix an integer $b \geq 2$ and let $h = 2\pi/\log b$. Suppose the support of A consists of powers of b . Write $Z_0 = \zeta(\lambda)$ and $Z_1 = \zeta(\lambda + ih)$. If $Z_0 \neq 0$, then*

$$g(A) \leq \frac{h^2|Z_0|^2}{h^2|Z_0|^2 + 4a^2|Z_1 - Z_0|^2}. \quad (68)$$

If $Z_0 = 0$, then $g(A) = 0$. The bound is independent of degree and coefficient magnitudes.

Proof. Here $A(s + ih) = A(s)$ exactly. The Hardy kernel is $k_w(s) = (s + \bar{w} - 1)^{-1}$. For $w = \lambda + ih$, the squared norm of the evaluation residual after projection onto k_λ is $h^2/[2a(4a^2 + h^2)]$. With $\kappa = 2a/(2a + ih)$, its value on F is

$$F(w) - \kappa F(\lambda) = \frac{aA(\lambda)(Z_1 - Z_0)}{2a + ih}.$$

Orthogonal projection onto the two kernels therefore gives

$$\|F\|^2 \geq 2a|F(\lambda)|^2 + \frac{|F(w) - \kappa F(\lambda)|^2}{h^2/[2a(4a^2 + h^2)]}.$$

Substitution yields

$$Q(A) \geq |A(\lambda)|^2 \left(|Z_0|^2 + \frac{4a^2}{h^2}|Z_1 - Z_0|^2 \right).$$

If $A(\lambda) = 0$ the gain vanishes; otherwise division proves (68). The kernel inequality is sharp in the ambient Hardy space; arithmetic attainment is not asserted. $\square$

Fix b and positive integers $d_1, \dots, d_m$. A geometric module here means the polynomials

$$A(s) = \sum_{r=1}^m d_r^{-s} P_r(b^{-s}), \quad (69)$$

where each P_r is any finite polynomial. Both its coefficients and degree may depend on T . Put $z_r = \exp(-ih \log d_r)$ and define

$$1 - \prod_{r=1}^m (1 - z/z_r) = \sum_{j=1}^m \ell_j z^j, \quad L = \sum_{j=1}^m |\ell_j| \sqrt{1 + (jh/(2a))^2}.$$

Repeated nodes are allowed. Elementary symmetric functions give

$$|\ell_j| \leq \binom{m}{j}, \quad L \leq (2^m - 1) \sqrt{1 + (mh/(2a))^2}. \quad (70)$$

Theorem 17.2 (Fixed modules and finite menus). *For A as in (69), let $Z_j = \zeta(\lambda + ijh)$ and assume $Z_j \neq 0$ for $1 \leq j \leq m$. Every nonzero anchored A satisfies*

$$g(A) \leq |Z_0|^2 \left(\sum_{j=1}^m \frac{|\ell_j| \sqrt{1 + (jh/(2a))^2}}{|Z_j|} \right)^2 \leq \frac{L^2 |Z_0|^2}{\min_{1 \leq j \leq m} |Z_j|^2}. \quad (71)$$

Let $G_{b,d}(T)$ be the supremum of true gain over all nonzero anchored polynomials in this module, with no degree bound. For every $\eta > 0$, the set of $T \geq 0$ for which $G_{b,d}(T) < \eta$ has positive lower density. The same assertion holds for the maximum over any finite predeclared collection of fixed modules, including different bases. In particular each such all-degree optimum has lower limit zero.

Proof. Set $A_j = A(\lambda + ijh)$. The defining polynomial takes value 1 at each z_r , so the exact shift relation is

$$A_0 = \sum_{j=1}^m \ell_j A_j.$$

Complete Hardy evaluation at each shifted point gives

$$Q(A) \geq \frac{4a^2}{4a^2 + (jh)^2} |Z_j A_j|^2.$$

The triangle inequality proves (71).

Use the standard Voronin–Reich–Bagchi universality theorem in its compact-set form: non-vanishing continuous targets, analytic in the interior of a compact subset of $1/2 < \Re s < 1$ with connected complement, are approximated within any fixed positive tolerance by a positive-lower-density set of vertical shifts; see [20, p. 59]. Choose small disjoint closed disks centered at $\alpha, \alpha + ih, \dots, \alpha + imh$, contained in that strip. Their union has connected complement. Take target $\varepsilon > 0$ on the central disk and target 1 on each other disk. These targets are nowhere zero. Approximation with tolerance $\varepsilon < 1/2$ gives

$$|Z_0| < 2\varepsilon, \quad |Z_j| > 1 - \varepsilon > 1/2 \quad (1 \leq j \leq m)$$

on a positive-lower-density set of actual T . Hence the supremum in (71) is less than $16L^2\varepsilon^2$. Choose also $\varepsilon < \sqrt{\eta}/(4L)$. For a finite menu, use the finite union of all distinct positive probe offsets and the maximum of the corresponding constants L ; coincident probes receive the same target 1. This is one universality application on a fixed compact set. Suprema are measurable: fixed anchored finite coefficient spaces admit countable dense sets of trials with continuous gain, and only countably many finite degrees are involved. $\square$

Proposition 17.3 (A paid escape condition). *Keep the fixed nodes and assume $\min_{1 \leq j \leq m} |Z_j| > 0$. For any finite anchored arithmetic polynomial, not necessarily supported in the module, set*

$$J(z) = \prod_{r=1}^m (1 - z/z_r), \quad E(A) = \sum_d c_d d^{-\lambda} J(e^{-ih \log d}).$$

Then

$$A_0 - E(A) = \sum_{j=1}^m \ell_j A_j, \quad |A_0 - E(A)| \leq \frac{L \sqrt{Q(A)}}{\min_{1 \leq j \leq m} |Z_j|}. \quad (72)$$

Any family with $|E(A)| \leq \delta|A_0|$ for a fixed $0 \leq \delta < 1$ satisfies (71) with an additional factor $(1 - \delta)^{-2}$, and therefore has the same all-height obstruction. Conversely, if $g(A) \geq \eta > 0$ and $Z_0 \neq 0$, then

$$\left|1 - \frac{E(A)}{A_0}\right| \leq \frac{L|Z_0|}{\sqrt{\eta} \min_{1 \leq j \leq m} |Z_j|}. \quad (73)$$

If off-module terms denote those indices outside $\bigcup_{r=1}^m \{d_r b^k : k \geq 0\}$, their coefficient cost satisfies

$$\sum_{d \text{ off module}} |c_d| d^{-\alpha} \geq 2^{-m} |A_0| \left[1 - \frac{L|Z_0|}{\sqrt{\eta} \min_{1 \leq j \leq m} |Z_j|}\right]_+.$$

Proof. The polynomial identity defining J gives the first part of (72) term by term. The same complete evaluations used above give its inequality. The relative-defect conclusion follows from $|A_0 - E(A)| \geq (1 - \delta)|A_0|$. For a successful trial, $A_0 \neq 0$ and $\sqrt{Q}/|A_0| = |Z_0|/\sqrt{g(A)}$; division proves (73). Module-supported terms contribute zero to E , and $|J(z)| \leq 2^m$ on the unit circle. These facts and the reverse triangle inequality prove the last bound. Exact module support implies $E = 0$; the converse is not claimed. $\square$

The conclusions hold for every geometric depth with the base and finite module generators fixed. Their positive-lower-density assertion is qualitative. On an admitted $N = o(T)$ support schedule, the additive corrected-model comparison transfers the lower-limit-zero conclusion to the *same restricted module subspace*. The paid escape condition quantifies the off-module evaluation required of a successful growing dictionary construction.

18 Reproducible complete native applications

All three cases use $a = 18667/50000$, $\alpha = 43667/50000$, $\sigma = 6333/50000$. The public verifier uses exact rational Bernoulli values, reduced-divisor covariance and discrete seam identities. Directed 512-bit Arb/Acb calculations then test the actual complete arithmetic head and both infinite mean/norm tails. Its three-coefficient family is defined analytically by

$$c_3 = 1, \quad c_2 = \frac{3^{-q} - 1/3}{1/2 - 2^{-q}}, \quad c_1 = -c_2/2 - 1/3.$$

Both $A(1) = A(q) = 0$ hold identically. The denominator cannot vanish because $|2^{-q}| = 2^{-\sigma} > 1/2$. Each head is accumulated from its same actual B_n states; the new covariance weights use the same complex coefficients and full mixed orders.

The mean uses Theorem 13.1 at order $R+2$. The accepting numerator is its finite corrected row minus its complete error. The denominator is $Q_M + U_R(c) + \delta_R S_0^2$. Direct evaluation of $\zeta(\lambda)A(\lambda)$ is an independent remainder control. The resulting rounded-down gains and height radii are

T	K	R	θ	complete gain lower	height halfwidth
1000	1024	23	1/2	.1971551245	.3685275677
10000	16384	30	1/2	.03521509868	.14171006664
1000	768	56	3/4	.1969254372	.3682598316

The halfwidths use (67) with $\eta = 81/10^8$. The first two complete norm-tail uppers are, respectively, less than .000176596358 and .000010561003. Compared with the scalar $25S_0^2/(16K)$ envelope, their tail tightenings exceed 42.1372 and 92.7486. The third uses the general $\theta = 3/4$ wall, below the

half-ratio wall; its comparison is the independently paid $(b_\theta + e_\theta)^2 S_0^2 / K$ envelope, with tightening greater than 42.1073. These are bounds on tail certification; the head and corrected mean remain common in each comparison.

The script verifies the direct whole-period mixed source against the divisor form, exact J_2 and first Bernoulli covariance cases, positive higher-Jordan inflation and its upper bounds, and independently summed tail segments through $4K - 1$. Normal and optimized executions agree in every mathematical output field. They pass 75,193 checks, including 2,354 exact rational identities; these controls accompany the analytic proofs of the whole tails and all-parameter statements.

The package includes the retained exact dual, null, gcd-tail, capacity, compensated-mean and robust-certificate verifiers, together with the new full mixed-order program. The reader instructions specify the tested dependencies and reproduction commands.

A A paired-wave scalar alternative

Here a short classical nonstationary argument supplies a scalar alternative to the mixed-order envelope. Under both nulls, the exact identity from Lemma 2.1 before integration by parts is

$$B_n = n^{-q} H_{0,c}(n) - q \int_n^\infty H_{0,c}(x) x^{-q-1} dx.$$

Assume $T \geq 0$, conjugating q and all coefficients otherwise. For $\omega = 2\pi k/d$, $k \geq 1$, the two integrals

$$I_\pm = \int_n^\infty x^{-\sigma-1} e^{i(\pm\omega x - T \log x)} dx$$

satisfy, when $n \geq N|q|/\pi$,

$$|I_\pm| \leq \frac{2n^{-\sigma-1}}{\omega \mp T/n}.$$

The quotients $x^{-\sigma}/(\omega x \mp T)$ are positive, decreasing to zero; integration by parts pays their endpoint and total variation. This is the standard monotone first-derivative estimate [14, 13].

Use the Abel Fourier series for the sawtooth only under the integral. Pair the signs before summing their coefficient magnitudes $1/(2\pi k)$:

$$\frac{|I_+| + |I_-|}{2\pi k} \leq \frac{dn^{-\sigma-1}}{\pi^2(k^2 - \rho^2)}, \quad \rho = dT/(2\pi n) \leq 1/2.$$

The majorant is summable and $\sum_{k \geq 1} (k^2 - 1/4)^{-1} = 2$ by telescoping. Abel averages have magnitude at most $1/2$; dominated convergence with $x^{-\sigma-1}$ returns the actual integral. The original inclusive endpoint $H_{0,c}(n)$ remains unchanged. Therefore

$$|B_n| \leq C_* S_0 n^{-\sigma}, \quad C_* = \frac{1}{2} + 2/\pi < 5/4, \quad n \geq N|q|/\pi.$$

Summing the complete positive p_n weights gives

$$Q - Q_M \leq C_*^2 S_0^2 / K, \quad |m - m_M| \leq C_* S_0 K^{-\alpha} \quad (K \geq N|q|/\pi).$$

This alternative credits classical height-uniform truncation; the mixed-order envelope retains the finer periodic covariance and resummed height-scale capacity.

Acknowledgment of assistance. AI assistants supported derivation, literature comparison, manuscript preparation and independent checks. The author is responsible for the mathematical statements and presentation.

References

- [1] M. Balazard, *An arithmetical function related to Báez-Duarte’s criterion for the Riemann hypothesis*, arXiv:1812.04309v1 (2018), Sections 2.1–2.2.
<https://arxiv.org/abs/1812.04309>.
- [2] halasz_resonator, *Harmonic–Möbius Unitriangular Factorization, $15/\pi^2$ Dual Norm Asymptotics, and Wavelet Head-to-Tail Reconstruction in the Nyman–Beurling–Báez-Duarte Criterion*, rStation paper 33, version 1 (19 October 2026), Theorems 2–3.
https://rstation.ai/problems/riemann_hypothesis/papers/33/v1.
- [3] halasz_resonator, *Step-Space Annihilator Duality in the Shifted Nyman-Beurling Criterion: Power-Saving Head Norm Asymptotics, Flat-Top Hermite Spline Zero Annihilators, Summation-by-Parts Periodic-Mean Tail Enclosures, and Interval-Certified Conjugate-Kernel Projections*, rStation paper 9, version 1 (16 October 2026), Sections 2.1–2.3.
https://rstation.ai/problems/riemann_hypothesis/papers/9.
- [4] halasz_resonator, *Bernoulli-2 Farey Large-Sieve Neutral Descent and Supercritical Truncated Divisor Covariance*, rStation paper 43, version 4 (24 October 2026), Theorem 1.
https://rstation.ai/problems/riemann_hypothesis/papers/43.
- [5] NIST Digital Library of Mathematical Functions, Sections 24.4 and 24.8, Bernoulli identities and Fourier series. <https://dlmf.nist.gov/24>.
- [6] R. G. Lorenz and S. P. Boyd, *Robust minimum variance beamforming*, IEEE Trans. Signal Process. **53** (2005), 1684–1696. <https://doi.org/10.1109/TSP.2005.845436>.
- [7] S. A. Vorobyov, A. B. Gershman and Z.-Q. Luo, *Robust adaptive beamforming using worst-case performance optimization*, IEEE Trans. Signal Process. **51** (2003), 313–324.
<https://doi.org/10.1109/TSP.2002.806865>.
- [8] C. Delaunay, E. Fricain, E. Mosaki and O. Robert, *Zero free regions for Dirichlet series*, Trans. Amer. Math. Soc. **365** (2013), 3227–3253; arXiv:1101.1199.
<https://arxiv.org/abs/1101.1199>.
- [9] H. L. Montgomery, *Extreme values of the Riemann zeta function*, Comment. Math. Helv. **52** (1977), 511–518. <https://doi.org/10.5169/seals-40013>.
- [10] C. Aistleitner, I. Berkes and K. Seip, *GCD sums from Poisson integrals and systems of dilated functions*, J. Eur. Math. Soc. **17** (2015), 1517–1546, Section 5; arXiv:1210.0741v5.
<https://doi.org/10.4171/JEMS/537>.
- [11] P. Lindqvist and K. Seip, *Note on some greatest common divisor matrices*, Acta Arith. **84** (1998), 149–154. <https://doi.org/10.4064/aa-84-2-149-154>.
- [12] S. Schubr, *Collision-uniform kernel asymptotics for Báez-Duarte zero packets*, edition 2, Luminary Research Archive (2026), with mathematical supplement.
<https://research.luminaryoddities.com/collision-uniform-kernel-asymptotics-baez-duarte-zero-packets/v/2>.
- [13] E. C. Titchmarsh, *The Theory of the Riemann Zeta-Function*, second edition, revised by D. R. Heath-Brown, Clarendon Press, Oxford, 1986, Theorem 4.11 and Section 4.14.

- [14] J. Arias de Reyna, *On the approximation of the zeta function by Dirichlet polynomials*, arXiv:2406.16667v1 (2024). <https://arxiv.org/abs/2406.16667v1>.
- [15] H. Maier and M. Th. Rassias, *Generalizations of a cotangent sum associated to the Estermann zeta function*, arXiv:1410.2145v1 (2014), Section 1.1. <https://arxiv.org/abs/1410.2145v1>.
- [16] QuasiRiemannTracker contributors, *QuasiRiemannTracker*, pull request 84, outer Dirichlet nonvanishing boundary 174797/200000, merged 27 October 2026. <https://github.com/GTimG/QuasiRiemannTracker/pull/84>.
- [17] H. L. Montgomery and R. C. Vaughan, *The large sieve*, *Mathematika* **20** (1973), 119–134, Section 2. <https://doi.org/10.1112/S0025579300004708>.
- [18] K. S. Kedlaya, *Notes on analytic number theory*, Chapter 17, Corollaries 17.2–17.3 and Section 17.3. <https://kskedlaya.org/ant/chapter-17.html>.
- [19] halasz_resonator, *Meissel–Báez–Duarte Floor Duality, Raabe–Möbius Binet–Bernoulli Convolution Algebras, and Farey–Nyquist Tail Large Sieve Bounds in Nyman–Beurling Space*, rStation paper 51, version 2 (25 October 2026), Sections 2–5. https://rstation.ai/problems/riemann_hypothesis/papers/51.
- [20] Ł. Pańkowski, *Hybrid joint universality theorem for Dirichlet L -functions*, *Acta Arithmetica* **141** (2010), no. 1, 59–72, p. 59. <https://doi.org/10.4064/aa141-1-3>.