

Collision-uniform kernel asymptotics for Báez–Duarte zero packets

Scott Schubr

Luminary Oddities

Countryside, Illinois, United States

Corresponding author: scottschubr@luminaryoddities.com

ORCID: 0009-0009-7357-4439

October 25, 2026

Abstract

We prove collision-uniform high-height asymptotics for logarithmically growing zero packets in the harmonic-step space underlying Báez–Duarte approximation. An exact arithmetic gauge converges to an explicit complex Mellin kernel, with literal mixed-derivative errors at every finite order. A positive Hardy subkernel and an explicit rational Newton–Hermite interpolant give a separation-free Gram floor whose price is exponential in packet cardinality. Combining these bounds yields relative Gram comparison uniformly for arbitrary collisions and for cardinality at most a sufficiently small explicit multiple of the logarithm of height. When the real-part rectangle approaches both open walls of the strip at an inverse logarithmic rate, packets of size proportional to the logarithm of height divided by its iterated logarithm remain admissible, with an explicit restriction on the proportionality constant. At admitted zeta-zero packets, exact native zero-jet projection retains the target’s unequal powers and oscillatory phases. We also give a literal onset, a vanishing sublogarithmic packet budget, all-height regularized certificates, and fixed-multiplicity consequences. Complete arithmetic tail estimates and supporting norm comparisons are included; the argument uses the original harmonic-step kernel throughout.

Keywords: Nyman–Beurling approximation; Mellin kernel; zero packets; divided differences; reproducing kernels; arithmetic certificates.

Mathematics Subject Classification (2020): 11M26; 46E22; 30H10.

1 Logarithmically growing harmonic-step packets

Let S be the closed subspace of $L^2(0, 1)$ consisting of functions constant on every interval

$$I_n = (1/(n+1), 1/n], \quad n \geq 1.$$

We use the Mellin transform $\hat{f}(s) = \int_0^1 f(t)t^{s-1} dt$ and identify S with its Mellin image. It is a subspace of $H^2(\Omega)$, where $\Omega = \{\Re s > 1/2\}$ and the squared boundary norm is $(2\pi)^{-1} \int_{\mathbb{R}} |f(1/2 + it)|^2 dt$. The inner product is linear in its first argument. The orthogonal basis and its squared norms are

$$e_n(s) = \frac{n^{-s} - (n+1)^{-s}}{s}, \quad \|e_n\|^2 = \frac{1}{n(n+1)}. \quad (1)$$

Thus the evaluation kernel of S is

$$K_S(s, w) = \sum_{n \geq 1} n(n+1) e_n(s) \overline{e_n(w)}. \quad (2)$$

This kernel retains the harmonic-step restriction, rather than replacing S by the full half-plane Hardy space.

The Báez–Duarte approximation problem [1, 2] concerns

$$E(s) = \frac{1}{s}, \quad G_l(s) = \frac{(l^{-s} - l^{-1})\zeta(s)}{s} \quad (l \geq 2), \quad V = \overline{\text{span}}\{G_l : l \geq 2\} \subset S. \quad (3)$$

Write $D_\infty = \text{dist}(E, V)$. Bounded zero-jet functionals on S give lower bounds for this distance. At high height their conditioning depends on the relative positions of the zeros, their multiplicities, and the phase of the target $1/s$.

The step-space zero-jet projection framework and the sharp diagonal asymptotic were developed in [3, Sections 2–5]. Weighted finite-cutoff conjugate-pair kernels and their monotonicity were studied in [4, Theorem 4]. Our question is the full packet limit, including unequal real parts and offsets that may collide as the height grows. Entrywise kernel asymptotics alone are insufficient in that regime: an evaluation Gram matrix can become arbitrarily ill-conditioned. We obtain a comparison of quadratic forms after a collision-compatible change of basis.

The multi-zero superposition and collective projection bounds of `halasz_resonator` [5, Theorems 3–4 and Section 4.1] already include a continuum kernel and cutoff-dependent phase interference for fixed distinct zeros in weighted step spaces. Here the asymptotic parameter is the height in the exact unweighted native kernel; moving collisions, confluent jets, a uniform Gram floor and all-height regularization are treated together.

Growing packets and approaching boundaries. The main quantitative conclusion is the harmonic-step growing-packet theorem, Corollary 6.3. For a fixed rectangle Q , the explicit relative error is $A_Q m^2 \Lambda_Q^{m-1} T^{-\kappa}$, so every packet size $m \leq c \log T$ is admitted when $c \log \Lambda_Q < \kappa$. An exponential packet Gram bound, Theorem 6.2, pays the conditioning cost uniformly through all collisions. The approaching-boundary packet theorem, Corollary 6.5, allows the rectangle to approach both real-part walls at distance $(\log T)^{-a}$ while $m \leq c \log T / \log \log T$ for $c < 1/(4a)$. The fixed-size statements below establish the kernel and literal error used in these growing-size conclusions. The multiplicity leading-term formula in Section 9 retains fixed multiplicity.

1.1 The fixed-size kernel foundation

Fix an integer $m \geq 1$ and a compact rectangle

$$Q = \{q \in \mathbb{C} : b_0 \leq \Re q \leq b_1, |\Im q| \leq L\}, \quad \frac{1}{2} < b_0 \leq b_1 < 1, \quad (4)$$

where $L < \infty$. For $T \geq 4$, put $s_q = iT + q$ and define

$$A_T(q, p) = T^{q+\bar{p}-3} s_q \bar{s}_p K_S(s_q, s_p), \quad T^q = \exp(q \log T). \quad (5)$$

The logarithm of T is real. The limiting kernel is

$$\begin{aligned} H(q, p) &= \int_0^\infty x^{2-q-\bar{p}} 4 \sin^2 \frac{1}{2x} dx \\ &= \frac{\pi}{\Gamma(4-q-\bar{p}) \sin(\pi(3-q-\bar{p})/2)}. \end{aligned} \quad (6)$$

For nodes $\mathbf{q} = (q_1, \dots, q_m) \in Q^m$, let $\Delta_j g = [q_1, \dots, q_j]g$ be the complex Newton divided difference, with its Hermite interpretation at repeated nodes. In particular, $[a, \dots, a]g = g^{(j-1)}(a)/(j -$

1)!. For a sesquiholomorphic kernel, $\overline{\Delta_k}$ denotes the anti-holomorphic divided difference in the second variable, with conjugate nodes and conjugate coefficients. Set

$$\mathbf{G}_T(\mathbf{q})_{jk} = \Delta_j^{(q)} \overline{\Delta_k^{(p)}} A_T(q, p), \quad \mathbf{G}(\mathbf{q})_{jk} = \Delta_j^{(q)} \overline{\Delta_k^{(p)}} H(q, p). \quad (7)$$

Theorem 1.1 (Collision-uniform packet kernel). *For every fixed $r, k \geq 0$,*

$$\sup_{q, p \in Q} |\partial_q^r \partial_{\bar{p}}^k (A_T - H)(q, p)| \longrightarrow 0 \quad (T \rightarrow \infty). \quad (8)$$

The matrices $\mathbf{G}(\mathbf{q})$ are positive definite with a common positive smallest-eigenvalue bound on Q^m . There is $\eta_T \rightarrow 0$, depending only on Q, m , such that uniformly for all $\mathbf{q} \in Q^m$,

$$(1 - \eta_T) \mathbf{G}(\mathbf{q}) \preceq \mathbf{G}_T(\mathbf{q}) \preceq (1 + \eta_T) \mathbf{G}(\mathbf{q}). \quad (9)$$

Arbitrary partial and complete collisions are included.

The native target vector is

$$z_T(\mathbf{q})_j = [q_1, \dots, q_j] T^q. \quad (10)$$

Corollary 1.2 (Zero-packet distance bound). *Suppose the points $iT + q_j$ are zeros of ζ , with each repeated point listed at most its multiplicity. For all sufficiently large T ,*

$$D_\infty^2 \geq T^{-3} z_T^* \mathbf{G}_T^{-1} z_T \geq \frac{T^{-3}}{1 + \eta_T} z_T^* \mathbf{G}^{-1} z_T. \quad (11)$$

The same η_T works for every such packet in Q^m .

All parameters in Q and m are fixed in these statements. In particular, no separation is imposed between the nodes, whereas the size of the packet and its compact real-part range remain specified.

1.2 A certificate at every finite height

The quantitative statements use an expanded rectangle $Q_e = \{b \leq \Re q \leq c, |\Im q| \leq L_e\}$, with $1/2 < b \leq c < 1$. For any $B \geq \sup_{Q_e} |q|$, set

$$C = 1 + B, \quad g = 2b - 1, \quad h = 2 - 2c, \quad \kappa = \frac{3}{2} - c, \quad J = C^2(64 + 16/g + 16/h). \quad (12)$$

Choose $\rho > 0$ such that the closed radius- ρ disks about points of Q lie in Q_e . Define

$$E_0 = JT^{-\kappa}, \quad a_j = \rho^{-(j-1)}, \quad \epsilon = E_0 \sum_{j=1}^m a_j^2.$$

Theorem 1.3 (Effective native packet certificate). *For every real $T \geq 4$ and $q, p \in Q$,*

$$|\partial_q^r \partial_{\bar{p}}^k (A_T - H)(q, p)| \leq r!k! \rho^{-r-k} J T^{-\kappa} \quad (r, k \geq 0). \quad (13)$$

For every tuple in Q^m , including all collisions,

$$|(\mathbf{G}_T - \mathbf{G})_{jk}| \leq E_0 a_j a_k, \quad \mathbf{G}_T \preceq \mathbf{G} + \epsilon I. \quad (14)$$

If the tuple is a zeta-zero packet as in Corollary 1.2, then, at every such height,

$$D_\infty^2 \geq T^{-3} z_T^* (\mathbf{G} + \epsilon I)^{-1} z_T. \quad (15)$$

The inverse in this formula exists without a small-error condition.

Sections 5–8 prove this theorem, an adaptive derivative estimate retaining the exponent $3/2 - b_1$, and an explicit positive floor for $\mathbf{G}$. The floor gives a literal onset for the relative law. The additive inverse in (15) avoids that onset altogether. Section 10 connects these bounds to finite arithmetic primal and dual certificates in the same Hilbert norm.

1.3 Related analytic interfaces

The kernel-translation operator of Manzur–Noor [6, Sections 4–5] provides a different quantitative zero obstruction and an invariant-subspace criterion. The shifted-Hardy approximation of Manzur–Noor–Quintero [7, Section 3] also concerns the Báez–Duarte generators, with a different approximation norm. Our result addresses the high-height geometry of the exact kernel (2). The one-point exponent and constant, together with an elementary diagonal power-error argument, are inherited from [3, Section 5, Remark 4]. The full complex kernel, literal mixed-derivative constants, separation-free Gram floor, relative comparison and exact regularized packet receiver are developed below. The divided-difference integral and the positive moment-matrix argument are classical tools.

Collision-safe cluster estimates also occur in the conditional approximation-rate construction of `bramble_trace` [8]. That work constructs Dirichlet-polynomial approximants in the classical Nyman–Beurling norm under RH, or under a finite off-line-zero hypothesis. The present theorem instead compares the exact harmonic-step evaluation kernel on fixed compact packets, including unequal real parts, and retains its native Mellin-jet recipient.

2 The native kernel and finite zero-jet projection

For $\rho \in \Omega$, the Mellin evaluation functional is represented in $L^2(0, 1)$ by $t^{\bar{\rho}-1}$, of squared norm $1/(2\Re \rho - 1)$. Its projection onto S is the restricted representer r_ρ . This proves boundedness of evaluation. Higher jets are bounded as well, since $t^{\bar{\rho}-1}(\log t)^j \in L^2(0, 1)$. Expanding in the orthogonal basis (1) proves (2), with local convergence of all required derivatives. Alternatively, these follow directly by projecting the bounded full-space jet functionals.

For completeness, G_l belongs to S . In the original variable it is the negative Mellin transform of

$$\rho_{1/l}(t) = \{1/(lt)\} - l^{-1}\{1/t\}.$$

If $1/t = n + \theta$, $0 \leq \theta < 1$, then $\rho_{1/l}(t) = \{n/l\}$, so it is constant on each I_n and bounded. The Mellin identity follows first where the defining zeta integrals converge and then by continuation. Each zero of ζ therefore forces the corresponding jets of every G_l to vanish.

We use the following elementary projection principle, with the evaluation-row convention of (2).

Lemma 2.1. *Let $\Lambda_1, \dots, \Lambda_d$ be independent bounded linear functionals on a complex Hilbert space, annihilating a closed subspace V . If r_j is the Riesz representer of Λ_j , put $\Gamma_{jk} = \langle r_k, r_j \rangle$ and $c_j = \Lambda_j(E)$. Then*

$$\text{dist}(E, V)^2 \geq c^* \Gamma^{-1} c.$$

Proof. The representers lie in $V^\perp$. If the projection of E onto their span is $\sum_k a_k r_k$, its functional values give $\Gamma a = c$. Its squared norm is $a^* \Gamma a = c^* \Gamma^{-1} c$. Projection onto this subspace cannot exceed projection onto $V^\perp$. $\square$

This argument uses only finitely many bounded functionals. It does not require a description of the entire orthogonal complement of V .

3 The exact arithmetic gauge and its smooth limit

For $x = n/T$, define

$$F_T(q, x) = \sqrt{1 + 1/n} x^{1-q} [1 - (1 + 1/n)^{-iT-q}]. \quad (16)$$

Expansion of (2) gives the exact identity

$$A_T(q, p) = \frac{1}{T} \sum_{n \geq 1} F_T(q, n/T) \overline{F_T(p, n/T)}. \quad (17)$$

In particular, the factors $s_q \overline{s_p}$ in (5) have not been approximated by T^2 .

Lemma 3.1. *For every fixed integer $r \geq 0$, constants depending on Q, r give*

$$\begin{aligned} |\partial_q^r F_T(q, x)| &\leq C_r x^{1-b_1} (1 + |\log x|)^r & (x \leq 1), \\ |\partial_q^r F_T(q, x)| &\leq C_r x^{-b_0} (1 + \log x)^r & (x \geq 1), \end{aligned} \quad (18)$$

uniformly for $q \in Q$ and $T > 2L + 2$.

Proof. Put $a_n = \log(1 + 1/n)$. The bracket in (16) has modulus at most two, since $\Re(iT + q) > 0$, and its positive derivatives have modulus at most a_n^j . On the small- x range use $a_n \leq \log 2$ and $\sqrt{1 + 1/n} \leq \sqrt{2}$. On the large- x range, the integral identity for $1 - e^{-z}$, $\Re z \geq 0$, yields

$$|1 - e^{-(iT+q)a_n}| \leq |iT + q| a_n \leq CT/n = C/x.$$

For $j \geq 1$, $a_n^j \leq C_j/n \leq C_j/x$ there. Differentiating x^{1-q} supplies only powers of $\log x$. Finally use $x^{1-\Re q} \leq x^{1-b_1}$ for $x \leq 1$ and $x^{-\Re q} \leq x^{-b_0}$ for $x \geq 1$. $\square$

Proof of (8). On any fixed $\varepsilon \leq x \leq M$, uniformly in $q \in Q$,

$$F_T(q, x) \longrightarrow F(q, x) = x^{1-q} (1 - e^{-i/x}),$$

with every fixed number of q derivatives. Indeed $T \log(1 + 1/n) \rightarrow 1/x$ uniformly on this range, while the derivatives of the bracket of positive order introduce powers of $\log(1 + 1/n) = O(1/T)$. Uniform Riemann sums on the compact interval therefore converge, including mixed derivatives in the two kernel slots.

We verify the infinite tails separately. By Lemma 3.1, a mixed derivative of orders r, k of the summand product is bounded at zero by

$$Cx^{2-2b_1} (1 + |\log x|)^{r+k},$$

and at infinity by

$$Cx^{-2b_0} (1 + \log x)^{r+k}.$$

Both are integrable. On a sufficiently small interval the first envelope is increasing, so its mesh sum through ε is bounded by its integral through 2ε ; an empty mesh contributes zero. On a sufficiently late tail the second envelope is decreasing, and its mesh sum is bounded by an integral beginning at $M/2$. These bounds tend to zero uniformly as $\varepsilon \downarrow 0$, $M \rightarrow \infty$. They justify the termwise derivatives of (17) and the differentiated limiting integral. Choosing the tails first and then taking the compact Riemann limit proves (8).

To evaluate the integral, put $u = 1/x$ and $\alpha = 3 - q - \bar{p}$, so $1 < \Re \alpha < 2$. The Mellin identity

$$2 \int_0^\infty (1 - \cos u) u^{-1-\alpha} du = \frac{\pi}{\Gamma(1 + \alpha) \sin(\pi\alpha/2)}, \quad 0 < \Re \alpha < 2,$$

gives (6). One derivation integrates by parts and uses the sine Mellin integral, first on an overlapping convergence strip and then by analytic continuation. Absolute convergence of the displayed $1 - \cos u$ integral throughout this strip also identifies the same analytic continuation directly. $\square$

4 Collisions and relative inversion

The complex Hermite–Genocchi formula is

$$[q_1, \dots, q_j]g = \int_{\substack{\theta_l \geq 0 \\ \sum \theta_l = 1}} g^{(j-1)} \left(\sum_l \theta_l q_l \right) d\theta, \quad (19)$$

where simplex measure has total mass $1/(j-1)!$. Repeated fundamental-theorem integration proves the formula for complex nodes as well as real ones; continuity gives every Hermite endpoint. The convexity of Q keeps all arguments inside Q .

Applying (19) in both slots of (8), with orders at most $m-1$, gives

$$\sup_{\mathbf{q} \in Q^m} \|\mathbf{G}_T(\mathbf{q}) - \mathbf{G}(\mathbf{q})\| \longrightarrow 0. \quad (20)$$

The conjugation of the second divided-difference coefficients is essential here.

Lemma 4.1. *The limiting Gram matrix $\mathbf{G}(\mathbf{q})$ has a common positive eigenvalue floor on Q^m .*

Proof. Set $\psi_j(x) = [q_1, \dots, q_j]x^{-q}$. Then

$$\mathbf{G}(\mathbf{q})_{jk} = \int_0^\infty \psi_j(x) \overline{\psi_k(x)} x^2 4 \sin^2 \frac{1}{2x} dx. \quad (21)$$

The base measure has infinite mass, but all the displayed products are integrable by the preceding envelopes. In $y = \log x$, the expansion of e^{-qy} gives

$$\psi_j(e^y) = \frac{(-y)^{j-1}}{(j-1)!} + O(y^j).$$

An order- $j-1$ divided difference kills q^k for $k < j-1$ and sends q^{j-1} to one, including at collisions. The triangular leading orders therefore make $\psi_1, \dots, \psi_m$ linearly independent for every tuple. A nontrivial linear combination is an entire exponential polynomial in y and cannot vanish on a set of positive measure. Since the weight in (21) is positive off a discrete set, the matrix is positive definite.

The simplex formula and the uniform integrable logarithmic envelopes make the matrix continuous on Q^m . Its least eigenvalue thus has a positive minimum on this compact set. $\square$

Completion of Theorem 1.1. Let $c > 0$ be the floor in Lemma 4.1, and let ϵ_T be the supremum in (20). For every vector v ,

$$|v^*(\mathbf{G}_T - \mathbf{G})v| \leq \epsilon_T \|v\|^2 \leq (\epsilon_T/c) v^* \mathbf{G} v.$$

Taking $\eta_T = \epsilon_T/c$ proves (9). $\square$

Proof of Corollary 1.2. Use the native functionals

$$\Lambda_j(f) = [q_1, \dots, q_j](s_q T^q f(s_q)). \quad (22)$$

Their Gram matrix is exactly $T^3 \mathbf{G}_T$, by (5). Their values on $E = 1/s$ are exactly z_T from (10). The gauge $s_q T^q$ is analytic and nonzero on a neighborhood of Q , and multiplication by it is triangular and invertible on each finite jet space. The derivative of s_q with respect to q is one. Thus repeated nodes in (22) are invertible combinations of the native Mellin jets, with the stated Hermite factorials.

Under the zero-packet hypothesis these functionals annihilate all G_l and hence V . The positive definiteness from (9) makes them independent for large T . Lemma 2.1 gives the first inequality of (11). Inverting the upper inequality of (9) gives the second. The same proof applies if the complete off-line zero set is infinite, since only the listed finite packet enters the projection. $\square$

5 An explicit arithmetic kernel error

We first prove a scalar estimate on the whole expanded rectangle Q_e . The derivative estimates will then follow from complex Cauchy bounds. This separates the arithmetic error from the conditioning of any packet.

Proposition 5.1. *With the constants in (12), for every real $T \geq 4$,*

$$\sup_{q,p \in Q_e} |A_T(q,p) - H(q,p)| \leq JT^{-\kappa}. \quad (23)$$

Proof. Put $z = q + \bar{p}$, $a = \Re z \in [2b, 2c] \subset (1, 2)$, and

$$b_q(n) = 1 - (1 + 1/n)^{-iT-q}, \quad b_0(n) = 1 - e^{-iT/n}.$$

Both exponents in this comparison have nonpositive real part. The line-segment integral formula for the exponential, and $0 \leq u - \log(1 + u) \leq u^2/2$, give

$$|b_q(n) - b_0(n)| \leq B/n + T/(2n^2) = d_n. \quad (24)$$

Also $|b_q(n)| \leq C \min(2, T/n)$ and $|b_0(n)| \leq \min(2, T/n)$. Let

$$R_T = \frac{1}{T} \sum_{n \geq 1} f_z(n/T), \quad f_z(x) = x^{2-z} 4 \sin^2(1/(2x)).$$

The discrepancy D_n between the brackets in the exact sum (17) and this limiting mesh is bounded by

$$D_n \leq 2C \min(2, T/n) d_n + \frac{C^2}{n} \min(2, T/n)^2.$$

In particular,

$$D_n \leq \begin{cases} 8C^2/n + 2CT/n^2, & n \leq T, \\ 4C^2T/n^2, & n \geq T. \end{cases} \quad (25)$$

For all n , the separate exact and limiting products give $D_n \leq 12$.

Set $N = \lceil \sqrt{T} \rceil$ and $x_0 = N/T$. Then $\sqrt{T} \leq N \leq (3/2)\sqrt{T}$ and $x_0 \leq 3/4$. The terms $n < N$, using the last bound, contribute at most $12(N/T)^{3-a} \leq 27T^{-(3-a)/2}$. On the range $N \leq n \leq T$, the first and second terms of (25) contribute, respectively, at most

$$\frac{8C^2}{(2-a)T}, \quad \frac{4C}{a-1} T^{-(3-a)/2}.$$

Here we used $\sum_{n \leq T} n^{1-a} \leq T^{2-a}/(2-a)$ and $\sum_{n \geq N} n^{-a} \leq 2N^{1-a}/(a-1)$. The remaining range $n \geq [T]$ contributes at most $8C^2/((a-1)T)$. A duplicated endpoint when T is an integer only enlarges these positive bounds. Thus

$$|A_T - R_T| \leq (27 + 4C/g)T^{-\kappa} + (8C^2/h + 8C^2/g)T^{-1}. \quad (26)$$

For completeness, the whole quadrature error is paid as follows. Writing $D = |2 - z| \leq 2C$, differentiation gives

$$f'_z(x) = (2 - z)x^{1-z} 4 \sin^2(1/(2x)) - 2x^{-z} \sin(1/x),$$

and hence

$$|f'_z(x)| \leq \begin{cases} 4Dx^{1-a} + 2x^{-a}, & x \leq 1, \\ (D+2)x^{-a-1}, & x \geq 1. \end{cases}$$

The head mesh and head integral through x_0 together cost at most $8x_0^{3-a} \leq 18T^{-(3-a)/2}$. On all remaining mesh cells, left-endpoint quadrature costs at most $T^{-1} \int_{x_0}^{\infty} |f'_z|$. Consequently

$$|R_T - H| \leq (18 + 2/g)T^{-\kappa} + (8C/h + 2C + 2)T^{-1}. \quad (27)$$

Since $T^{-1} \leq T^{-\kappa}$, the total constant is at most

$$47 + 2C + (8C^2 + 4C + 2)/g + (8C^2 + 8C)/h \leq C^2(49 + 14/g + 16/h) \leq J.$$

The last inequalities use $C \geq 1$. This proves (23). $\square$

This square-root split and variation estimate extend the diagonal rate argument of [3, Section 5, Remark 4]; the constants above are uniform in both complex slots. The sums at zero and infinity are both included in the estimate.

Proof of (13) and (14). The arithmetic series is locally normally convergent for $\Re q, \Re p > 1/2$. The limiting integral is locally normally convergent when both real parts are in $(1/2, 1)$. Thus $A_T - H$ is holomorphic in q and in the independent variable $\bar{p}$. Cauchy's formula on the two radius- ρ circles proves (13). Apply (19) in both slots. The simplex masses cancel $r!k!$, so the resulting entry bound is $E_0\rho^{-(j+k-2)} = E_0a_ja_k$. For every vector v ,

$$|v^*(G_T - G)v| \leq E_0 \left(\sum_j a_j |v_j| \right)^2 \leq \epsilon \|v\|^2. \quad (28)$$

This proves the stated Loewner comparison without node separation. $\square$

Corollary 5.2 (Adaptive native exponent). *For the rectangle Q of (4), define*

$$\begin{aligned} d &= \min(b_0 - 1/2, 1 - b_1), & g_0 &= 2b_0 - 1, & h_0 &= 2 - 2b_1, \\ \kappa_0 &= 3/2 - b_1, & \bar{C} &= 1 + b_1 + L + d, & \bar{J} &= \bar{C}^2(64 + 32/g_0 + 32/h_0), \\ M(T) &= \max(2/d, \log T). \end{aligned}$$

For every real $T \geq 4$,

$$\sup_{q,p \in Q} |\partial_q^r \partial_{\bar{p}}^k (A_T - H)| \leq 3\bar{J}r!k!T^{-\kappa_0} M(T)^{r+k}. \quad (29)$$

In (14) one may instead take $E_0 = 3\bar{J}T^{-\kappa_0}$ and $a_j = M(T)^{j-1}$.

Proof. Choose $\rho = 1/M(T) = \min(d/2, 1/\log T)$. The expanded rectangle has $g \geq g_0/2$, $h \geq h_0/2$ and $C \leq \bar{C}$, so $J \leq \bar{J}$. Its exponent is $\kappa_0 - \rho$, and $T^\rho \leq e < 3$. Cauchy's formula and the same factorial cancellation give the result. $\square$

6 Logarithmically growing packets

A direct Hardy interpolation argument improves the dimension dependence of the separation-free floor. This permits the packet size to grow with height while retaining the same native kernel and exact target.

Collision-uniform divided-difference bounds for exponential families at fixed cardinality are classical; see Avdonin–Ivanov [9, Lemma 5]. Here an explicit rational Hardy interpolant supplies a dimension-dependent floor that can be combined with the complete arithmetic kernel error.

Lemma 6.1 (A dimension-priced Hardy interpolation floor). *Let $z_1, \dots, z_m$ lie in $\Re z > 0$, with $|z_j| \leq S$. Let $\mathbf{C}(\mathbf{z})$ be the Gram matrix of $F \mapsto [z_1, \dots, z_j]F$, $1 \leq j \leq m$, in the Hardy space with kernel $(z + \bar{w})^{-1}$ and squared boundary norm $(2\pi)^{-1} \int_{\mathbb{R}} |F(it)|^2 dt$. For arbitrary collisions,*

$$\mathbf{C}(\mathbf{z}) \succeq \frac{2}{m(S+2)^{2m}(1+S^2)^{m-1}} I. \quad (30)$$

Proof. Write $L_j F = [z_1, \dots, z_j]F$ and prescribe arbitrary data $b \in \mathbb{C}^m$. Put $d(z) = (z+1)^m$ and define the lower triangular matrix

$$A_{kl} = [z_l, \dots, z_k]d \quad (l \leq k), \quad A_{kl} = 0 \quad (l > k).$$

The divided-difference product rule gives

$$L_k(dF) = \sum_{l=1}^k (L_l F) A_{kl}.$$

This identity follows from the finite recurrence for distinct nodes and extends continuously to every Hermite configuration. Its diagonal entries are $d(z_k) \neq 0$.

For $r = k - l$, the complete-homogeneous formula gives

$$A_{kl} = h_{m-r}(z_l + 1, \dots, z_k + 1), \quad |A_{kl}| \leq \binom{m}{r} (S+1)^{m-r}.$$

Every absolute row sum and column sum is at most $\sum_{r=0}^m \binom{m}{r} (S+1)^{m-r} = (S+2)^m$. The row–column norm bound therefore gives $\|A\| \leq (S+2)^m$. Set $\alpha = Ab$ and form

$$P(z) = \sum_{k=1}^m \alpha_k \prod_{j < k} (z - z_j), \quad F_b(z) = \frac{P(z)}{(z+1)^m}.$$

The Newton data of P are exactly α . Applying the product rule to $dF_b = P$ and using invertibility of A proves $L(F_b) = b$. The rational function F_b is strictly proper with its only pole at -1 , so $F_b \in H^2(\Re z > 0)$.

For real t ,

$$|it - z_j| \leq |t| + S \leq \sqrt{1+S^2} \sqrt{1+t^2}.$$

Consequently, for $1 \leq k \leq m$,

$$\left| \frac{\prod_{j < k} (it - z_j)}{(it+1)^m} \right| \leq \frac{(1+S^2)^{(m-1)/2}}{\sqrt{1+t^2}}.$$

Cauchy–Schwarz in the numerator sum and $\|\alpha\| \leq (S+2)^m \|b\|$ give

$$\|F_b\|_{H^2}^2 \leq \frac{m}{2} (S+2)^{2m} (1+S^2)^{m-1} \|b\|^2,$$

since $(2\pi)^{-1} \int_{\mathbb{R}} (1+t^2)^{-1} dt = 1/2$. Thus $b \mapsto F_b$ is a right inverse R for the Newton data map L , with the displayed squared norm bound. The identity $R^*L^* = I$ implies $\|v\| \leq \|R\| \|L^*v\|$. Since $\|L^*v\|^2 = v^*Cv$, this proves (30), including independence of the functionals, without first assuming that their Gram is invertible. $\square$

Put

$$c_* := \frac{529}{576}, \quad S_Q := \sup_{q \in Q} |q - 1/2|, \quad D_Q := S_Q + 2, \quad C_Q := 1 + S_Q^2.$$

Theorem 6.2 (Exponential packet Gram bound). *For every integer $m \geq 1$ and every tuple in Q^m ,*

$$\mathbf{G}(\mathbf{q}) \succeq \delta_m I, \quad \delta_m := \frac{2c_*}{m D_Q^{2m} C_Q^{m-1}}. \quad (31)$$

Keep the fixed expanded rectangle, radius ρ and literal constants J, κ of (12). Set

$$\beta_\rho := \max(1, \rho^{-2}), \quad \Lambda_Q := D_Q^2 C_Q \beta_\rho, \quad A_Q := \frac{J D_Q^2}{2c_*}.$$

For every real $T \geq 4$, the relative comparison (9) holds with

$$\eta_{T,m} := A_Q m^2 \Lambda_Q^{m-1} T^{-\kappa}. \quad (32)$$

All partial and complete complex collisions are included.

Proof. For $x \geq 1$, the inequality $\sin u \geq u - u^3/6 \geq (23/24)u$ at $u = 1/(2x) \leq 1/2$ gives

$$4x^2 \sin^2(1/(2x)) \geq c_*.$$

Restrict the positive feature integral (21) to $x \geq 1$. It follows, as a positive-kernel comparison, that

$$H(q, p) \succeq_{\text{kernel}} c_* \int_1^\infty x^{-q-\bar{p}} dx = \frac{c_*}{q + \bar{p} - 1}.$$

Divided differences preserve this comparison: apply the same finite linear combination to the features in both slots, and use continuity for repeated nodes. With $z_j = q_j - 1/2$, the right side is the Hardy kernel of Lemma 6.1. That lemma proves (31).

The literal mixed Cauchy bound and the two simplex formulas hold at every finite order. Their factorials cancel exactly, even if the chosen order varies with height. Hence (28) gives

$$|v^*(\mathbf{G}_T - \mathbf{G})v| \leq J T^{-\kappa} \left(\sum_{j=0}^{m-1} \rho^{-2j} \right) \|v\|^2 \leq J m \beta_\rho^{m-1} T^{-\kappa} \|v\|^2.$$

Divide by the floor (31) to obtain (32). Both signs of (9) follow. Positivity of the native Gram follows whenever this explicit error is below one. $\square$

Corollary 6.3 (Schubr's Uniform Growing-Packet Theorem). *For every integer schedule $m(T) \geq 1$ with $m(T) = o(\log T)$, the relative error (32) tends to zero, uniformly over all tuples in $Q^{m(T)}$. More explicitly, fix $c > 0$ with $c \log \Lambda_Q < \kappa$ and set $d = \kappa - c \log \Lambda_Q > 0$. For $1 \leq m(T) \leq c \log T$,*

$$\eta_{T,m(T)} \leq A_Q c^2 (\log T)^2 T^{-d} \longrightarrow 0.$$

For a prescribed $0 < \varepsilon < 1$, a literal sufficient onset is

$$T \geq \max \left(4, e^{1/c}, \left[\frac{16A_Q c^2}{e^2 d^2 \varepsilon} \right]^{2/d} \right). \quad (33)$$

At each such height, if the tuple is a zeta-zero packet with every repetition bounded by the corresponding multiplicity, then

$$D_\infty^2 \geq T^{-3} z_T^* \mathbf{G}_T^{-1} z_T \geq \frac{T^{-3}}{1 + \eta_{T,m(T)}} z_T^* \mathbf{G}^{-1} z_T. \quad (34)$$

Proof. Taking logarithms of (32) proves the sublogarithmic assertion. In the stated linear logarithmic range, $\Lambda_Q^{m-1} \leq T^{c \log \Lambda_Q}$ and $m^2 \leq c^2 (\log T)^2$. The elementary maximum $\sup_{u \geq 0} u^2 e^{-du/2} = 16/(e^2 d^2)$ proves the sufficient onset. For example, $c = \kappa/(2 \log \Lambda_Q)$ is admissible.

The exact gauge functionals (22) have Gram $T^3 \mathbf{G}_T$ and target z_T in every finite dimension. The zero-packet hypothesis makes their representers lie in $V^\perp$. At the stated onset $\eta < 1$, their native Gram is positive definite. The same finite projection argument as in Corollary 1.2, and inversion of its complete upper Gram comparison, prove (34). $\square$

Corollary 6.4 (Sublogarithmic packet budget). *For $m(T) = o(\log T)$, both the scalar relative certificate and the additive-inverse certificate are at most*

$$T^{2b_1 - 3 + o(1)} \rightarrow 0.$$

Thus the fixed finite approximation-space obstruction following (51) continues to hold for these growing packets.

Proof. The exact target estimate gives, in every finite dimension,

$$T^{-3} z_T^* \mathbf{G}^{-1} z_T \leq \frac{m D_Q^{2m} C_Q^{m-1}}{2c_*} T^{2b_1 - 3} \sum_{j=0}^{m-1} \frac{(\log T)^{2j}}{(j!)^2}.$$

For $2 \leq m \leq 1 + \log T$, the sum is at most $m[e \log T / (m-1)]^{2m-2}$, by monotonicity of its terms and $j! \geq (j/e)^j$. When $m = o(\log T)$, the logarithm of the entire extra factor is $o(\log T)$, since $u \log(1/u) \rightarrow 0$ as $u \downarrow 0$. The case $m = 1$ is immediate. The scalar relative factor only decreases this upper, and $(\mathbf{G} + \epsilon I)^{-1} \preceq \mathbf{G}^{-1}$ gives the same ceiling for the additive-inverse certificate. Combine this with the finite head-supported dual lower bound (49) whenever its numerator is nonzero. $\square$

Corollary 6.5 (Approaching-boundary packet theorem). *Fix $L \geq 0$ and $a > 0$. Put $\ell = \log T$, $\delta_T = \ell^{-a}$ and, whenever $\delta_T \leq 1/4$, let*

$$Q_T = \{q : 1/2 + \delta_T \leq \Re q \leq 1 - \delta_T, |\Im q| \leq L\}.$$

The relative native Gram law holds uniformly over all tuples in $Q_T^{m(T)}$ with error tending to zero if

$$m(T) = o\left(\frac{\log T}{\log \log T}\right).$$

It also holds for $1 \leq m(T) \leq c \log T / \log \log T$ whenever $0 < c < 1/(4a)$. At admitted zero packets the same native receiving inequality (34) follows.

Proof. Use the expanded rectangle

$$Q_{e,T} = \{q : 1/2 + \delta_T/2 \leq \Re q \leq 1 - \delta_T/2, |\Im q| \leq L + \delta_T/2\}$$

and radius $\rho_T = \delta_T/2$. The literal scalar error has $g = h = \delta_T$ and $\kappa_T = 1/2 + \delta_T/2$. Its constant satisfies

$$J_T \leq (L+3)^2(64 + 32/\delta_T) \leq \frac{48(L+3)^2}{\delta_T}.$$

The first inequality uses $\sup_{Q_{e,T}} |q| \leq L+2$; the second uses $\delta_T \leq 1/4$. The original-node bound can be kept fixed:

$$S_L = \sqrt{1/4 + L^2}, \quad D_L = S_L + 2, \quad C_L = 1 + S_L^2.$$

Set $\Lambda_L = 4D_L^2 C_L$ and $A_L = 24(L+3)^2 D_L^2 / c_*$. Theorem 6.2, applied separately at each height with its displayed finite constants, gives

$$\eta_{T,m} \leq A_L m^2 \delta_T^{-1} (\Lambda_L \delta_T^{-2})^{m-1} T^{-1/2 - \delta_T/2}. \quad (35)$$

Taking logarithms and discarding the last additional decay gives

$$\log \eta_{T,m} \leq -\frac{1}{2}\ell + a(2m-1) \log \ell + (m-1) \log \Lambda_L + 2 \log m + \log A_L.$$

This tends to $-\infty$ in the sublogarithmic quotient range. For $m \leq c\ell / \log \ell$, it is $[-1/2 + 2ac + o(1)]\ell$, which tends to $-\infty$ under the stated strict condition. This uses the all-order literal error, not a fixed-rectangle asymptotic with changing parameters. The exact finite-dimensional native projection return is unchanged. $\square$

7 A literal floor for every colliding tuple

The compactness proof of Lemma 4.1 does not specify a numerical floor. The following formula supplies one. For $m \geq 1$ and $R > 0$, define

$$\begin{aligned} c_m &= \frac{1}{(2m)^{m-1}} \prod_{k=0}^{m-1} \frac{2^{2k+1}}{(2k+1) \binom{2k}{k}^2}, & b_m &= \frac{1}{\sqrt{m}(1+R)^{m-1}(m-1)!}, \\ K_m(R)^2 &= \sum_{j=0}^{m-1} \left(\frac{R^{m-j}}{j!(m-j)!} \right)^2, & a &= \min \left(\frac{1}{4}, \frac{\sqrt{c_m} b_m}{2\sqrt{2}e^R K_m(R)} \right), \\ \delta_m(R) &= \frac{c_m b_m^2 a^{2m-1}}{8}. \end{aligned} \quad (36)$$

Theorem 7.1 (Explicit separation-free floor). *If $|q_j| \leq R$ and the nodes are in the strip $1/2 < \Re q < 1$, then $\mathbf{G}(\mathbf{q}) \succeq \delta_m(R)I$. The same formula covers arbitrary complex collisions.*

Proof. Index the limiting features by $0 \leq j < m$: $\psi_j(e^y) = [q_1, \dots, q_{j+1}]e^{-qy}$. The complete-homogeneous identity for divided differences gives

$$\psi_j(e^y) = \sum_{k \geq j} \frac{(-y)^k}{k!} h_{k-j}(q_1, \dots, q_{j+1}).$$

For a coefficient vector v , truncate its feature combination at degree $m-1$, obtaining $P_v(y) = \sum_{k < m} (Bv)_k y^k$. Here $B = DH$, where $D_{kk} = (-1)^k / k!$ and $H_{kj} = h_{k-j}(q_1, \dots, q_{j+1})$ for $j \leq k$. The inverse Newton conversion is

$$(H^{-1})_{kj} = (-1)^{k-j} e_{k-j}(q_1, \dots, q_k).$$

Indeed expansion of the Newton products $\prod_{i=1}^j (z - q_i)$ and of $z^k = \sum_{j=0}^k h_{k-j}(q_1, \dots, q_{j+1}) \prod_{i=1}^j (z - q_i)$ proves both conversion formulas. Thus each row of H^{-1} has absolute sum at most $(1 + R)^k$, and $\|B^{-1}\| \leq 1/b_m$.

The monomial moment matrix on $[-1, 1]$ has trace at most $2m$. Its determinant is the product in (36), by monic Legendre orthogonalization: the degree- k Legendre polynomial has leading coefficient $2^{-k} \binom{2k}{k}$ and squared norm $2/(2k + 1)$. Its smallest eigenvalue is therefore at least c_m , by bounding each of the other eigenvalues by the trace. For $0 < a \leq 1$, rescaling $y = at$ yields

$$\|P_v\|_{L^2(-a,a)} \geq \sqrt{c_m} a^{m-1/2} \|Bv\| \geq \sqrt{c_m} b_m a^{m-1/2} \|v\|. \quad (37)$$

Since $|h_{k-j}| \leq \binom{k}{j} R^{k-j}$ and $\sum_{l \geq d} u^l/l! \leq e^u u^d/d!$, the discarded tail of feature j on this interval is at most $e^{Ra} R^{m-j} a^m/(j!(m-j)!)$. The L^2 norm of the combined tail is consequently at most $\sqrt{2} e^R K_m(R) a^{m+1/2} \|v\|$. The choice of a makes this at most half the lower bound (37).

In logarithmic coordinates the positive weight in (21) is $w(y) = 4e^{3y} \sin^2(e^{-y}/2)$. For $|y| \leq 1/4$, $t = e^{-y}/2 \leq 2/3$, $\sin t \geq 5t/6$, and $e^y \geq 3/4$. Hence $w(y) \geq 25/48 > 1/2$. Restricting the integral to $[-a, a]$, and squaring the half-amplitude estimate, proves the stated floor. $\square$

A rational alternative can be useful. For any integer $R \geq 1$ bounding the nodes, put

$$b = \frac{1}{m(1+R)^{m-1}(m-1)!}, \quad d_* = \min(1, c_m), \quad K = \frac{(1+R)^m}{m!}, \quad a_0 = \min\left(\frac{1}{4}, \frac{d_* b}{4 \cdot 3^R K}\right).$$

Then

$$\mathbf{G} \succeq \delta_0 I, \quad \delta_0 = c_m b^2 a_0^{2m-1}/8 > 0. \quad (38)$$

All these constants are rational. To check the same proof, note $b \leq b_m$, $d_* \leq \sqrt{c_m}$, $K_m \leq \sum_j R^{m-j}/(j!(m-j)!) \leq K$, $e^R \leq 3^R$ and $2\sqrt{2} \leq 4$.

Corollary 7.2 (Explicit relative onset). *For any proved floor $\delta > 0$ from the preceding theorem, set*

$$\eta_T = \frac{J \sum_{j=0}^{m-1} \rho^{-2j}}{\delta T^\kappa}.$$

Then (9) holds for all $T \geq 4$. For a prescribed $0 < \eta < 1$, the literal sufficient onset is

$$T \geq \max\left(4, \left[\frac{J \sum_{j=0}^{m-1} \rho^{-2j}}{\eta \delta}\right]^{1/\kappa}\right).$$

The adaptive constants of Corollary 5.2 give the alternative $\eta_T = 3\bar{J}T^{-\kappa_0} \sum_{j=0}^{m-1} M(T)^{2j}/\delta$.

Proof. Apply (28) and $\|v\|^2 \leq v^* \mathbf{G} v/\delta$. Both displayed errors tend to zero for fixed m, Q . $\square$

8 All-height regularized receiving bounds

Proof of (15). The native functionals (22) have Gram $\Gamma = T^3 \mathbf{G}_T$ and target z_T , exactly, at every $T \geq 4$. The zero-packet hypothesis makes their representers r_j belong to $V^\perp$. With the inner product linear in its first argument, $\Gamma_{jk} = \langle r_k, r_j \rangle$. For any vector u set $h = \sum_j u_j r_j$. Positivity of $\|(E - v) - h\|^2$, for $v \in V$, and then taking the infimum give

$$D_\infty^2 \geq 2\Re(u^* z_T) - u^* \Gamma u \geq 2\Re(u^* z_T) - T^3 u^* (\mathbf{G} + \epsilon I) u.$$

The limiting Gram is positive semidefinite by its integral, and $\epsilon > 0$. The concave quadratic on the right is maximized at $u = T^{-3}(\mathbf{G} + \epsilon I)^{-1} z_T$, proving the result. This argument does not invert Γ and remains valid if its functionals are dependent. $\square$

If $\mathbf{G} \succeq \delta I$, then $\mathbf{G} + \epsilon I \preceq (1 + \epsilon/\delta)\mathbf{G}$. Thus the additive-inverse bound is at least as strong as the scalar relative bound with error ϵ/δ . It remains available before the relative comparison establishes positive definiteness by perturbation.

Corollary 8.1 (Weighted and chosen-vector certificates). *For any positive numbers ω_j , put*

$$R_\omega = E_0 \left(\sum_j a_j^2 / \omega_j \right) \text{diag}(\omega_1, \dots, \omega_m).$$

Under the zero-packet hypothesis, at every real $T \geq 4$,

$$D_\infty^2 \geq T^{-3} z_T^* (\mathbf{G} + R_\omega)^{-1} z_T, \quad (39)$$

$$D_\infty^2 \geq T^{-3} \left[2\Re(v^* z_T) - v^* \mathbf{G} v - E_0 \left(\sum_j a_j |v_j| \right)^2 \right] \quad (v \in \mathbb{C}^m). \quad (40)$$

Proof. Weighted Cauchy–Schwarz bounds the first expression in (28) by $v^* R_\omega v$. Repeat the preceding maximization. Retaining the original absolute-value penalty and choosing $u = T^{-3}v$ gives (40) directly. $\square$

The choices $\omega_j = 1$ and $\omega_j = a_j$ are explicit examples. Alternatively, (40) permits a certified test vector without computing an inverse. Every formula retains the exact unequal powers and oscillatory phases in z_T .

9 Phase interference and multiple zeros

For a shared real part, write $q = \beta + i\tau$ and

$$H_\beta(a) = \frac{\pi}{\Gamma(4 - 2\beta - ia) \sin(\pi(3 - 2\beta - ia)/2)}. \quad (41)$$

Then $H(q, p) = H_\beta(\tau - \sigma)$ and $T^q = T^\beta e^{i\tau \log T}$. For distinct offsets τ_j , cancellation of the invertible Newton change of basis in (11) gives

$$D_\infty^2 \geq \frac{T^{2\beta-3}}{1 + \eta_T} z^* \mathbf{H}^{-1} z, \quad \mathbf{H}_{jk} = H_\beta(\tau_j - \tau_k), \quad z_j = e^{i\tau_j \log T}. \quad (42)$$

The relative error is uniform even when the minimum separation tends to zero. At coincident nodes, (11), rather than the inverse of a singular evaluation matrix, defines the endpoint.

The following formula is the elementary two-point Hermitian inversion for the limiting native kernel. Phase-sensitive multi-zero inversion has the preceding weighted-cutoff counterpart [5, Section 4.1].

Corollary 9.1 (Two-point interference). *Put $h = H_\beta(0)$ and $\ell = \log T$. For two distinct zeros $\beta + iT$, $\beta + i(T + \delta)$ in a bounded-offset packet, the limiting factor in (42) is*

$$z^* \mathbf{H}^{-1} z = \frac{2h - 2\Re(H_\beta(-\delta)e^{i\delta\ell})}{h^2 - |H_\beta(\delta)|^2}. \quad (43)$$

The denominator is positive for $\delta \neq 0$.

Proof. The matrix is $\begin{pmatrix} h & H_\beta(-\delta) \\ H_\beta(\delta) & h \end{pmatrix}$, positive definite by independence of its two limiting features. Direct Hermitian inversion gives (43). $\square$

The target phase in (43) is part of the receiving formula. In particular, close distinct zeros are not replaced by a multiple zero merely from their geometric proximity.

For a fixed multiplicity m , set $r = m - 1$ and

$$\mu_j(\beta) = \int_0^\infty (\log x)^j x^{2-2\beta} 4 \sin^2 \frac{1}{2x} dx, \quad M_\beta = (\mu_{a+b}(\beta))_{a,b=0}^r. \quad (44)$$

Every moment exists and M_β is positive definite.

Corollary 9.2 (Multiplicity factor). *Along a sequence of zeros $\rho = \beta + iT$ of multiplicity at least m , with $T \rightarrow \infty$ and β in a fixed compact subinterval of $(1/2, 1)$,*

$$D_\infty^2 \geq (1 - o(1))(M_\beta^{-1})_{r,r} T^{2\beta-3} (\log T)^{2m-2}. \quad (45)$$

The error is uniform in that real-part interval.

Proof. Use the offset variable τ in the chart $q = \beta + i\tau$. Then $\partial_\tau = i\partial_q$; at fully repeated offset zero the Newton–Hermite functionals are $\partial_\tau^a/a!$. Differentiating the limiting feature and target in this chart gives the limiting Gram and target

$$\mathbf{G}_\beta = A M_\beta A^*, \quad A_{aa} = \frac{(-i)^a}{a!}, \quad z_a = \frac{(i\ell)^a}{a!}.$$

Consequently

$$z^* \mathbf{G}_\beta^{-1} z = (M_\beta^{-1})_{r,r} \ell^{2r} + O(\ell^{2r-1})$$

for $m \geq 2$; for $m = 1$ it equals $1/\mu_0(\beta)$. The factorials cancel in the highest-degree term. The moment matrix and its inverse vary continuously on a compact real-part interval and have positive eigenvalue floors there, so the estimate and its positive leading coefficient are uniform. Apply (11) in the shared-beta chart. $\square$

The coefficient $(M_\beta^{-1})_{r,r}$ is also the reciprocal squared norm of the monic degree- r polynomial orthogonal to lower powers for the positive measure in (44). This follows from the Schur complement of the lower-order moment block. When $m = 1$, (45) recovers the sharp single-zero constant of [3, Section 5, Remark 4]. Monic square norms also retain collision geometry in the signed-measure packet estimates of [10]. Here the measure is the native positive logarithmic measure in (44), and the receiving functionals are Mellin jets; the signed-measure recovery theorem is not used as an input.

10 Finite certificates in the same arithmetic norm

The lower bounds above can be compared with a trial approximation only when its full norm is certified in the same space. We give the relevant finite formulas, including their infinite-tail payments.

10.1 A periodic primal residual

In the step realization, $E = 1$ and $G_k|_{I_n} = -\{n/k\}$, as follows by taking the Mellin transform. For rational coefficients b_k let

$$r_n = 1 - \sum_{k=2}^N b_k \{n/k\}, \quad U(b) = \sum_{n \geq 1} \frac{|r_n|^2}{n(n+1)}.$$

Then $D_\infty^2 \leq d_N^2 \leq U(b)$, where $d_N = \text{dist}(E, \text{span}\{G_2, \dots, G_N\})$. For real coefficients the sequence $f_n = r_n^2$ is periodic. Its mean is

$$\begin{aligned} \mu &= 1 - \sum_k b_k \frac{k-1}{k} \\ &\quad + \sum_{j,k} b_j b_k \left[\frac{(j-1)(k-1)}{4jk} + \frac{\gcd(j,k)^2 - 1}{12jk} \right]. \end{aligned} \quad (46)$$

To see the pair formula, average the residues modulo $\text{lcm}(j, k)$; their covariance after scaling by j, k is $(\gcd(j, k)^2 - 1)/(12jk)$. The single means are $(k-1)/(2k)$. This is the periodic-mean mechanism of [4, Theorem 3] in the native norm.

One explicit bound on every interval sum of $f_n - \mu$ is

$$B_f = \sum_k \frac{|b_k| \lfloor k/2 \rfloor \lceil k/2 \rceil}{k} + \sum_{j,k} \frac{|b_j b_k| \text{lcm}(j, k)}{4}. \quad (47)$$

Indeed the centered sawtooth has prefix-sum range $\lfloor k/2 \rfloor \lceil k/2 \rceil / (2k)$. For a sequence in $[0, 1]$ of period P and mean m_0 , the total positive centered mass is at most $Pm_0(1 - m_0) \leq P/4$, bounding its prefix-sum range. Expansion of r_n^2 now proves (47). Abel summation with the decreasing weights $1/(n(n+1))$ yields

$$\left| \sum_{n > M} \frac{f_n}{n(n+1)} - \frac{\mu}{M+1} \right| \leq \frac{B_f}{(M+1)(M+2)}. \quad (48)$$

Thus the finite head, rational mean and tail error supply a full-norm primal certificate without traversing the entire period.

10.2 A finite dual and a continuous-tail majorant

There is also an elementary head-supported annihilator for every finite approximation space. With μ_{Mob} denoting the Möbius function, put

$$t_n = \frac{1}{n} \sum_{r \leq N/n} \frac{\mu_{\text{Mob}}(r)}{r} \quad (1 \leq n \leq N), \quad t_{N+1} = 0, \quad \lambda_n = t_n - t_{n+1}.$$

Möbius inversion gives $\sum_{j \leq N/k} t_{jk} = 1/k$ for $1 \leq k \leq N$. Summation by parts consequently gives

$$\sum_{n \leq N} \lambda_n \lfloor n/k \rfloor = 1/k, \quad \sum_{n \leq N} n \lambda_n = 1, \quad \sum_{n \leq N} \lambda_n \{n/k\} = 0 \quad (2 \leq k \leq N).$$

The functional $L(f) = \sum_{n \leq N} \lambda_n f_n$ annihilates the first- N span and has squared norm $\sum_{n \leq N} n(n+1) \lambda_n^2$. Therefore, for all complex approximation coefficients,

$$d_N^2 \geq H_N := \frac{|\sum_{n \leq N} \lambda_n|^2}{\sum_{n \leq N} n(n+1) \lambda_n^2}. \quad (49)$$

The denominator is nonzero by the normalization identity. A positive lower bound requires a nonzero numerator, which is checked at the chosen N rather than assumed universally.

For a finite point packet $s_1, \dots, s_m$ in Ω , define

$$(A_M)_{jk} = \sum_{n \leq M} n(n+1) e_n(s_j) \overline{e_n(s_k)}, \quad (C_M)_{jk} = \frac{(M+1)^{1-s_j-\bar{s}_k}}{s_j + \bar{s}_k - 1}.$$

The continuous evaluation representers on $(0, 1/(M+1))$ have Gram C_M . Their orthogonal step projections give the remaining native tail Gram. Projection is a contraction for every linear combination, so the exact native Gram satisfies

$$\Gamma \preceq A_M + C_M. \quad (50)$$

For distinct nodes C_M is positive definite, by independence of distinct exponentials on an interval. The variational argument used in Section 8, with target $c_j = 1/s_j$, gives

$$D_\infty^2 \geq c^*(A_M + C_M)^{-1}c$$

whenever the nodes are zeta zeros. Derivative functionals are treated by differentiating the continuous-tail Gram, with the appropriate Hermite factorials. This extends the conjugate-pair projection payment of [4, Theorem 4] by the same Hilbert-space argument.

10.3 The fixed-packet approximation budget

The complex simplex formula gives $||[q_1, \dots, q_{j+1}]T^q| \leq T^{b_1}(\log T)^j/j!$. For a proved floor $G \succeq \delta I$, the scalar relative certificate obeys the uniform upper ceiling

$$\frac{T^{-3}z_T^*G^{-1}z_T}{1 + \eta_T} \leq \delta^{-1}T^{2b_1-3} \sum_{j=0}^{m-1} \frac{(\log T)^{2j}}{(j!)^2} \rightarrow 0. \quad (51)$$

The additive-inverse certificate has the same ceiling, since $(G + \epsilon I)^{-1} \preceq G^{-1}$. Thus any fixed first- N space with $H_N > 0$ eventually cannot produce a full-norm upper smaller than this particular fixed- m , fixed- Q certificate. This follows directly from (49); it is an accounting constraint on these certificates, not an upper bound on D_∞ or on all possible native dual tests.

An exact one-point version is available without an asymptotic error. For $s = \beta + iT$, $1/2 < \beta \leq 1$, $T \geq 1$, let $M = \lceil T \rceil$. On $M \leq n \leq 2M - 1$, $\theta = T \log(1 + 1/n) \leq 1$ and $a = (1 + 1/n)^{-\beta} \geq 1/2$. Since $1 - \cos \theta \geq \theta^2/4$ and $\log(1 + 1/n) \geq 1/(2n)$,

$$|1 - ae^{-i\theta}|^2 \geq T^2/(16n^2), \quad |s|^2 K_S(s, s) \geq \frac{T^{3-2\beta}}{16 \cdot 4^{2\beta}} \geq \frac{T^{3-2\beta}}{256}.$$

The exact native one-point requirement consequently satisfies

$$\frac{1}{|s|^2 K_S(s, s)} \leq 256T^{2\beta-3} \leq 256/T. \quad (52)$$

In particular, if $H_N > 256/T$, no full-norm trial in that fixed approximation space can beat this exact one-point requirement. The comparison always uses the same arithmetic norm: a small head residual alone is not a bound for the full residual.

A complementary whole-line recipient. The accompanying mathematical supplement gives a height-adapted Hardy norm with a complete arithmetic upper, and proves why an unweighted compact boundary band cannot replace a full norm. It includes the exact exterior-weight payment and a prefix-Schur signal identity. Those localized criteria use their own recipients; they are not applications of the native kernel theorem with an upper computed in a different norm.

Algebraic controls. The accompanying standard-library program checks the leading Newton identities at rational complex nodes, mixed and complete collisions, the native-jet factorial convention and the two-point Hermitian inverse identity. Its exact finite controls supplement the analytic proof; the derivative estimates, explicit floor and variational receiver are established above independently of computation. Additional exact controls check the scalar error payments, Newton factorial cancellation, rational floor constants and completed-square identities, including singular Grams.

Statements and Declarations

Funding. This research received no external funding.

Competing interests. The author owns Luminary Oddities and the Luminary Research Archive, which hosts the preprint and its source package. The author declares no other competing interests.

AI assistance. OpenAI ChatGPT and Codex assisted with mathematical derivations, proof development and checking, literature discovery and comparison, verification-code development, manuscript drafting and revision. These tools are not authors. Scott Schubr is the sole author and is responsible for the mathematical content, references, code and final manuscript.

Data and code availability. The source package accompanying the preprint contains the complete proofs, verification programs and input data. <https://research.luminaryoddities.com/collision-uniform-kernel-asymptotics-baez-duarte-zero-packets/v/4>.

References

- [1] L. Báez-Duarte, A strengthening of the Nyman–Beurling criterion for the Riemann hypothesis, *Rend. Lincei Mat. Appl.* **14** (2003), 5–11. <https://arxiv.org/abs/math/0202141>.
- [2] B. Bagchi, On Nyman, Beurling and Baez-Duarte’s Hilbert space reformulation of the Riemann hypothesis, *Proc. Indian Acad. Sci. Math. Sci.* **116** (2006), 137–146. <https://doi.org/10.1007/BF02829783>.
- [3] u/span_of_dilates, *The Báez-Duarte subspace has codimension equal to the number of zeta zeros off the critical line, with distance and rate bounds not assuming RH*, rStation, version 12, 27 October 2026. https://rstation.ai/problems/riemann_hypothesis/papers/1.
- [4] u/halasz_resonator, *Step-Space Annihilator Duality in the Shifted Nyman–Beurling Criterion*, rStation, version 1, 16 October 2026, Theorems 3–4. https://rstation.ai/problems/riemann_hypothesis/papers/9.

- [5] u/halasz_resonator, *Smoothed Biorthogonal Tail Extensions and Multi-Zero Step-Space Superposition in the Shifted Nyman-Beurling Criterion: Two-Sided Divisor-Sum Operator Bounds, Cutoff Jump Elimination, and Interval-Certified Collective Zero Exclusion*, rStation, version 1, 17 October 2026.
https://rstation.ai/problems/riemann_hypothesis/papers/11.
- [6] J. Manzur and W. Noor, *A Hardy-space operator acting on ζ -zeros*, arXiv:2610.03568v1 (2026). <https://arxiv.org/abs/2610.03568v1>.
- [7] J. Manzur, W. Noor and G. Quintero, *A Hardy space approximation supporting zero-free half-planes for the ζ -function*, arXiv:2606.16097v1 (2026).
<https://arxiv.org/abs/2606.16097v1>.
- [8] u/bramble_trace, *Adaptive zero clusters and optimal Nyman-Beurling approximation rates*, rStation, version 2, 24 October 2026.
https://rstation.ai/problems/riemann_hypothesis/papers/48.
- [9] S. A. Avdonin and S. A. Ivanov, Exponential Riesz bases of subspaces and divided differences, *Algebra i Analiz* **13**(3) (2001), 1–17; English translation, *St. Petersburg Math. J.* **13**(3) (2002), 339–351. <https://arxiv.org/abs/math/0103160v1>.
- [10] S. Schubr, *Growing-Index Recovery and Tail Scales for Signed Measures*, author preprint (2026). <https://research.luminaryoddities.com/growing-index-recovery-tail-scales-signed-measures/v/1>.