

Fourier computation of empirical diffusion metrics

Scott Schubr

Luminary Oddities

Countryside, Illinois, United States

Corresponding author: `scottschubr@luminaryoddities.com`

ORCID: <https://orcid.org/0009-0009-7357-4439>

October 30, 2026

Abstract

We give a finite Fourier algorithm for computing empirical diffusion-metric entries together with their joint uncertainty estimates. The metric is defined through an inverse weighted Laplacian and estimated by empirical heat-kernel Sinkhorn normalization. For a fixed smooth, strictly positive density on a fixed-dimensional flat torus and fixed smooth test functions, the algorithm preserves the exact estimator’s population-centered Gaussian limit, joint covariance consistency and ordinary nonparametric bootstrap. For every fixed $\eta > 0$, its deterministic work is $O(n^{1+\eta})$ in the exact-real model, using sufficiently high fixed-order bias extrapolation. The proof controls the closing spectral gap as bandwidth decreases, the actual normalization iterations, and the errors in both the inverse metric and its influence covariance under the original empirical weights. Positive Fourier features and reduced linear solves provide the algorithm. The work bound includes feature-Gram construction and factorization; explicit finite schedules illustrate the resulting resource counts.

Keywords: Sinkhorn normalization, diffusion metric, Fourier features, statistical computation, influence covariance.

Mathematics Subject Classification (2020): 62G20, 65C60, 65T50.

1 Introduction

An empirical diffusion metric involves both kernel normalization and an inverse whose nonconstant spectral gap closes as the bandwidth tends to zero. Fast kernel multiplication alone therefore gives an incomplete computational guarantee. For statistical inference, the numerical metric must approximate its exact counterpart more accurately than the root-sample fluctuation, and the influence calculation must retain the response of the normalization and the inverse. A small error in the metric does not by itself control these quadratic covariance calculations.

We resolve this computational question for the periodic heat-kernel Sinkhorn metric and the population-centered estimator of [7]. The algorithm forms positive finite Fourier features, performs a prescribed logarithmic number of half-scaling iterations from the constant vector, and solves reduced feature systems. It computes metric entries and their complete joint influence covariance under the original empirical law. The main theorem gives the total operation count, including feature Gram formation and factorization, and returns the inferential limits of the exact estimator. The same construction applies to a fixed ordinary bootstrap replicate.

The operation bound is pointwise at each fixed positive smooth density. For every fixed $\eta > 0$, sufficiently high but fixed extrapolation order permits a bandwidth for which the work is $O(n^{1+\eta})$. All constants may depend on the density, dimension, prescribed scores, extrapolation order and chosen exponent. We count real arithmetic and elementary function evaluations at unit cost. This declared model specifies the theorem’s computational guarantee.

1.1 Relation to previous work

The symmetry-preserving geometric half iteration is classical; see Knight–Ruiz–Uçar [4, Section 3]. Symmetric Sinkhorn normalization as a diffusion-kernel construction is developed by Kessler–Magnet–Feydy [3]. We use this established normalization and bound the number of iterations on the actual sampled trajectory uniformly as the bandwidth shrinks. Wormell–Reich [8, Section 4] already establish a well-conditioned accelerated symmetric Sinkhorn algorithm and small-bandwidth initialization on the flat torus. Their local contraction and initialization results already give fast normalization in this setting. Here the prescribed half iteration is analyzed together with finite Fourier approximation and the full inverse-metric and influence-covariance return. Fourier acceleration of Sinkhorn iterations is established; Lakshmanan–Pichler–Potts [5] develop an NFFT implementation, while Altschuler–Bach–Rudi–Niles-Weed [1] combine low-rank Nyström approximation with scaling stability and explicit transport-cost guarantees. We use finite Fourier features and the Woodbury identity. The additional issue here is the simultaneous shrinking bandwidth, inverse metric, actual iteration count and original-law influence covariance.

The exact statistical target and its root- n inference are proved in [7]. We state the specific inputs needed from that paper before bounding the numerical errors. Differentiability and inference for fixed-regularization Sinkhorn geometric operations are developed by Saidi–Gunsilius–Mena [6]. Dong et al. [2] study sensitivity and conditioning for finite-support inverse cost parameters observed through transition plans. These provide related operator and statistical perspectives; the present theorem concerns fixed-score inverse diffusion energies of an unknown sampled density as the heat bandwidth vanishes.

The proof has three computational obligations. First, a feature-covariance comparison gives an empirical gap of order bandwidth. Second, positive semidefiniteness gives local contraction of the actual half iterations, after one smoothing step has entered a small normalization neighborhood.

Third, direct original-law product estimates pay the complete inverse response and covariance, including their mixed terms. Sections 3–5 carry out these steps.

2 Target, exact estimator and main theorem

We first define the population metric and its exact sample estimator. The main theorem then states which statistical conclusions survive the finite computation and the work required to preserve them. Let $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$ have volume one. Fix a strictly positive $f \in C^\infty(\mathbb{T}^d)$ with $\int f = 1$, put $\mu = f \, dx$, and fix real $u_1, \dots, u_p \in C^\infty(\mathbb{T}^d)$. Dimension d and the number p of scores are fixed. If $u_a^0 = u_a - \mu u_a$, define

$$D_f = -f^{-1} \operatorname{div}(f \nabla), \quad D_f \phi_a = u_a^0, \quad \mu \phi_a = 0, \quad G_{ab}^0 = \mu(u_a^0 \phi_b). \quad (2.1)$$

The prescribed metric is equivalently $G_{ab}^0 = \mu(\nabla \phi_a \cdot \nabla \phi_b)$. Its influence and joint covariance are

$$\begin{aligned} \psi_{ab}^0 &= u_a^0 \phi_b + u_b^0 \phi_a - \nabla \phi_a \cdot \nabla \phi_b - G_{ab}^0, \\ \Sigma_{ab,cd} &= \mu(\psi_{ab}^0 \psi_{cd}^0), \quad a \leq b, \, c \leq d. \end{aligned} \quad (2.2)$$

For $t > 0$, let $H_t = e^{t\Delta/2}$ and write its periodic kernel as

$$k_t(x) = \sum_{r \in \mathbb{Z}^d} e^{-2\pi^2 t |r|^2} e^{2\pi i r \cdot x}. \quad (2.3)$$

At a probability law ν , the positive symmetric scale and Markov operator satisfy

$$a_{t,\nu} H_t(a_{t,\nu} \nu) = 1, \quad P_{t,\nu} v = a_{t,\nu} H_t(a_{t,\nu} v \nu). \quad (2.4)$$

For finite positive laws the scale is the unique finite Sinkhorn scale. Let $\Pi_\nu v = (\nu v)1$ and put

$$T_{t,\nu} = \frac{t}{2} \left\{ (I - P_{t,\nu} + \Pi_\nu)^{-1} - (I + P_{t,\nu})^{-1} - \frac{\Pi_\nu}{2} \right\}, \quad G_{ab}(t, \nu) = \nu(u_a T_{t,\nu} u_b). \quad (2.5)$$

Thus $T1 = 0$ and, on mean-zero functions, $T = tP(I - P^2)^{-1}$.

For an integer $s > d/2$ define

$$M = 2 + \frac{5s}{2}, \quad 0 < \beta < \min \left\{ \frac{1}{2M}, \frac{\eta}{4d} \right\}, \quad 2K > M, \quad \frac{1}{4K} < \beta, \quad (2.6)$$

where K is an integer and $\eta > 0$ is prescribed. Such choices exist: choose β first and then increase K . For $n \geq 2$ set $h_n = n^{-\beta}$ and $t_\ell = h_n/2^\ell$, $0 \leq \ell < K$. The fixed extrapolation weights are determined by

$$\sum_{\ell=0}^{K-1} c_\ell = 1, \quad \sum_{\ell=0}^{K-1} c_\ell 4^{-\ell j} = 0 \quad (1 \leq j < K). \quad (2.7)$$

Let $X_1, \dots, X_n$ be iid from μ and $\mu_n = n^{-1} \sum_i \delta_{X_i}$. The exact estimator is

$$\widehat{G}_n = \sum_{\ell=0}^{K-1} c_\ell G(t_\ell, \mu_n). \quad (2.8)$$

Theorem 2.1 (Computational inference). *Under the fixed model above, the algorithm in Section 3 returns a matrix $\widetilde{G}_n$, a joint covariance matrix $\widetilde{\Sigma}_n$ and a failure flag. Its deterministic work on every sample is*

$$O(nR_n^2 + R_n^3 + nR_n \log n) = O(n^{1+\eta}), \quad R_n = O(h_n^{-d/2} (\log n)^{d/2}), \quad (2.9)$$

and its memory is $O(nR_n + R_n^2)$. Real arithmetic, elementary exponentials, logarithms, trigonometric functions, square roots and prescribed score evaluations have unit cost. At the fixed density f ,

the probability of a zero failure flag tends to one, and

$$\sqrt{n} \|\tilde{G}_n - \hat{G}_n\| \longrightarrow 0, \quad \tilde{\Sigma}_n \longrightarrow \Sigma \quad \text{in probability.} \quad (2.10)$$

Consequently

$$\sqrt{n} \operatorname{vec}(\tilde{G}_n - G^0) \Longrightarrow N(0, \Sigma), \quad (2.11)$$

where vectorization uses entries $a \leq b$. Every fixed contrast with positive limiting variance admits the corresponding asymptotically standard normal studentization.

For one ordinary nonparametric bootstrap replicate, use the same bandwidths, weights and iteration budget, merging repeated sites. If $\tilde{G}_n^*$ is its output, then

$$d_{\text{BL}} \left(\mathcal{L}[\sqrt{n} \operatorname{vec}(\tilde{G}_n^* - \tilde{G}_n) \mid X_1, \dots, X_n], N(0, \Sigma) \right) \longrightarrow 0 \quad (2.12)$$

in probability. Each replicate has the work bound (2.9).

The theorem fixes p, K, η before n varies. The covariance may be singular; studentization is asserted for fixed positive-variance contrasts. The implementation uses sampled positions and score values, with its bandwidth and budgets set in advance.

2.1 Statistical inputs from the exact estimator

We use the following results from [7, Sections 3–7]. They identify the statistical problem to which the computational return will be applied.

Proposition 2.2 (Exact-estimator inputs). *For the fixed model and choices (2.6):*

1. *The population scale obeys $a_{t,\mu} = f^{-1/2} + O(t)$ uniformly. Its normalized positive selfadjoint Markov operator has a nonconstant $L^2(\mu)$ gap at least ct for small t .*

2. With $q = 1 + s/2$ and $\epsilon = \|\nu - \mu\|_{H^{-s}} \leq ct^{s+q}$,

$$\|a_{t,\nu}/a_{t,\mu} - 1\|_{H^s} \leq C\epsilon t^{-s}. \quad (2.13)$$

The local scale at a positive atomic law agrees with its finite scale.

3. The exact estimator (2.8) has asymptotic linear representation with influence (2.2) and covariance Σ , and its exact multinomial bootstrap is consistent. For a scalar score u , let $g = u - \nu u$ and

$$z_- = (I - P_\nu)^{-1}g, \quad \nu z_- = 0, \quad z_+ = (I + P_\nu)^{-1}g, \quad C_\nu = (I - P_\nu)(I + P_\nu)^{-1}. \quad (2.14)$$

The exact centered influence is the centered version of

$$\Psi_{t,\nu} = \frac{t}{2} \{g(z_- - z_+) + C_\nu(z_- P_\nu z_- + z_+ P_\nu z_+)\}. \quad (2.15)$$

Products here are pointwise. At the base bandwidth h_n , polarization of these exact empirical influences has a consistent joint covariance under μ_n ; each fixed influence has $L^2(\mu_n)$ norm $O_{\mathbb{P}}(1)$.

The local neighborhood is reached because $\mathbb{E}\|\mu_n - \mu\|_{H^{-s}}^2 = O(n^{-1})$ and $M > s + q$. These inputs retain the simultaneous vanishing bandwidth and the moving empirical normalization. The remaining proof concerns the finite algorithm and its error relative to these exact quantities.

3 A finite algorithm

We describe one band t . The same construction is used at all K bands. Merge repeated sample sites and write the resulting law as $\nu = \sum_{i=1}^m w_i \delta_{x_i}$, with $m \leq n$ and positive integer multiples w_i of $1/n$. Put $W = \text{diag}(w_i)$, $e = (\sqrt{w_i})_{i=1}^m$ and $\mathcal{P} = I - ee^T$.

3.1 Positive normalized features

Fix $A = 30$, $B = 20$ and $C_0 = A + d + 1$. Let

$$L = \left\lceil \sqrt{C_0 \log n/t} \right\rceil, \quad a = 2\pi^2 t, \quad E_t = d(1 + \sqrt{\pi/a})^{d-1} \frac{e^{-aL^2}}{aL}. \quad (3.1)$$

Let $k_{t,L}$ be (2.3) restricted to $|r_j| \leq L$ for all j , and define

$$q_t = \frac{k_{t,L} + E_t + n^{-A-2}}{1 + E_t + n^{-A-2}}. \quad (3.2)$$

Its real sine-cosine features give an $m \times R$ matrix F such that $q_t(x_i - x_j) = (FF^T)_{ij}$. The constant feature is one. For one representative of each pair $\{r, -r\}$ of nonzero retained frequencies, the two other features are

$$\sqrt{\frac{2e^{-2\pi^2 t|r|^2}}{1 + E_t + n^{-A-2}}} \cos(2\pi r \cdot x), \quad \sqrt{\frac{2e^{-2\pi^2 t|r|^2}}{1 + E_t + n^{-A-2}}} \sin(2\pi r \cdot x).$$

3.2 Scaling and projected solves

Use the symmetry-preserving half iteration [4], starting at $b_0 = 1$, and perform exactly

$$J = 1 + \left\lceil \frac{(B+2) \log n}{\log(4/3)} \right\rceil, \quad b_{j+1} = \left\{ \frac{b_j}{F[F^T(w b_j)]} \right\}^{1/2} \quad (0 \leq j < J), \quad (3.3)$$

componentwise. Form

$$V_0 = W^{1/2} \text{diag}(b_J)F, \quad V = \mathcal{P}V_0, \quad H = V^T V, \quad S = VV^T. \quad (3.4)$$

Cholesky-factor $I - H$ and $I + H$. If $I - H$ is not positive definite, return zero metric and covariance matrices with a nonzero failure flag. Otherwise use

$$\tilde{R}_- = I + V(I - H)^{-1}V^T, \quad \tilde{R}_+ = I - V(I + H)^{-1}V^T, \quad \tilde{C} = 2\tilde{R}_+ - I - ee^T. \quad (3.5)$$

Only their actions are formed. The last projector makes $\tilde{C}e = 0$.

For each scalar score needed in polarization, compute

$$\begin{aligned} g &= u - \nu u, & g_E &= W^{1/2}g, \\ z_{-,E} &= \tilde{R}_- g_E, & z_{+,E} &= \tilde{R}_+ g_E, & z_{\pm} &= W^{-1/2}z_{\pm,E}, \\ \tilde{\Phi}_t(u) &= \frac{t}{2}g_E^T(z_{-,E} - z_{+,E}), \\ U &= z_-(W^{-1/2}S z_{-,E}) + z_+(W^{-1/2}S z_{+,E}), \\ \tilde{\psi}_E(u) &= \mathcal{P} \frac{t}{2} \left\{ W^{1/2}[g(z_- - z_+)] + \tilde{C}W^{1/2}U \right\}. \end{aligned} \quad (3.6)$$

All products in U and in the square brackets are products of function values. Thus the conversion by $W^{-1/2}$ occurs before multiplication.

Polarize with $(u_a + u_b, u_a - u_b)$, dividing the differences by four, to obtain $\tilde{G}_{ab}(t, \nu)$ and $\tilde{\psi}_{ab,E}(t, \nu)$.

Combine the metrics with (2.7). At the base band h_n form

$$\tilde{\Sigma}_{ab,cd} = \tilde{\psi}_{ab,E}^T \tilde{\psi}_{cd,E}. \quad (3.7)$$

This is the original-weight empirical covariance, because the Euclidean influence vectors include $W^{1/2}$. A one-site law has centered inputs zero and returns zero quantities by these same formulas.

4 Gap and scaling bounds

The inverse calculation needs a lower bound on the sample spectral gap. The finite algorithm also needs a prescribed number of normalization iterations to reach sufficient accuracy. We prove these two estimates after bounding the Fourier approximation error.

Lemma 4.1 (Feature accuracy). *For a sufficiently large fixed C_0 , the kernel (3.2) is real, symmetric, strictly positive, positive semidefinite and has integral one. Uniformly over the fixed bands,*

$$R = O(t^{-d/2}(\log n)^{d/2}), \quad \|q_t - k_t\|_\infty \leq Cn^{-A}. \quad (4.1)$$

Proof. For $a > 0$, monotonicity and a Gaussian integral bound give

$$\sum_{j \in \mathbb{Z}} e^{-aj^2} \leq 1 + \sqrt{\pi/a}, \quad 2 \sum_{j > L} e^{-aj^2} \leq \frac{e^{-aL^2}}{aL}.$$

A union bound over the coordinates therefore makes E_t a bound for the discarded Fourier absolute sum. The added constant makes the retained series strictly positive since $k_t > 0$. All its Fourier coefficients are nonnegative, giving positive semidefiniteness. The denominator fixes the constant coefficient to one. Also $\|k_{t,L}\|_\infty \leq Ct^{-d/2}$; hence the normalization error is bounded by a constant times $(E_t + n^{-A-2})(1 + t^{-d/2})$. The choice $C_0 = A + d + 1$ gives $E_t \leq Ct^{-d/2}n^{-2\pi^2 C_0}$; using $\beta d < 1$ proves (4.1). Counting the retained cube gives the rank bound. $\square$

Lemma 4.2 (Empirical gap). *With probability tending to one, the exact empirical normalized heat operator at every fixed band has a nonconstant $L^2(\mu_n)$ gap at least $ct/2$, for a constant $c > 0$*

depending on the fixed model.

Proof. Write $a_t = a_{t,\mu}$ and $a_{t,n} = a_{t,\mu_n}$. Sobolev embedding and (2.13) give

$$\|a_{t,n}/a_t - 1\|_\infty = O_{\mathbb{P}}(n^{-1/2}t^{-s}). \quad (4.2)$$

Use the heat feature vector $\varphi_t(x) = (e^{-\pi^2 t |r|^2} e^{2\pi i r \cdot x})_r \in \ell^2$ and put $g_t(x) = a_t(x)\varphi_t(x)$. The covariance $C_t = \mathbb{E}[g_t(X) \otimes g_t(X)]$ has the population Markov operator's nonzero spectrum. Its largest eigenvalue is one and its second is at most $1 - ct$, by Proposition 2.2. Since $\|g_t(x)\|^2 \leq Ck_t(0) \leq Ct^{-d/2}$, Hilbert-space variance gives

$$\mathbb{E}\|C_{t,n} - C_t\|_{\text{HS}}^2 \leq Cn^{-1}t^{-d}, \quad C_{t,n} = n^{-1} \sum_i g_t(X_i) \otimes g_t(X_i).$$

The nonzero spectrum of $C_{t,n}$ is that of the sampled Gram matrix with population scales. Replacing them by $a_{t,n}$ is a diagonal congruence by $I + O_{\mathbb{P}}(n^{-1/2}t^{-s})$. Its norm error is of the same order, and $s > d/2$ absorbs the covariance error. Weyl comparison thus gives

$$\lambda_2(P_{t,\mu_n}) \leq 1 - ct + O_{\mathbb{P}}(n^{-1/2}t^{-s}).$$

The exact normalized matrix is a positive selfadjoint Markov matrix. Its constant eigenvalue is exactly one. Finally $\beta(s+1) < 1/2$, so the last error is $o_{\mathbb{P}}(t)$. A fixed union over the bands proves the assertion. $\square$

Lemma 4.3 (Actual scaling trajectory). *For the fixed iteration count (3.3), uniformly over the fixed bands,*

$$\|\log b_J - \log a_{t,n}\|_\infty = O_{\mathbb{P}}(n^{-B}). \quad (4.3)$$

Proof. First use the exact heat matrix K_t with the empirical weights. In logarithmic coordinates its half map is $\mathcal{F}_k(u) = \{u - \log(K_t(we^u))\}/2$. It is sup-norm nonexpansive: its derivative is half of identity minus a stochastic matrix and has row absolute sums at most one. If $u_* = \log a_{t,n}$, heat smoothing $H^{-s} \rightarrow H^s \rightarrow C^0$ and (4.2) show that its first iterate from zero satisfies

$$\|u_1 - u_*\|_\infty = O_{\mathbb{P}}(t + n^{-1/2}t^{-s}) = o_{\mathbb{P}}(1).$$

Both u_1 and u_* approach $-\log f/2$. The whole later trajectory stays in the same sup-norm box about u_* .

Let P_* be the exact normalized operator in $L^2(w)$. At $u = u_* + v$, the stochastic matrix in the derivative is

$$P_u = \text{diag}((P_*e^v)^{-1})P_*\text{diag}(e^v).$$

Positive semidefiniteness and the Markov property give $\|I - P_*\|_{\text{op}} \leq 1$. For $\|v\|_\infty \leq r$,

$$\|P_u - P_*\|_{\text{op}} \leq e^{2r} - 1, \quad \|D\mathcal{F}_k(u)\|_{\text{op}} \leq e^{2r}/2.$$

Choose a fixed small r for which the last bound is at most $3/4$. The straight segment to u_* lies in the box, so the actual iterates contract in $L^2(w)$ there. Since $w_i \geq 1/n$,

$$\|u_J - u_*\|_\infty \leq \sqrt{n}r(3/4)^{J-1} = O(n^{-B}).$$

The fixed iteration count is independent of the density-dependent asymptotic onset of this event.

Now compare the actual finite-kernel map $\mathcal{F}_q$ with $\mathcal{F}_k$ along the exact-kernel trajectory. All scaling vectors on that trajectory are bounded above and below on the good event, and $H_t\mu_n$ is uniformly bounded below. Equation (4.1) therefore gives $\|\mathcal{F}_q(u_j) - \mathcal{F}_k(u_j)\|_\infty \leq Cn^{-A}$. The finite-

kernel map is also sup-norm nonexpansive. Induction over all actual steps yields path difference at most CJn^{-A} . Since $A > B + 3$, this proves (4.3). $\square$

5 Inverse and covariance error bounds

We now propagate the preceding errors through the inverse operators and the influence calculation. The covariance requires products of influence functions, so both linear cross errors and their product must be retained. All comparisons use the original empirical weights.

Lemma 5.1 (Reduced inverse accuracy). *On an event of probability tending to one, the Cholesky test succeeds at every band. In original function coordinates, write $P_0 = P_* - \Pi_w$ and use S for its trial counterpart under the isometry $W^{1/2}$. For $\delta_0 = Cn^{-B}$,*

$$\begin{aligned} \|S - P_0\|_{\text{op}} &\leq \delta_0, \\ \|\tilde{R}_- - R_-\|_{\text{op}} &\leq C\delta_0 t^{-2}, \quad \|\tilde{R}_+ - R_+\|_{\text{op}} \leq C\delta_0, \\ \|\tilde{C} - C_*\|_{\text{op}} &\leq C\delta_0. \end{aligned} \tag{5.1}$$

The resolvents in the second line are compared on centered inputs; the response in the last line vanishes on constants.

Proof. Before projection, the finite scaled Gram equals DP_*D plus a kernel error, with $D = \text{diag}(b_J/a_{t,n}) = I + O_{\mathbb{P}}(n^{-B})$. The bounded scales and the weighted Schur test bound that kernel error by Cn^{-A} . Orthogonal projection has norm one, giving the first line. Also $S \succeq 0$, and Lemma 4.2 gives $\|S\| \leq 1 - ct/4$ on the good event. Nonzero eigenvalues of VV^T and V^TV agree, so $I - H$ is positive definite. The Woodbury identities (3.5) are therefore valid.

The resolvent identity pays two minus inverses, each of norm $O(t^{-1})$, and two bounded plus inverses. On the mean-zero space this proves the second line of (5.1). The exact zero-on-constants

response is $2(I + P_0)^{-1} - I - \Pi_w$; hence its error is bounded by the plus-resolvent error. This also verifies the last line. $\square$

Lemma 5.2 (Metric and full influence errors). *At each band, for every fixed scalar score and its centered exact and trial influences,*

$$\begin{aligned} |\tilde{\Phi}_t - \Phi_t| &\leq C\delta_0/t, \\ \|\tilde{\psi}_t - \psi_t\|_{L^2(w)} &\leq C\sqrt{n}\delta_0 t^{-2} \end{aligned} \tag{5.2}$$

on the good event. The resulting original-law joint covariance error is $o_{\mathbb{P}}(1)$.

Proof. The metric bound follows from the prefactor $t/2$, a bounded centered score and the minus-resolvent error. For the influence write all quantities in $L^2(w)$. Exact and trial minus outputs have norm $O(t^{-1})$, their difference is $O(\delta_0 t^{-2})$, and plus outputs are bounded with difference $O(\delta_0)$. For every vector $\|z\|_{\infty} \leq \sqrt{n}\|z\|_{L^2(w)}$.

For pointwise products the exact identity

$$zPz - xSx = (z - x)Pz + xP(z - x) + x(P - S)x$$

pays both output changes and the operator change. Applied to $U = z_- P_* z_- + z_+ P_* z_+$ and its trial version it gives

$$\|U - \tilde{U}\|_2 \leq C\sqrt{n}\delta_0 t^{-3}, \quad \|U\|_2 + \|\tilde{U}\|_2 \leq C\sqrt{n}t^{-2}.$$

Together with (5.1) and the prefactor $t/2$, this proves the second line of (5.2). Centering at the same law is an $L^2(w)$ contraction.

The exact influence norms are $O_{\mathbb{P}}(1)$ by Proposition 2.2. For two scalar or polarized entries,

the covariance difference is bounded by

$$\|\tilde{\psi}_a - \psi_a\|_2 \|\psi_b\|_2 + \|\tilde{\psi}_b - \psi_b\|_2 \|\psi_a\|_2 + \|\tilde{\psi}_a - \psi_a\|_2 \|\tilde{\psi}_b - \psi_b\|_2.$$

This includes the two linear mixed terms and their product. Since $\sqrt{n} n^{-B} h_n^{-2} \rightarrow 0$, it proves covariance consistency after the fixed polarization. Every pairing uses the original law w . $\square$

Proof of Theorem 2.1. The fixed multiband metric error is $O_{\mathbb{P}}(n^{-B} h_n^{-1})$ by Lemma 5.2, so it is $o_{\mathbb{P}}(n^{-1/2})$. The same lemma and the exact covariance input prove (2.10). Failure has probability tending to zero by Lemma 5.1, so the specified fallback leaves these convergences unchanged. The exact estimator's asymptotic linear representation and Slutsky's lemma give (2.11) and the stated studentization.

For completeness the operation budget is

Operation at one band	Work
Real feature formation and storage	$O(nR)$
J actual scaling multiplications	$O(nR \log n)$
Projection and feature Gram $V^T V$	$O(nR^2)$
Two reduced Cholesky factorizations	$O(R^3)$
Fixed metric and influence right-hand sides	$O(nR + R^2)$
Fixed joint covariance	$O(n)$

The projection is performed as $V_0 - e(e^T V_0)$, and S is applied as $V(V^T x)$; neither needs a dense sample-by-sample matrix. There are only K fixed bands. The stated memory and deterministic work bounds follow, including on fallback samples. Now $nR_n^2 = O(n^{1+\beta d}(\log n)^d)$ and $R_n^3 = O(n^{3\beta d/2}(\log n)^{3d/2})$. Equation (2.6) absorbs both powers and logarithms into $O(n^{1+\eta})$.

For a bootstrap law μ_n^* , conditional Hilbert-space variance and the original empirical bound

give $\mathbb{E}\|\mu_n^* - \mu\|_{H^{-s}}^2 \leq C/n$. The heat feature-covariance discrepancy likewise has joint squared mean $O(n^{-1}t^{-d})$: condition on the original sample for the bootstrap variance and add the original sampling variance. Every preceding good-event argument therefore holds under the joint original/bootstrap law. Merged weights remain at least $1/n$. Consequently $\sqrt{n}(\tilde{G}_n^* - \hat{G}_n^*) \rightarrow 0$ jointly in probability, as does the original numerical error. For each $\epsilon > 0$, Markov's inequality applied to the conditional error probability turns this into conditional negligibility in original-sample probability. The exact bootstrap input then gives (2.12). Each replicate repeats the same finite budget. $\square$

5.1 Inference from the computed covariance

The full covariance output permits fixed-dimensional joint Gaussian calibration after the metric computation. This is a standard consequence of covariance consistency; it makes explicit what can be done with the computed joint influence covariance.

Corollary 5.3 (Fixed-contrast calibration). *Write $q = p(p+1)/2$ and fix a real $k \times q$ matrix A . Under Theorem 2.1, set $C_n = A\tilde{\Sigma}_n A^T$ and $C = A\Sigma A^T$. Given the observations, let $Z_n^* = C_n^{1/2}g$, where $g \sim N(0, I_k)$ is independent. Then*

$$d_{\text{BL}}(\mathcal{L}[Z_n^* \mid X_1, \dots, X_n], N(0, C)) \longrightarrow 0 \quad \text{in probability.}$$

This statement allows singular C . If C is positive definite, put $Y_n = \sqrt{n}A \text{vec}(\tilde{G}_n - G^0)$. With probability tending to one C_n is invertible, and

$$Y_n^T C_n^{-1} Y_n \Longrightarrow \chi_k^2,$$

where the statistic is defined as zero when C_n is singular. From the computed covariance, forming C_n costs $O(kq^2 + k^2q)$ operations, and factoring it costs $O(k^3)$ exact-real operations. Each subsequent

calibration vector uses $O(k^2)$ operations given an independent standard Gaussian vector. For fixed p, k , these costs are independent of sample size.

Proof. The matrix $\tilde{\Sigma}_n$ is a Gram matrix, including on the failure branch, so C_n is positive semidefinite. Theorem 2.1 gives $C_n \rightarrow C$ in probability. Continuity of the positive semidefinite square root and coupling with the same g bound the bounded-Lipschitz distance by $\min\{2, \mathbb{E}\|g\| \|C_n^{1/2} - C^{1/2}\|_{\text{op}}\}$, which tends to zero in probability. When C is positive definite, continuity of inversion and the theorem's Gaussian limit give the chi-square assertion by Slutsky's theorem.

A semidefinite Cholesky algorithm forms a factor L_n with $L_n L_n^T = C_n$ using arithmetic, square roots and exact zero comparisons. At a zero pivot the corresponding row and column of the positive semidefinite Schur complement are zero, so the algorithm fills that factor column with zeros and proceeds. This costs $O(k^3)$, also for singular matrices. Multiplication $L_n g$ costs $O(k^2)$ and has the same conditional Gaussian law as $C_n^{1/2} g$. $\square$

6 Examples and computational resources

The uniform circle gives a direct normalization example. Let $d = 1$, $f = 1$ and $u(x) = \cos(2\pi x)$.

Then $D_f = -\partial_x^2$, $\phi = u/(4\pi^2)$ and

$$G^0 = \frac{1}{8\pi^2}, \quad \psi^0(x) = \frac{3 \cos(4\pi x)}{8\pi^2}, \quad \Sigma = \frac{9}{128\pi^4}. \quad (6.1)$$

Indeed $\mu u^2 = 1/2$ and $2u\phi - |\phi'|^2 - G^0 = 3(2 \cos^2(2\pi x) - 1)/(8\pi^2)$. At finite population bandwidth,

$P_t u = e^{-2\pi^2 t} u$ and hence

$$G(t, \mu) = \frac{t}{4 \sinh(2\pi^2 t)}. \quad (6.2)$$

This gives the target, covariance and even-bias normalization in closed form.

The accompanying Python reference implementation follows (3.2)–(3.7), including merged multiplicities, the constant projector and conversion to function coordinates. Its optional dense comparison checks the same finite-feature metric and influence actions by a second linear-algebra route. The analytic proof above establishes the asymptotic and operation claims; the floating-point comparisons illustrate the finite formulas. The program prints its output and writes a file only when the caller supplies an explicit output path.

6.1 A finite-schedule resource check

The constants associated with fixed extrapolation order can be large. For $d = s = 1$ and $n = 10^6$, Table 1 evaluates two admissible schedules in (2.6). At the smallest band $t_{K-1} = n^{-\beta}/2^{K-1}$, the actual feature count in (3.1) is $R = 2\lceil\sqrt{32\log n/t_{K-1}}\rceil + 1$. The table records the leading unstructured Gram-formation term nR^2 for that band. The prescribed iteration count is $J = 1058$ for both schedules. These finite counts expose the fixed-order factors in the asymptotic work bound.

η	β	K	t_{K-1}	R	nR^2
0.1	0.02	13	1.851996×10^{-4}	3093	9.566649×10^{12}
0.5	0.10	3	6.279716×10^{-2}	169	2.8561×10^{10}

Table 1: Feature and work-term counts for two admissible fixed schedules at one million samples.

They describe the stated feature-Gram implementation; selection of an application-specific schedule requires quantitative bias and conditioning information as well as this operation count.

Funding

This research received no external funding.

Competing interests

The author declares no competing financial or nonfinancial interests.

Data availability

The analytic results are proved in the article. The editable source and verification code are available in the accompanying source package at the Luminary Research Archive.

Acknowledgements

AI tools assisted mathematical exploration, verification code, literature comparison and manuscript preparation. The author is responsible for the stated results and references.

References

- [1] J. Altschuler, F. Bach, A. Rudi and J. Niles-Weed, Massively scalable Sinkhorn distances via the Nyström method, *Advances in Neural Information Processing Systems* **32** (2019). <https://arxiv.org/abs/1812.05189v3>.
- [2] H. Dong, J. Li, Y. Gong, R. Li and Y. Liu, Sinkhorn linearization and the spectral proxy: unifying the statistical and algorithmic theory of feature-parameterized inverse optimal transport via a single spectral sandwich, preprint (2026). <https://arxiv.org/abs/2608.13201v1>.
- [3] N. Kessler, R. Magnet and J. Feydy, Sinkhorn normalization of diffusion kernels, *Proceedings of Machine Learning Research* **306** (2026), 56764–56795. <https://proceedings.mlr.press/v306/kessler26a.html>.
- [4] P. A. Knight, D. Ruiz and B. Uçar, A symmetry preserving algorithm for matrix scaling, *SIAM J. Matrix Anal. Appl.* **35** (2014), 931–955. doi:10.1137/110825753.
- [5] R. Lakshmanan, A. Pichler and D. Potts, Nonequispaced fast Fourier transform boost for the Sinkhorn algorithm, *Electron. Trans. Numer. Anal.* **58** (2023), 289–

315. doi:10.1553/etna_vol58s289. The inspected preprint is <https://arxiv.org/abs/2201.07524v5>.
- [6] T. L. Saidi, F. Gunsilius and G. Mena, Second-order Sinkhorn geometry and $\sqrt{n}$ -inference, preprint (2026). <https://arxiv.org/abs/2610.36488v1>.
- [7] S. Schubr, Statistical inference for empirical heat-kernel Sinkhorn metrics, Luminary Research Archive, edition 3 (2026). <https://research.luminaryoddities.com/statistical-inference-empirical-heat-kernel-sinkhorn-metrics/v/3>.
- [8] C. L. Wormell and S. Reich, Spectral convergence of diffusion maps: improved error bounds and an alternative normalization, *SIAM J. Numer. Anal.* **59** (2021), 1687–1734. doi:10.1137/20M1344093. The inspected preprint is <https://arxiv.org/abs/2006.02037v2>.

List of Tables

- 1 Feature and work-term counts for two admissible fixed schedules at one million samples. 18