

Moment certificates for homogeneous stable count laws

Scott Schubr
Luminary Oddities LLC

Abstract

We compare a homogeneous real-stable count law P with the multinomial law M having the same mean. The normalized covariance deficit D yields finite chi-square bounds retaining both $x = \text{Tr } D$ and $s_2 = \text{Tr}(D^2)$. In particular, the likelihood ratio has a covariance-determined quadratic component of exact squared norm $ns_2/[2(n-1)]$, with a uniform higher-order remainder. A hybrid Gaussian coefficient envelope gives chi-square convergence when x stays bounded and $s_2 \rightarrow 0$, without a lower bound on retained category means. Matching mean and covariance forces total-variation agreement of order $x^{3/2}$ as $x \rightarrow 0$; a cubic example makes that exponent sharp. Conversely, nonfactorizable positive stable cubic laws rule out a dimension-uniform third-chaos bound by a constant times $s_2^{3/2}$ on the full class.

The main input is the optimal real coefficient inequality $|e_k(z)|^2 \leq \binom{n}{k}$ for centered real vectors of squared norm n , proved in every order by classical same-degree root extrema, a refined contour estimate and complete rational polynomial certificates. We classify equality and retain the optimal universal bound $\text{TV}(P, M) \leq \min\{1, x\}$. We also prove the sharp complex bound $|e_5(z)|^2 \leq \|z\|_2^{10}/1728$ for centered $z \in \mathbb{C}^6$. A reciprocal-phase weighted-median reduction, a classical affine real-root extremum theorem and exact rational polynomial certificates give this finite-dimensional result. Ternary determinantal representation and Gaussian coefficient orthogonality yield the sharp bound $\|U_5\|_2^2 \leq (3/4)h_5(D)$ in degree six at every covariance rank, attained by positive stable product laws. Thus every order in degree six satisfies the complete-homogeneous factor-one target. For general degrees we retain a factor-24 envelope in all orders, strengthen order five to factor 2 and to $10/9$ for $n \geq 256$, and add an improved derivative branch. Stronger structured subclasses and explicit spectral regions are included; unrestricted higher-interior factor one in degrees at least seven remains open. A Gaussian mutual-TV limit along powers of each fixed stable law is also proved. Classical root-extremum and determinantal inputs and the centered coefficient, tensor and orthogonal-expansion methods of Han and Niles-Weed are credited. Standalone standard-library code reconstructs the finite exact certificates.

1 Setting and relation to prior work

A nonzero polynomial is *real stable* if it has real coefficients and no zero when every variable has positive imaginary part. Let

$$p(x) = \sum_{|a|=n} P(a)x^a, \quad P(a) \geq 0, \quad p(\mathbf{1}) = 1$$

be homogeneous and real stable, with integer $n \geq 1$. Here $|a| = \sum_i a_i$ and A has law P . Put $t_i = \mathbb{E}A_i > 0$, $q = t/n$, $L(x) = q \cdot x$, and $M = \text{Mult}(n, q)$. Total variation is $\text{TV}(P, M) = \frac{1}{2} \sum_a |P(a) - M(a)|$. Define

$$B = \frac{1}{2} \left(\text{diag}(t) - \frac{tt^\top}{n} - \text{Cov}(A) \right), \quad W = \text{Tr}(\text{diag}(t)^{-1}B). \quad (1)$$

We will prove $B \succeq 0$. The identity $\sum_i t_i = n$ gives $W = \frac{1}{2}(m - 1 - \sum_i \text{Var}(A_i)/t_i)$.

Multinomial approximation for dependent random variables is classical; Loh's Stein-method treatment [7] is a central predecessor. Ehm's binomial approximation [3] already treats the bi-variate stable subclass, whose generating polynomials factor into linear forms. Roos [9] develops approximation of convolution products, including independent categorical sums. We do not claim that dependent multinomial approximation, or moment-based approximation in these subclasses, is new.

Loh's Theorems 5 and 6 treat, respectively, independent categorical summands and dependent multivariate hypergeometric sampling. Those displayed estimates do not give the uniform covariance-deficit certificate proved here.

The closest methodological predecessor is Han and Niles-Weed [5]. Their analysis of permutation mixtures combines centered elementary symmetric polynomials, Banach transfer and orthogonal chi-square expansions. Their Theorem 4.3(2), equation (27), supplies the real coefficient estimate refined below. Related Banach and coefficient methods occur in Roos [10]; later centered-vector and matrix inequalities appear in [6]. Our argument adapts these tools to the centered hyperbolic expansion of an arbitrary homogeneous stable count law. For categorical observations, a single permutation mixture of independent coordinates has the same count generating polynomial as the underlying product of categorical linear forms. A general exchangeable mixture representation instead yields a mixture of independent reference laws; it does not directly give our one mean-matched multinomial or the deficit in (1).

The main refinement is the covariance-spectrum bound and uniform quadratic likelihood normal form in Theorem 7, together with the sharp matching exponent and the full-stable cubic obstruction in Theorem 21. The complete real scalar coefficient-one certificate, its equality classification, optimal TV coefficient two and fixed-law Gaussian mutual-TV limit supply the surrounding certification theory. Banach's theorem, the predecessor centered ESP method, hyperbolic-slice extremality, and the orthogonal-expansion architecture are credited inputs rather than new general mechanisms. Fixed degree and category count are separate parameters throughout.

Spectral and Frobenius improvements are already present in Han–Niles-Weed [5, Section 5] and [6, Corollary 1.3] for permanent/permutation-product models. Roos [17] already studies signed multivariate Krawtchouk corrections for independent categorical sums, and multinomial orthogonal projections have classical reproducing-kernel descriptions [18]. We claim neither those architectures nor a new general tensor contraction principle. Our bounds cover arbitrary nonfactorizable homogeneous stable count laws, with explicit spectral-square remainder and a sharp obstruction to extending a product-class Frobenius power. The Gaussian tensor-norm and hyperbolicity tools in that obstruction, and quantitative AM–GM at the scalar endpoint, are also classical and explicitly attributed below.

The all-real two-moment estimate of Derksen, Ivanov, Lee and Viola [14, Lemma 36] is another important predecessor. On the centered squared-norm- n slice it gives $|e_k| \leq \binom{n}{k}((k-1)/(n-1))^{k/2}$. Its squared ratio to $\binom{n}{k}$ is $32/25$ at $n = 6, k = 3$, so that displayed estimate alone does not give the value-one bound proved below. Their real-rooted truncation argument and its earlier ancestry are established methods. For binomial-orthonormal biased Krawtchouk polynomials, Hsieh et al. [15, Lemma 3.12] give an $O(k^{1/4})e^{y^2/4}$ bound under fixed-bias and spatial-window hypotheses. Our statement concerns the exact finite bound one at every admissible integer mean and every order. We do not claim the classical ESP–Krawtchouk correspondence, root reduction, recurrence or Hermite picture as new, and make no general all-degree complex-vector or matrix coefficient-one claim. The fixed complex case $(n, k) = (6, 5)$ is proved separately below.

Theorem 1. Let $n \geq 1$ and let $z \in \mathbb{R}^n$ satisfy $\sum_i z_i = 0$. For $1 \leq k \leq n$,

$$|e_k(z)|^2 \leq \binom{n}{k} \left(\frac{\sum_i z_i^2}{n} \right)^k. \quad (2)$$

Also $e_0 = 1$. The universal squared coefficient one is optimal.

Theorem 2. For every integer $n \geq 1$ and every normalized homogeneous real-stable count law with nonnegative coefficients,

$$\chi^2(P\|M) \leq \sum_{k=2}^n (2W)^k, \quad \text{TV}(P, M) \leq \min\{1, 2W\}. \quad (3)$$

Mean-zero categories are deleted before defining W using the retained positive means. The conclusion allows arbitrary category count, repeated roots, zero coefficients and singular covariance deficit. The coefficient two is optimal uniformly over the degree.

The proof is computer-assisted: the exact polynomial identities, infinite-domain reductions and exhaustive finite domains are specified below, with standalone code reconstructing their complete certificates. The scalar theorem is proved by the complete domain partition below, rather than assumed from finite samples.

2 Centered expansion and the exact density identity

Degree one is exactly multinomial. In the lemmas below we therefore assume $n \geq 2$.

Lemma 3. There are homogeneous polynomials R_k of degree k , $2 \leq k \leq n$, invariant under $x \mapsto x + a\mathbf{1}$, such that

$$p(x) = L(x)^n + \sum_{k=2}^n R_k(x) L(x)^{n-k}. \quad (4)$$

Moreover $B \succeq 0$, $B\mathbf{1} = 0$, and $R_2(x) = -x^\top Bx$.

Proof. For real x , set $y = x - L(x)\mathbf{1}$. The polynomial $p(z\mathbf{1} + y)$ is monic of degree n . Every root is real: an upper-half-plane root contradicts stability, and a lower-half-plane root has an upper conjugate. Its coefficient of z^{n-1} is $t \cdot y = 0$. Write its roots as $\lambda_1, \dots, \lambda_n$, so $\sum_j \lambda_j = 0$, and let $R_k(x)$ be its coefficient of z^{n-k} . This defines homogeneous polynomials, invariant under addition of $a\mathbf{1}$ because y is unchanged. Substitution $z = L(x)$ gives (4).

Write $H = \nabla^2 p(\mathbf{1})$. The coefficient of z^{n-2} is $\frac{1}{2}y^\top Hy$. Euler's identity gives $H\mathbf{1} = (n-1)t$ and $\mathbf{1}^\top H\mathbf{1} = n(n-1)$, so

$$R_2(x) = \frac{1}{2}x^\top \left(H - \frac{n-1}{n} tt^\top \right) x.$$

Since $H = \text{Cov}(A) + tt^\top - \text{diag}(t)$, this is $-x^\top Bx$. On the other hand the root identity gives

$$R_2(x) = -\frac{1}{2} \sum_j \lambda_j^2, \quad \sum_j \lambda_j^2 = 2x^\top Bx. \quad (5)$$

Thus $B \succeq 0$; the displayed formula or translation invariance gives $B\mathbf{1} = 0$. □

If $W = 0$, positive t and $B \succeq 0$ imply $B = 0$. Equation (5) then makes every centered root zero for every real x , so all R_k vanish and $P = M$. This proves rigidity without dividing by W . The cases $n = 1$ or $m = 1$ are also exact multinomial laws. Zero-mean categories can be deleted because their nonnegative counts vanish almost surely.

For a homogeneous polynomial R_k , let T_k denote its symmetric k -linear form, with $R_k(x) = T_k(x, \dots, x)$. Our ordered-index convention means that a monomial coefficient at α is $(k!/\alpha!)T_{k,\alpha}$.

Lemma 4. *Let ν_k be the signed coefficient measure of $R_k L^{n-k}$. There exist mutually orthogonal mean-zero densities $U_k = d\nu_k/dM$ satisfying*

$$\frac{P}{M} - 1 = \sum_{k=2}^n U_k, \quad \chi^2(P||M) = \sum_{k=2}^n \mathbb{E}_M U_k^2, \quad (6)$$

and

$$\mathbb{E}_M U_k^2 = \binom{n}{k}^{-1} \sum_{i_1, \dots, i_k} \frac{T_k(e_{i_1}, \dots, e_{i_k})^2}{q_{i_1} \cdots q_{i_k}}. \quad (7)$$

Proof. Translation invariance of R_k gives $T_k(\mathbf{1}, v_2, \dots, v_k) = 0$: first differentiate the diagonal polynomial and then polarize its remaining arguments. Let $X_1, \dots, X_n$ be independent categorical variables of law q and set

$$H_k(i_1, \dots, i_k) = \frac{T_k(e_{i_1}, \dots, e_{i_k})}{q_{i_1} \cdots q_{i_k}}, \quad U_k = \binom{n}{k}^{-1} \sum_{|S|=k} H_k(X_S).$$

Symmetry makes U_k a function of the count vector. Summing any one argument of H_k against q gives zero by the contraction identity. Hence the kernel is completely conditionally degenerate. For each fixed subset S ,

$$\mathbb{E}[H_k(X_S) x^{\text{counts}(X)}] = R_k(x) L(x)^{n-k}.$$

Coefficient comparison after symmetrization proves $\nu_k(a) = M(a)U_k(a)$. All $M(a)$ on $|a| = n$ are positive. In a product of two kernels associated with different subsets, an index in the symmetric difference occurs in only one kernel; integrate that independent categorical observation to obtain zero. This applies both within an order and between distinct orders. Thus only identical subsets contribute to the square norm, giving (7), while (4) gives (6). All operations are finite sums. $\square$

3 Hilbert tensor transfer and a universal certificate

We use Banach's classical real Hilbert-space theorem [1, 2]: the norm of a symmetric multilinear form on arbitrary unit-vector tuples equals its norm on diagonal tuples. It is also the input in [5, Lemma 4.2].

Lemma 5. *If $|R_k(x)| \leq c(x^\top Bx)^{k/2}$ for all real x , then*

$$\mathbb{E}_M U_k^2 \leq \frac{c^2 (nW)^k}{\binom{n}{k}}. \quad (8)$$

Proof. For $v \in \ker B$, the polynomial $R_k(x + sv)$ is bounded independently of real s , hence is constant. Differentiation and polarization give $T_k(v, v_2, \dots, v_k) = 0$. Thus T_k descends to the real Hilbert quotient by $\ker B$, with squared norm $x^\top Bx$. Banach's theorem there implies

$$|T_k(v_1, \dots, v_k)| \leq c \prod_j \sqrt{v_j^\top B v_j}.$$

Apply this to coordinate vectors in (7) and sum the products to obtain

$$\mathbb{E}_M U_k^2 \leq \binom{n}{k}^{-1} c^2 \left(\sum_i \frac{B_{ii}}{q_i} \right)^k = \binom{n}{k}^{-1} c^2 (nW)^k.$$

No inverse of B is used. □

Han and Niles-Weed [5, Theorem 4.3(2)] prove the uniform real centered coefficient bound $|e_k(z)|^2 \leq 10 \binom{n}{k}$ when $\sum_j z_j^2 = n$. Below we refine their binary contour and Beta-integral argument to a constant below three. The classical extremum reduction needed for that refinement is stated first; the bound with constant ten serves as a comparison, rather than as a tail input in our final theorem.

The next elementary bound retains positivity of the actual density. It is the upper-distance branch of the classical sharp relation between chi-square divergence and total variation; see Guntuboyina, Saha and Schiebinger [4, Corollary 5.3], which credits Gilardoni's Proposition 2.1. We give the short proof needed here and make no claim of a new general divergence inequality.

Lemma 6. *For probability measures $P \ll M$, write $v = \chi^2(P \| M)$ and $\delta = \text{TV}(P, M)$. If $\delta > 1/2$, then $v \geq \delta/(1 - \delta)$.*

Proof. Let $Y = dP/dM - 1$ and $a = M(Y < 0)$. Then $\mathbb{E}Y_+ = \mathbb{E}Y_- = \delta$ and $Y_- \leq 1$, so $a \geq \delta$. Cauchy–Schwarz on the two parts of the partition gives

$$v \geq \frac{\delta^2}{a} + \frac{\delta^2}{1-a} = \frac{\delta^2}{a(1-a)}.$$

For $a \geq \delta > 1/2$, $a(1-a) \leq \delta(1-\delta)$. The degenerate partition cases force $\delta = 0$ and do not occur here. □

We first establish the contour and finite-coefficient tools. The central and boundary estimates then prove the scalar theorem in every order, after which the orthogonal transfer and density positivity give the count-law bounds.

4 Covariance-spectrum bounds and a quadratic normal form

Put $T = \text{diag}(t)$ and

$$D = 2T^{-1/2}BT^{-1/2}, \quad x = \text{Tr } D = 2W, \quad s_2 = \text{Tr}(D^2), \quad a_n = \frac{n}{2(n-1)} \quad (n \geq 2).$$

The matrix D annihilates $\sqrt{t}$ and satisfies $0 \preceq D \preceq I - \sqrt{t}\sqrt{t}^\top/n$, by covariance positivity. For $r \geq 0$ define the nonnegative finite spectral polynomials

$$g_r(D) = [z^r] \det(I - 2zD)^{-1/2}, \quad b_r(D) = \min\{x^r, g_r(D)\}.$$

These coefficients are defined for every PSD D , regardless of convergence at $z = 1$.

Theorem 7. *For $n \geq 2$, the exact quadratic density in Lemma 4 satisfies*

$$\|U_2\|_2^2 = a_n s_2. \tag{9}$$

For $3 \leq k \leq n$,

$$\|U_k\|_2^2 \leq \min\{s_2 x^{k-2}, s_2 g_{k-2}(D), g_k(D)\}. \quad (10)$$

Consequently, with $J_n(D) = \sum_{r=1}^{n-2} b_r(D)$,

$$\chi^2(P\|M) = a_n s_2 + \|P/M - 1 - U_2\|_2^2, \quad (11)$$

$$0 \leq \|P/M - 1 - U_2\|_2^2 \leq s_2 J_n(D), \quad (12)$$

$$a_n s_2 \leq \chi^2(P\|M) \leq s_2(a_n + J_n(D)). \quad (13)$$

All norms are in $L^2(M)$ and all sums are finite; no smallness assumption on x is needed. If $x = 0$, all terms vanish by rigidity. Empty sums apply at $n = 2$.

The gain over trace-only control retains s_2/x^2 , the reciprocal effective rank when $x > 0$. Spectral permanent bounds already occur in Han–Niles–Weed [5, Section 5]; their later matrix theorem and Corollary 1.3 also retain Frobenius information [6]. The result here concerns the full homogeneous stable class and the explicit covariance-controlled remainder, rather than a new spectral or orthogonal-expansion principle.

Lemma 8. *If S is a symmetric real k -linear form of multilinear norm at most one and $\Lambda \succeq 0$ has eigenvalues λ_i , then, for $k \geq 2$,*

$$\sum_{i_1, \dots, i_k} \lambda_{i_1} \cdots \lambda_{i_k} S(e_{i_1}, \dots, e_{i_k})^2 \leq \text{Tr}(\Lambda^2) (\text{Tr} \Lambda)^{k-2}.$$

At $k = 2$ the right side means $\text{Tr}(\Lambda^2)$.

Proof. Fix the last $k - 2$ unit basis arguments. The symmetric matrix $C_{ij} = S(e_i, e_j, \dots)$ is an operator contraction. Thus

$$\sum_{i,j} \lambda_i \lambda_j C_{ij}^2 \leq \frac{1}{2} \sum_{i,j} (\lambda_i^2 + \lambda_j^2) C_{ij}^2 = \sum_i \lambda_i^2 \sum_j C_{ij}^2 \leq \sum_i \lambda_i^2.$$

Symmetry justifies the equality and the row norms are at most one. Sum the remaining nonnegative weights. This is an elementary weighted Hilbert–Schmidt contraction argument. $\square$

Proof of Theorem 7. The scalar theorem gives $|R_k(v)|^2 \leq \binom{n}{k} (2/n)^k (v^\top B v)^k$. Use the quotient and Banach argument of Lemma 5. The weighted coordinate family has covariance

$$\Lambda = B^{1/2} \text{diag}(q)^{-1} B^{1/2}.$$

Its nonzero spectrum equals that of $(n/2)D$ by the $XX^\top, X^\top X$ identity. Lemma 8 and the exact tensor identity cancel the factors $\binom{n}{k}$ and $(2/n)^k (n/2)^k$, giving $\|U_k\|_2^2 \leq s_2 x^{k-2}$. For $k = 2$, use $T_2 = -B$ directly:

$$\|U_2\|_2^2 = \frac{n^2}{\binom{n}{2}} \text{Tr}((T^{-1/2} B T^{-1/2})^2) = a_n s_2.$$

No inverse of B is used.

For clarity we give the Gaussian steps, with all normalizations. The whitened polynomial $F_k(y) = R_k(\text{diag}(q)^{-1/2} y)$ has ordered symmetric tensor A_k and

$$|F_k(y)|^2 \leq \binom{n}{k} (y^\top D y)^k, \quad \|U_k\|_2^2 = \|A_k\|_{\text{F}}^2 / \binom{n}{k}.$$

For a standard real Gaussian G , the degree- k Hermite component of $F_k(G)$ has squared norm $k!\|A_k\|_{\mathbb{F}}^2$. Indeed the coefficient of the product probabilists' Hermite H_α is $(k!/\alpha!)A_{k,\alpha}$, and $\mathbb{E}H_\alpha^2 = \alpha!$. Lower-degree chaos squares are nonnegative. Hence

$$\|U_k\|_2^2 \leq \mathbb{E}(G^\top DG)^k/k! = g_k(D).$$

The last identity follows by integrating independent Gaussian squares after diagonalization, initially near $z = 0$, and comparing finite coefficients.

For the stronger hybrid bound, kernel descent expresses $A_k = \sqrt{\binom{n}{k}} S \circ (D^{1/2})^{\otimes k}$ with S symmetric of multilinear norm at most one on $\text{ran } D$. Work in an eigenbasis and put $r = k - 2$ and

$$P_{ij}(G) = S(e_i, e_j, D^{1/2}G, \dots, D^{1/2}G).$$

Its highest Hermite chaos has squared norm $r!$ times the remaining r -index weighted tensor square. Multiply by $\lambda_i \lambda_j$, sum, and discard the nonnegative lower-chaos norms. Pointwise Lemma 8's matrix argument bounds the inner bilinear sum by $s_2(G^\top DG)^r$. Thus

$$r!\|U_k\|_2^2 \leq s_2 \mathbb{E}(G^\top DG)^r, \quad \|U_k\|_2^2 \leq s_2 g_r(D).$$

For $r = 0$ this is just the matrix argument, while the exact quadratic identity is sharper. Orthogonality proves the remaining assertions. $\square$

Corollary 9. *If $\rho = \|D\|_{\text{op}} < 1/2$, then*

$$\chi^2(P\|M) \leq s_2\{a_n + \det(I - 2D)^{-1/2} - 1\} \leq s_2\{a_n + \exp(x + s_2/(1 - 2\rho)) - 1\}.$$

For any sequence in the full stable class with bounded x and $s_2 \rightarrow 0$, one has $\chi^2(P\|M) \rightarrow 0$ and $\text{TV}(P, M) \rightarrow 0$, even when degree, category count and positive means vary.

Proof. The convergent generating series sums the hybrid coefficients. Also

$$\log \det(I - 2D)^{-1/2} = x + \sum_{j \geq 2} \frac{2^{j-1}}{j} \text{Tr}(D^j) \leq x + \frac{s_2}{1 - 2\rho},$$

since $\text{Tr}(D^j) \leq \rho^{j-2}s_2$ and $2^{j-1}/j \leq 2^{j-2}$. Finally $\rho \leq \sqrt{s_2}$, so the required spectral-radius condition holds eventually. $\square$

For the product/permutation-mixture subclass, convergence from small Frobenius norm already follows from [6, Corollary 1.3]. Our assertion extends the domain, not the concept of this convergence criterion. The determinant above is not the Gaussian-likelihood expression $\det(I - D^2)^{-1/2} - 1$: for $p = (XY + XZ + YZ)/3$ the latter is $1/3$, whereas the actual chi-square is $1/2$ and D has spectrum $(1/2, 1/2, 0)$.

4.1 Quadratic approximation and covariance matching

Put $H_n(x) = \sum_{r=1}^{n-2} x^r$. Since $J_n(D) \leq H_n(x)$,

$$0 \leq \frac{\chi^2(P\|M)}{a_n s_2} - 1 \leq \frac{H_n(x)}{a_n} \leq \frac{2x}{1-x} \quad (s_2 > 0, 0 \leq x < 1).$$

This is a small-deficit likelihood normal form, uniform in degree, categories and positive means, rather than a central-limit or sampling theorem. The exact quadratic is

$$U_2(a) = -\frac{n}{n-1} \left[(a-t)^\top T^{-1} B T^{-1} (a-t) - \sum_i \frac{B_{ii} a_i}{t_i^2} \right]. \quad (14)$$

To verify it, the ordered distinct observation pairs in the degenerate kernel have multiplicities $a_i a_j - \delta_{ij} a_i$. Their contraction is divided by $n(n-1)$; substitute $q = t/n$ and use $B\mathbf{1} = 0$ to center a . The signed measure $M(1 + U_2)$ need not be nonnegative. Signed multivariate Krawtchouk approximations for independent categorical sums are classical [17], as are multinomial orthogonal projections [18]; the new estimate controls this correction's full-stable remainder.

With $\delta_2 = \frac{1}{2} \mathbb{E}_M |U_2|$, reverse triangle and Cauchy give

$$|\text{TV}(P, M) - \delta_2| \leq \frac{1}{2} \sqrt{s_2 J_n(D)}, \quad \text{TV}(P, M) \leq \frac{1}{2} \sqrt{s_2 (a_n + J_n(D))}.$$

If P, Q have the same n, t and covariance, they share U_2, D and M , whence

$$\text{TV}(P, Q) \leq \min\{1, \sqrt{s_2 J_n(D)}\} \leq \frac{x^{3/2}}{\sqrt{1-x}} \quad (0 \leq x < 1). \quad (15)$$

The first bound follows by applying Cauchy to each higher-chaos remainder; the last uses $J_n \leq x/(1-x)$ and $s_2 \leq x^2$.

The exponent 3/2 in this uniform matching conclusion is sharp. For $0 < \epsilon < 1/4$, take three independent Bernoulli probabilities $(1/2 + \epsilon, 1/2 + \epsilon, 1/2 - 2\epsilon)$, and let P_- reverse the sign of ϵ in P_+ . With $L = (X + Y)/2$,

$$p_\pm = L^3 - 3\epsilon^2 (X - Y)^2 L \mp 2\epsilon^3 (X - Y)^3.$$

They have $t = (3/2, 3/2)$, $B = 3\epsilon^2 \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$, $x = 8\epsilon^2$, and $s_2 = x^2$. Direct use of the exact tensor norm gives

$$\|U_2\|_2^2 = 48\epsilon^4, \quad \|U_3\|_2^2 = 256\epsilon^6 = x^3/2, \quad \frac{\chi^2(P_+ \| M)}{a_3 s_2} - 1 = \frac{2}{3}x.$$

The absolute coefficient sum of $p_+ - p_- = -4\epsilon^3 (X - Y)^3$ is $32\epsilon^3$, so $\text{TV}(P_+, P_-) = 16\epsilon^3 = x^{3/2}/\sqrt{2}$. Thus neither the relative $O(x)$ remainder nor the matching $O(x^{3/2})$ exponent improves uniformly; the upper constants are not claimed optimal.

4.2 Rare cells and a nonproduct illustration

Small deficit does not uniformly imply the local Taylor ratios $\text{KL}/\chi^2 \rightarrow 1/2$ and $H^2/\chi^2 \rightarrow 1/4$, where $H^2 = \sum (\sqrt{P} - \sqrt{M})^2$. For $n = 2$ and $0 < q < 1/2$, independent Bernoulli probabilities $0, 2q$ give

$$P = (1 - 2q, 2q, 0), \quad M = ((1 - q)^2, 2q(1 - q), q^2), \quad x = q/(1 - q), \quad \chi^2 = x^2.$$

Here

$$\text{KL}(P \| M) = (1 - 2q) \log \frac{1 - 2q}{(1 - q)^2} + 2q \log \frac{1}{1 - q} = q^2 + O(q^3),$$

and $H^2 = q^2 + O(q^3)$ as well. The latter follows by expanding the affinity $(1 - q)\sqrt{1 - 2q} + 2q\sqrt{1 - q}$; the absent count-two mass provides its leading deficit. Both ratios therefore tend to one. By contrast, probabilities $1/2 \pm \epsilon$ have a fixed positive reference law and yield the ordinary ratios

$1/2, 1/4$ by Taylor expansion. This is a barrier to a uniform local expansion, not to classical divergence inequalities.

For a diffuse nonproduct example, let $p = e_d(X_1, \dots, X_m)/\binom{m}{d}$, $2 \leq d < m$. This classical polynomial is real stable, with $t_i = d/m$ and D eigenvalue $(d-1)/(m-1)$ repeated $m-1$ times. Thus $x = d-1$ and $s_2 = (d-1)^2/(m-1)$. For fixed $d \geq 3$, $x \geq 2$ yet Corollary 9 applies as $m \rightarrow \infty$. Directly, with $c = (m)_d/m^d$, the multinomial conditional on no collisions is P , giving the classical birthday identities

$$\text{TV}(P, M) = 1 - c, \quad \chi^2(P||M) = c^{-1} - 1 \sim d(d-1)/(2m) \sim a_d s_2.$$

No categorical-product representation exists: multiaffinity forces distinct linear factors to have disjoint variable supports, which cannot generate every d -subset when $m > d$. These formulas illustrate the domain rather than constitute new birthday results.

4.3 The complete-homogeneous target in low orders

Let $h_k(D) = [z^k] \det(I - zD)^{-1}$. The stronger stable-specific target $\|U_k\|_2^2 \leq h_k(D)$ remains open in general. The following partial conclusion makes its present domain precise.

Proposition 10. *The target holds at $k = 2, 3$ for every admissible degree, and at $k = 4$ for every $n \geq 5$. At $k = 3$, or $k = 4, n \geq 5$, it is strict when $D \neq 0$. Rank-one D is covered at every order. The top-order theorem below supplies the remaining quartic degree $n = 4$.*

Proof. The exact quadratic energy is at most $s_2 \leq h_2 = (x^2 + s_2)/2$. Newton gives $h_3 - xs_2 = e_3(\lambda(D)) \geq 0$, so Theorem 7 already proves the non-strict cubic assertion. For strictness, the centered scalar cubic maximum from the binary calculation in Lemma 27 is $\beta_{n,3} = 2n(n-2)/[3(n-1)^2] < 2/3$. A squared scalar constant β multiplies both spectral-square and hybrid bounds by β , since the normalized form's norm is at most $\sqrt{\beta}$. Thus $\|U_3\|_2^2 \leq \beta_{n,3}s_2x < h_3$ for nonzero D .

For every nonzero PSD spectrum,

$$h_4(D) > \frac{3}{5}s_2g_2(D), \quad g_2 = x^2/2 + s_2. \quad (16)$$

To prove it normalize $x = 1$, and put $p = \text{Tr } D^2$, $q = \text{Tr } D^3$, $v = \text{Tr } D^4$. Here $0 < p \leq 1$, $q \geq p^2$ and $v \geq p^3$ by Cauchy/Jensen with weights given by the eigenvalues. Newton gives $24h_4 = 1 + 6p + 3p^2 + 8q + 6v$. Therefore five times the lower bound for $24[h_4 - (3/5)p(1/2 + p)]$ is

$$5 - 6p - 17p^2 + 30p^3 = \frac{3}{2} - \frac{1}{2}w + (28 + 30w)w^2 \geq 13(w - 1/52)^2 + \frac{3}{2} - \frac{1}{208} > 0, \quad w = p - \frac{1}{2} \in [-1/2, 1/2].$$

Homogeneity restores (16).

For $n \geq 6$ the centered scalar quartic bound of Lemma 27 is at most $3/5$: its comparison reduces to $9n^2 - 66n + 72 \geq 0$, zero at six and increasing thereafter. At $n = 5$ the two reflected multiplicities $r = 1, 2$ give squared scalar ratios $45/256$ and $5/9$, respectively, both below $3/5$; these follow directly from the raw generating products. Scaling the hybrid bound by these scalar constants and applying (16) proves the quartic assertion. If rank $D = 1$, then $h_k(D) = x^k$ and the trace theorem suffices at every order. $\square$

4.4 Top-order complete-homogeneous bound by ternary restriction

Theorem 11. *For every degree $n \geq 2$ and every admissible covariance deficit, including singular D ,*

$$\|U_n\|_2^2 \leq h_n(D).$$

Consequently the complete-homogeneous target holds at $k = 4$ for every admissible degree. More strongly, for every $w \in \mathbb{C}^m$,

$$|R_n(w)|^2 \leq \left(\frac{2}{n} w^* B w \right)^n. \quad (17)$$

Proof. We use the classical Helton–Vinnikov theorem: every degree- n ternary hyperbolic form has a definite real symmetric degree-sized linear determinantal representation. See Hanselka [22, Theorem 1], including repeated-root and nonsmooth forms. This deep ternary theorem is an input; no global determinantal representation of p is assumed.

Fix $w = a + ib$. Put $a_0 = a - L(a)\mathbf{1}$, $b_0 = b - L(b)\mathbf{1}$ and $F(t, s, u) = p(t\mathbf{1} + sa_0 + ub_0)$. Real stability and conjugation imply hyperbolicity in $(1, 0, 0)$; its leading t coefficient is one even if these directions are dependent. The centered expansion is

$$F(t, s, u) = t^n + \sum_{j=2}^n R_j(sa_0 + ub_0)t^{n-j}.$$

The classical theorem and congruence normalization give $F = \det(tI_n + sA + uC)$ with real symmetric A, C . Indeed the definite coefficient at $(1, 0, 0)$ may be chosen positive, and its determinant is $F(1, 0, 0) = 1$, so the congruence introduces no scalar factor. Vanishing of the first centered coefficient gives $\text{Tr } A = \text{Tr } C = 0$. Comparison of the second coefficient yields

$$-\frac{1}{2} \text{Tr}(sA + uC)^2 = -(sa_0 + ub_0)^\top B(sa_0 + ub_0).$$

In particular $\text{Tr } A^2 = 2a^\top B a$ and $\text{Tr } C^2 = 2b^\top B b$, using $B\mathbf{1} = 0$. Translation invariance and the determinant identity at $(t, s, u) = (0, 1, i)$ imply $R_n(w) = \det(A + iC)$. For its singular values σ_j , AM–GM gives

$$|R_n(w)|^2 = \prod_{j=1}^n \sigma_j^2 \leq \left(\frac{\sum_j \sigma_j^2}{n} \right)^n, \quad \sum_j \sigma_j^2 = \text{Tr}[(A - iC)(A + iC)] = 2w^* B w.$$

The commutator has zero trace. This proves (17), including zero right side, without inverting B .

Put $Q = \text{diag}(q)$. Let Z be standard circular complex Gaussian with independent coordinates of density $\pi^{-1}e^{-|z|^2}$. For the ordered symmetric tensor A_n of $R_n(Q^{-1/2}z)$, complex monomial orthogonality gives

$$\mathbb{E}|R_n(Q^{-1/2}Z)|^2 = n! \|A_n\|_F^2, \quad \|U_n\|_2^2 = \|A_n\|_F^2.$$

The second identity uses the exact density denominator $\binom{n}{n} = 1$. Since $D = (2/n)Q^{-1/2}BQ^{-1/2}$, integration of (17) yields

$$\|U_n\|_2^2 \leq \frac{\mathbb{E}(Z^* D Z)^n}{n!} = h_n(D).$$

For the last equality, unitary diagonalization gives independent unit-mean exponential squared moduli, with generating function $\prod_i (1 - z\lambda_i)^{-1}$. This is a finite coefficient identity at arbitrary PSD D , not a Gaussian approximation or a spectral-radius restriction. The quartic conclusion now follows from Proposition 10 and the case $n = 4$ just proved. At $n = 1$ the centered density is zero. $\square$

For clarity, applying the same argument to the normalized degree- k derivative $(k!/n!)D_1^{n-k}p$ only gives

$$\|U_k\|_2^2 \leq \binom{n}{k} \left(\frac{k-1}{n-1} \right)^k h_k(D).$$

Its top coefficient is $R_k/\binom{n}{k}$ and its normalized deficit is $(k-1)D/(n-1)$. The displayed factor can exceed one. Thus this derivative argument alone leaves higher interior orders open; the degree-six case is closed below, while general degrees $n \geq 7$ remain unresolved outside the stated covered regions. This proof neither complexifies the real scalar theorem nor supplies a complex coefficient-one inequality.

Theorem 12 (All-order comparison from the known complex factor). *For every $2 \leq k \leq n$,*

$$\|U_k\|_2^2 \leq \min\{24, \beta_{n,k}^{\mathbb{C}}\} h_k(D), \quad \beta_{n,k}^{\mathbb{C}} = \binom{n}{k} \left(\frac{k-1}{n-1} \right)^k. \quad (18)$$

The factor 24 is the known complex-vector constant of Han–Niles–Weed, not a new scalar estimate. All coefficients are finite at arbitrary PSD D . In particular, if $\rho(D) < 1$,

$$\begin{aligned} \chi^2(P\|M) &\leq 24\{\det(I-D)^{-1} - 1 - x\}, \\ \chi^2(P\|M) &\leq a_n s_2 + 24\{\det(I-D)^{-1} - 1 - x - h_2(D)\}. \end{aligned} \quad (19)$$

Proof. For each complex direction w , the ternary pencil in the preceding proof gives $M_w = A + iC$ with $\text{Tr } M_w = 0$ and $\|M_w\|_F^2 = 2w^* B w$. Its characteristic coefficients give $R_k(w) = e_k(\lambda(M_w))$, even if the matrix is not diagonalizable. Schur triangularization gives $\sum_j |\lambda_j(M_w)|^2 \leq \|M_w\|_F^2$. Apply the scaled complex centered-vector theorem of Han–Niles–Weed [6, Theorem 1.1]:

$$|R_k(w)|^2 \leq 24 \binom{n}{k} \left(\frac{2}{n} w^* B w \right)^k.$$

If all eigenvalues vanish the assertion is immediate. For standard circular complex Gaussian Z , substitute $w = Q^{-1/2}Z$. Complex monomial orthogonality and the exact density normalization give

$$k! \|A_k\|_F^2 \leq 24 \binom{n}{k} \mathbb{E}(Z^* D Z)^k = 24 \binom{n}{k} k! h_k(D), \quad \|U_k\|_2^2 = \frac{\|A_k\|_F^2}{\binom{n}{k}}.$$

Cancellation proves the factor-24 branch, and the derivative argument above proves the other branch. Thus finite orthogonality gives $\chi^2(P\|M) \leq 24 \sum_{k=2}^n h_k(D)$. If $\rho(D) < 1$, extending the nonnegative generating series to all orders proves the first determinant bound. Keeping the exact quadratic energy $a_n s_2$ and extending only orders $k \geq 3$ proves the second. The determinant has power minus one, distinct from the real-Gaussian envelope’s power minus one half. $\square$

One may combine all these branches order by order:

$$\chi^2(P\|M) \leq a_n s_2 + \sum_{k=3}^n \min\{s_2 x^{k-2}, s_2 g_{k-2}(D), g_k(D), 24 h_k(D), \beta_{n,k}^{\mathbb{C}} h_k(D)\}.$$

The general higher-interior factor-one target in degrees $n \geq 7$ remains open; the known complex factor 24 is not an exact-one substitute. The representation is still chosen separately for each ternary restriction.

4.5 A sharper full-spectrum bound at order five

Order five admits a stronger uniform comparison in general degree. The proof uses the same real scalar-to-hybrid transfer, with an explicit fifth-order scalar constant, and elementary symmetric-polynomial inequalities. It does not require a complex coefficient-one theorem.

Lemma 13. *On the centered real squared-norm- n slice, let $\beta_{n,5} = \max |e_5|^2 / \binom{n}{5}$. For every $n \geq 6$,*

$$\beta_{n,5} \leq \frac{3}{4}.$$

The uniform constant is sharp, attained at $n = 6$ and binary multiplicity $r = 2$. For $n \geq 8$ one also has

$$\beta_{n,5} \leq B_n := \frac{4n(5n-12)^3}{1215(n-1)(n-2)(n-3)(n-4)}. \quad (20)$$

Proof. By the classical two-root reduction in Lemma 22, take multiplicities $r, s = n - r$. Put $c = (s - r)/\sqrt{rs}$ and $u = c^2 \in [0, U]$, where $U = (n - 2)^2/(n - 1)$. Newton identities give

$$e_5 = \frac{nc(6c^2 + 12 - 5n)}{30}, \quad \beta = \frac{n^2 u(6u + 12 - 5n)^2}{900 \binom{n}{5}}.$$

For $A = 5n - 12$, the derivative of $u(6u - A)^2$ is $(6u - A)(18u - A)$. Its maximum on $[0, U]$ is therefore bounded by the larger of its value at U and its local-maximum value at $A/18$, even when the latter lies outside the interval. Their normalized contributions are

$$E_n = \frac{2n(n-2)(n-3)(n-4)}{15(n-1)^4} < \frac{2}{15}, \quad B_n.$$

For $n \geq 8$, $5n - 12 \geq (7/2)n$ and the denominator product in (20) is less than n^4 , so $B_n > 343/2430 > 2/15$. This proves (20). Moreover $B_n < 3/4$ for $n \geq 8$: after clearing positive denominators, the difference is

$$\begin{aligned} & 3645(n-1)(n-2)(n-3)(n-4) - 16n(5n-12)^3 \\ &= 251944 + 468998y + 195495y^2 + 30590y^3 + 1645y^4 > 0, \quad y = n - 8 \geq 0. \end{aligned}$$

At $n = 6$, reflected multiplicities $r = 1, 2, 3$ give $96/3125, 3/4, 0$;

at $n = 7$ they give $7/162, 6776/9375, 7/64$. These direct rational evaluations finish the proof. $\square$

Lemma 14 (Sharp fifth spectral comparison). *For every PSD D ,*

$$h_5(D) \geq \frac{3}{8}s_2g_3(D). \quad (21)$$

For nonzero D , equality holds exactly when its positive spectrum consists of two equal eigenvalues. Thus $3/8$ is the best universal constant.

Proof. The zero case is immediate. Normalize $x = 1$ and write $p = p_2, q = p_3, r = p_4, s = p_5$, where $p_j = \text{Tr}(D^j)$. Newton identities give

$$g_3 = \frac{1 + 6p + 8q}{6}, \quad 120h_5 = 1 + 10p + 15p^2 + 20q + 20pq + 30r + 24s.$$

Hence $E = 120[h_5 - (3/8)pg_3]$ equals

$$E = 1 + \frac{5}{2}p - 30p^2 + (20 - 40p)q + 30r + 24s. \quad (22)$$

If $p \leq 1/2$, Cauchy/Jensen with the eigenvalues as probability weights yields $q \geq p^2, r \geq p^3, s \geq p^4$, and the q coefficient is nonnegative. Thus

$$E \geq (1 - 2p)(1 + \frac{9}{2}p - p^2 - 12p^3) \geq 0.$$

The second factor is concave on $[0, 1/2]$ and has endpoint values $1, 3/2$, so this is strict if $p < 1/2$.

For $p \geq 1/2$, let e_j be the elementary symmetric functions of the normalized eigenvalues. Another Newton rearrangement of (22) gives

$$E = \frac{15}{2}(2p-1)(4-3p) + (240-60p)e_3 - 240e_4 + 120e_5.$$

Since $e_3 = xe_3 \geq 4e_4$ and $p \leq 1$,

$$E \geq \frac{15}{2}(2p-1)(4-3p) + (180-60p)e_3 + 120e_5 \geq 0.$$

Equality forces $p = 1/2$ and $e_3 = 0$, hence exactly two equal positive eigenvalues. Conversely that spectrum gives equality. For eigenvalues (a, a) , $h_5 = 6a^5$, $s_2 = 2a^2$ and $g_3 = 8a^3$, proving sharpness. Homogeneity restores arbitrary x . $\square$

Theorem 15 (Fifth-order full-spectrum refinement). *For every admissible stable count law of degree $n \geq 6$,*

$$\|U_5\|_2^2 \leq \min\{2, \beta_{n,5}^C\} h_5(D). \quad (23)$$

For $n \geq 8$ the sharper coefficient is

$$\|U_5\|_2^2 \leq C_n h_5(D), \quad C_n = \min\{2, \beta_{n,5}^C, (8/3)B_n\}. \quad (24)$$

In particular, $\|U_5\|_2^2 \leq (10/9)h_5(D)$ for every $n \geq 256$ and every spectrum. These bounds include singular D and impose no spectral-radius restriction. The third branch tends to $800/729 > 1$; no unrestricted factor-one conclusion follows. At $n = 5$ the top-order factor one is stronger.

Proof. A scalar squared constant β multiplies the hybrid bound by β , as in Proposition 10. Lemmas 13 and 14 give respectively the factors 2 and $(8/3)B_n$. The derivative branch is already established in (18). At $n = 6$ it is $6144/3125 < 2$, so it is retained.

To obtain the stated high-degree corollary, $B_n \leq 5/12$ is equivalent to

$$P(n) = 6075(n-1)(n-2)(n-3)(n-4) - 48n(5n-12)^3 \geq 0.$$

Its exact shift at $n = 256 + y$ is

$$P(256 + y) = 34765845384 + 1638253434y + 16121745y^2 + 59250y^3 + 75y^4.$$

Every coefficient is positive, proving the assertion on the entire real ray $n \geq 256$ and hence every integer degree there. The limit of $(8/3)B_n$ is $(8/3)(500/1215) = 800/729$. The threshold is a convenient sufficient one, not an optimality claim. Sharpness of the separate scalar and spectral inputs does not assert simultaneous attainment by a stable law, so factor 2 is not claimed optimal for the count-law problem. $\square$

Corollary 16 (Two fifth-order factor-one regions). *Let $n \geq 6$ and $D \neq 0$. The factor-one bound holds if either*

$$\frac{s_2}{x^2} \leq \frac{1}{11} \quad \text{or} \quad \frac{\lambda_{\max}(D)}{x} \geq \frac{7}{8}.$$

The corresponding gaps $h_5 - \|U_5\|_2^2$ are at least $17x^5/53240$ and $x^5/4096$, respectively. For general degrees $n \geq 7$, an intermediate spectral region remains uncovered.

Proof. Normalize $x = 1$. For $p \leq 1/11$, use $\|U_5\|_2^2 \leq (3/4)pg_3$ and

$$\begin{aligned} 120[h_5 - (3/4)pg_3] &= 1 - 5p - 75p^2 + (20 - 100p)q + 30r + 24s \\ &\geq 1 - 5p - 55p^2 - 70p^3 \geq 51/1331. \end{aligned}$$

Here the q coefficient is positive, $q \geq p^2, r \geq p^3$, and the last polynomial decreases for $p \geq 0$. For $\alpha = \lambda_{\max}(D) \geq 7/8$, the spectral-square branch and the monomials of h_5 give

$$\|U_5\|_2^2 \leq \frac{3}{4}p \leq \frac{3}{4}[\alpha^2 + (1 - \alpha)^2], \quad h_5 \geq \alpha^5 + \alpha^4(1 - \alpha) = \alpha^4.$$

Their difference is $f(\alpha) = \alpha^4 - (3/2)\alpha^2 + (3/2)\alpha - 3/4$, with $f(7/8) = 1/4096$ and $f'(\alpha) > 0$ on $[7/8, 1]$. Rescale by x^5 . $\square$

In any orderwise chi-square bound, the $k = 5$ summand may therefore be replaced by these stronger branches. The ingredients are the credited real root reduction and tensor transfer, together with the explicit fifth-degree calculation and sharp spectral comparison; no new general symmetric-function or Gaussian method is claimed. In particular, the unrestricted target at $5 \leq k < n$ remains a separate problem.

4.6 A sharp complex fifth coefficient in dimension six

Theorem 17. *For every $z \in \mathbb{C}^6$ with $\sum_i z_i = 0$,*

$$|e_5(z)|^2 \leq \frac{(\sum_i |z_i|^2)^5}{1728}. \quad (25)$$

The constant is optimal. Consequently every degree-six homogeneous stable count law, in every covariance-deficit rank, satisfies the sharp bound

$$\|U_5\|_2^2 \leq \frac{3}{4}h_5(D). \quad (26)$$

In particular, together with the lower-order and top-order results, every order in degree six satisfies the complete-homogeneous factor-one target.

Proof. The zero vector is immediate. Otherwise homogeneity gives $\sum |z_i|^2 = 1$. Two zero coordinates give $e_5 = 0$; exactly one gives $|e_5|^2 \leq 5^{-5} < 1/1728$ by AM–GM. Otherwise write $z_i = r_i u_i$, where $r_i > 0$, $|u_i| = 1$ and $\sum r_i u_i = 0$. With $P_6 = \prod r_i$, for every real c ,

$$|e_5(z)| = P_6 \left| \sum_i \frac{\overline{u_i}}{r_i} \right| \leq P_6 \sum_i |r_i^{-1} - cr_i|.$$

The right side is minimized at a weighted median $c = t^{-2}$ of the numbers r_i^{-2} with weights r_i , where $t = r_j$ is one radius. Choose $\sigma_i = 1$ for $r_i < t$ and $\sigma_i = -1$ for $r_i > t$. Signs at ties are free. If there are m tied radii, including j , the median condition bounds the signed strict-group sum J by $|J| \leq mt$. The other $m - 1$ tie signs give offsets $-(m - 1)t, -(m - 3)t, \dots, (m - 1)t$; intervals of radius t around these offsets cover $[-mt, mt]$. Hence the signs can be chosen so that, for the five real numbers $x_i = \sigma_i r_i$ ($i \neq j$),

$$L = \sum x_i, \quad |L| \leq t, \quad t^2 + \sum x_i^2 = 1. \quad (27)$$

The minimized majorant is exactly

$$P_6 \sum_i |r_i^{-1} - r_i/t^2| = |te_4(x) - (L/t)e_5(x)|. \quad (28)$$

Indeed $e_4(x) = e_5(x) \sum 1/x_i$, and every tied sign contributes zero. We may discard all remaining sign and ordering restrictions on x .

Fix $L, R = \sum x_i^2$ and t . The maximum of the absolute affine coefficient functional in (28) on the compact real-rooted slice with fixed $e_1 = L$ and $e_2 = (L^2 - R)/2$ occurs with at most two distinct roots. This is the classical hyperbolic two-slice principle, directly covered by Riener-Schabert [8, Theorem 2.8 and Corollary 2.10]. For completeness, maximize either sign and select a maximizing quintic f with the largest number d of distinct roots. If $3 \leq d < 5$, write $f = hg$, with g containing each distinct root once. The perturbations $h(g + \epsilon) = f + \epsilon h$ preserve real-rootedness for both sufficiently small signs and preserve the first two coefficients because $\deg h \leq 2$. A nonzero change of the functional improves one sign. A zero change preserves the maximum but splits a repeated root, increasing d . Both are contradictions. If $d = 5$, perturb by ϵX ; both small signs preserve real-rootedness, while the nonzero coefficient t of e_4 improves one sign. Thus $d \leq 2$.

Label the two values a, b , with minority multiplicity $r \in \{1, 2\}$ and majority multiplicity $s = 5 - r$. The one-value case is included by $a = b$. If $b = 0$, at least three coordinates vanish and the objective is zero. Otherwise reversing every sign allows $b > 0$. Put

$$v = a/b, \quad \tau = t/b, \quad L = rv + s, \quad R = rv^2 + s, \quad P = v^r, \quad A = rv^{r-1} + sv^r.$$

(Here L, R are the rescaled quantities.) The normalized objective squared is

$$\frac{(\tau A - LP/\tau)^2}{(\tau^2 + R)^5}, \quad \tau > 0, \quad \tau^2 \geq L^2.$$

It suffices to prove, for every real v ,

$$f_r(v, X) = X(X + R)^5 - 1728(AX - LP)^2 \geq 0 \quad (X \geq L^2). \quad (29)$$

Write $X = L^2 + y$, $V = L^2 + R > 0$, and $f_r = \sum_{j=0}^6 c_j y^j$. Then

$$\begin{aligned} c_0 &= L^2 J, & J &= V^5 - 1728(LA - P)^2, \\ c_1 &= V^4(6V - 5R) - 3456AL(LA - P), & c_2 &= 5V^3(3V - 2R) - 1728A^2, \\ c_3 &= 10V^2(2V - R), & c_4 &= 5V(3V - R), \\ c_5 &= 6V - R, & c_6 &= 1. \end{aligned}$$

In particular $c_3, c_4, c_5, c_6 > 0$. For $r = 1$, putting $w = (v + 2)^2$ gives

$$J = 32w(w^4 + 30w^3 + 360w^2 + 1296w + 11664) \geq 0.$$

For $r = 2$,

$$J = 7776(v + 2)^2 Q(v + 1), \quad Q(w) = 1 - 2w + 8w^2 - 14w^3 + 22w^4 - 14w^5 + 8w^6 - 2w^7 + w^8.$$

For $w < 0$, $Q(w) > 0$ term by term, and $Q(0) = 1$. For $w > 0$, $Q(w)/w^4 = z^4 - 2z^3 + 4z^2 - 8z + 8$, where $z = w + w^{-1} \geq 2$; substitution $z = 2 + h$ gives $8 + 16h + 16h^2 + 6h^3 + h^4 > 0$. Thus $c_0 \geq 0$ in both cases.

The remaining five univariate positivity certificates are given explicitly below. For $r = 1$ they prove $c_1, c_2 > 0$ on $\mathbb{R}$; hence every coefficient of f_1 is nonnegative. For $r = 2$ they prove $c_1, c_3, \Delta = 4c_1c_3 - c_2^2 > 0$ on $\mathbb{R}$. Therefore

$$f_2 = c_0 + y(c_1 + c_2y + c_3y^2) + c_4y^4 + c_5y^5 + y^6 \geq 0 \quad (y \geq 0).$$

This proves (29), then the radial bound and (25). Sharpness follows from $z = (2, 2, -1, -1, -1, -1)$: $\sum |z_i|^2 = 12$ and $e_5(z) = -12$.

For the stable-law assertion, the existing ternary restriction supplies, separately for every complex direction w , real symmetric matrices A, C such that $R_5(w) = e_5(A + iC)$, $\text{Tr}(A + iC) = 0$ and $\|A + iC\|_F^2 = 2w^*Bw$. The six eigenvalues are centered and their squared-modulus sum is at most this Frobenius square by Schur triangularization. Consequently

$$|R_5(w)|^2 \leq (2w^*Bw)^5/1728.$$

For standard circular Gaussian Z , substitute $w = Q^{-1/2}Z$ and use $D = \frac{1}{3}Q^{-1/2}BQ^{-1/2}$. Gaussian coefficient orthogonality and the exact density normalization give

$$\|U_5\|_2^2 \leq \frac{6^5 5! h_5(D)}{1728 5! \binom{6}{5}} = \frac{3}{4} h_5(D).$$

Matrices may depend on w ; no global determinantal representation in arbitrarily many variables is used. Singular deficits cause no division. $\square$

Exact certificates on the full real line. The following table specifies complete Bernstein partitions for the five polynomials used above. The variables in the partition column are denoted w and belong to $[0, 1]$.

r	polynomial	degree	partition endpoints
1	c_1	10	$0, \frac{1}{2}, \frac{5}{8}, \frac{11}{16}, \frac{3}{4}, 1$
1	c_2	8	$0, \frac{1}{2}, \frac{3}{4}, 1$
2	c_1	10	$0, \frac{1}{2}, \frac{5}{8}, \frac{11}{16}, \frac{3}{4}, 1$
2	c_3	6	$0, \frac{1}{2}, \frac{3}{4}, 1$
2	$4c_1c_3 - c_2^2$	16	$0, \frac{1}{2}, \frac{5}{8}, \frac{3}{4}, 1$

Here is the exact verification method, implemented in the accompanying standard-library program `verify_complex_six.py`. Reconstruct each $p(v) = \sum_{i=0}^d a_i v^i$ from the displayed formulas. All a_i are nonnegative and $a_0 > 0$, proving positivity for $v \geq 0$. For $v < 0$ set $v = -w/(1 - w)$; the Bernstein controls of $(1 - w)^d p(-w/(1 - w))$ are

$$\beta_i = \frac{(-1)^i a_i}{\binom{d}{i}}, \quad 0 \leq i \leq d.$$

Apply exact rational midpoint de Casteljau subdivision, retaining both children, to the listed intervals. Every entry of every resulting control array is strictly positive. These are twenty full arrays, and their intervals cover $[0, 1]$. Since the Bernstein basis is nonnegative and sums to one, this certifies positivity for every negative real v , without a cutoff or sampling argument. The program also verifies the two exact factorizations of J . Thus the computational part consists only of finite integer and rational identities and strict sign checks.

Attainment by positive stable count laws. For $0 < \epsilon < 1/4$ consider

$$p_\epsilon(X, Y) = [(\tfrac{1}{2} + 2\epsilon)X + (\tfrac{1}{2} - 2\epsilon)Y]^2 [(\tfrac{1}{2} - \epsilon)X + (\tfrac{1}{2} + \epsilon)Y]^4.$$

This is a normalized stable product law with positive coefficients, means $(3, 3)$ and $q = (1/2, 1/2)$. Direct calculation gives

$$B = 6\epsilon^2 \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, \quad D = 4\epsilon^2 \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, \quad R_5 = -12\epsilon^5 (X - Y)^5.$$

The nonzero eigenvalue of D is $8\epsilon^2$, so

$$h_5(D) = 32768\epsilon^{10}, \quad \|U_5\|_2^2 = \frac{144\epsilon^{10}}{6} (2 + 2)^5 = 24576\epsilon^{10} = \tfrac{3}{4}h_5(D).$$

Thus the constant is attained within the stated count-law class.

Corollary 18. *For every degree $n \geq 6$, the fifth-order spectral bound has the additional branch*

$$\|U_5\|_2^2 \leq b_n h_5(D), \quad b_n = \frac{3125}{8} \frac{\binom{n}{5}}{(n-1)^5}.$$

This branch is sharp at $n = 6$; no optimality is asserted for $n > 6$.

Proof. Use the classical complex derivative-root contraction, in the credited derivative argument of Han–Niles–Weed [6, Section 1.1]. For centered $z \in \mathbb{C}^n$ with $\sum |z_i|^2 = n$, differentiate its monic root polynomial $n - 6$ times and normalize to obtain six roots w . Coefficient differentiation and the iterated contraction give

$$e_5(w) = \frac{6}{\binom{n}{5}} e_5(z), \quad \sum_{j=1}^6 |w_j|^2 \leq \frac{30}{n-1}.$$

Theorem 17 yields

$$|e_5(z)|^2 \leq \left(\frac{\binom{n}{5}}{6} \right)^2 \frac{(30/(n-1))^5}{1728} = b_n \binom{n}{5}.$$

Apply the same pointwise ternary/Schur and Gaussian transfer as above, now with

$$D = (2/n)Q^{-1/2}BQ^{-1/2},$$

to obtain the spectral assertion. □

This improves the previous degree-five derivative branch by the factor $3125/8192$. It does not resolve unrestricted higher-degree factor one: $b_7 = 21875/20736 > 1$ and $\lim_{n \rightarrow \infty} b_n = 625/192 > 1$. The branch can be combined by minimum with the other established bounds.

4.7 An exact rank-two matrix formulation

The degree-six, rank-two sharp bound has an exact finite-dimensional matrix formulation. The converse below realizes every admissible matrix pair by an actual positive stable count law, so this reformulation does not enlarge the problem to arbitrary symmetric tensors.

Theorem 19 (Exact rank-two matrix reformulation). *The following assertions are equivalent.*

1. Every admissible stable count law of degree 6 with rank $D = 2$ satisfies $\|U_5\|_2^2 \leq (3/4)h_5(D)$.
2. For a 6×6 matrix M , write $e_5(M) = [t] \det(tI_6 + M)$, the ordinary fifth elementary function of its eigenvalues. For every pair of real symmetric 6×6 matrices H, K with

$$\operatorname{Tr} H = \operatorname{Tr} K = \operatorname{Tr}(HK) = 0, \quad \operatorname{Tr} H^2 = \operatorname{Tr} K^2 = 1,$$

write

$$e_5(zH + wK) = \sum_{j=0}^5 c_j z^{5-j} w^j.$$

Then, for every $a, b \geq 0$,

$$S_{H,K}(a, b) := \sum_{j=0}^5 c_j^2 (5-j)! j! a^{5-j} b^j \leq \frac{5}{72} h_5(a, b), \quad h_5(a, b) = \sum_{j=0}^5 a^{5-j} b^j. \quad (30)$$

It is enough to establish the second assertion for $a, b > 0$. Both assertions hold by Theorem 17. The equivalence proof also realizes every admissible matrix pair by a positive stable count law.

Proof. First suppose the matrix assertion holds. Each centered coefficient R_j is constant along $\ker B$. To see this directly, fix $v \in \ker B$. The centered roots associated with $x + sv$ have squared norm $2x^\top Bx$, independent of real s , by (5). Their elementary coefficients are therefore uniformly bounded in s . The polynomial $R_j(x + sv)$ is consequently constant. This argument only requires a finite root-norm bound, not the desired spectral inequality.

Let $a, b > 0$ be the two nonzero eigenvalues of D , with orthonormal eigenvectors u_1, u_2 , and put $Q = \operatorname{diag}(q)$. Since $D\sqrt{q} = 0$, the vectors $Q^{-1/2}u_i$ are centered for L . The kernel descent just proved shows that all whitened R_j depend only on these two coordinates. The full ternary restriction

$$F(t, z, w) = p(t\mathbf{1} + Q^{-1/2}(zu_1 + wu_2))$$

is monic of degree six in t , hyperbolic in $(1, 0, 0)$, and has zero t^5 coefficient and quadratic coefficient $-3(az^2 + bw^2)$. The classical ternary representation used in Theorem 11 gives

$$F(t, z, w) = \det(tI + zA + wC)$$

with real symmetric matrices satisfying

$$\operatorname{Tr} A = \operatorname{Tr} C = \operatorname{Tr}(AC) = 0, \quad \operatorname{Tr} A^2 = 6a, \quad \operatorname{Tr} C^2 = 6b.$$

Thus $H = A/\sqrt{6a}$ and $K = C/\sqrt{6b}$ satisfy the matrix hypotheses.

For independent standard circular complex Gaussians Z, W , monomial orthogonality and the exact density identity give

$$\|U_5\|_2^2 = \frac{\mathbb{E}|e_5(\sqrt{6a}ZH + \sqrt{6b}WK)|^2}{5! \binom{6}{5}} = \frac{6^5}{720} S_{H,K}(a, b) = \frac{54}{5} S_{H,K}(a, b).$$

Here other Gaussian coordinates disappear by kernel descent, and $h_5(D) = h_5(a, b)$. Equation (30) proves the first assertion.

Conversely, fix any matrices in the second assertion and $a, b > 0$. Set

$$\begin{aligned} H_1 &= \sqrt{a} H, \\ H_2 &= -\frac{1}{2}\sqrt{a} H + \frac{1}{2}\sqrt{3b} K, \\ H_3 &= -\frac{1}{2}\sqrt{a} H - \frac{1}{2}\sqrt{3b} K. \end{aligned}$$

For sufficiently small $\epsilon > 0$, all three matrices $A_i = I/3 + \epsilon H_i$ are positive definite and sum to I . Define

$$p(X_1, X_2, X_3) = \det \left(\sum_{i=1}^3 X_i A_i \right).$$

This is homogeneous of degree six and normalized at $\mathbf{1}$. Its stability follows because $\sum_i (\Im X_i) A_i$ is positive definite when every $\Im X_i > 0$. Its coefficients are nonnegative by Cauchy–Binet applied to concatenated square-root columns of the A_i . They are in fact strictly positive: write $A_i = \delta I + B_i$ with a common $\delta > 0$ and $B_i \succeq 0$; the same expansion contains the positive contribution $\delta^6 (X_1 + X_2 + X_3)^6$.

The means are $t_i = \text{Tr } A_i = 2$, so $q_i = 1/3$. Direct second-coefficient comparison gives

$$B = \frac{\epsilon^2}{2} \text{Gram}(H_1, H_2, H_3), \quad D = B.$$

In the orthonormal H, K coordinates, the three frame vectors are $(\sqrt{a}, 0)$, $(-\sqrt{a}/2, \sqrt{3b}/2)$ and $(-\sqrt{a}/2, -\sqrt{3b}/2)$. Their frame operator is $\text{diag}(3a/2, 3b/2)$. Hence D has exactly two positive eigenvalues $(3\epsilon^2/4)a, (3\epsilon^2/4)b$.

Whitening multiplies the category coordinates by $\sqrt{3}$. An orthogonal change among independent circular Gaussian coordinates therefore makes the whitened fifth-coefficient matrix equal in law to

$$\epsilon \sqrt{9/2} (\sqrt{a} ZH + \sqrt{b} WK).$$

Consequently

$$\|U_5\|_2^2 = \frac{\epsilon^{10}(9/2)^5}{720} S_{H,K}(a, b), \quad h_5(D) = \epsilon^{10}(3/4)^5 h_5(a, b).$$

Their ratio is $(54/5)S_{H,K}(a, b)/h_5(a, b)$. The first assertion applied to this genuine rank-two stable law proves (30) for positive a, b . Polynomial continuity proves the nonnegative endpoints as well; both sides vanish at $a = b = 0$. $\square$

The realization retains the simultaneous matrix constraints. The coupled weighted inequality (30) is therefore a consequence of the full stable-law theorem, with the stronger coefficient $5/72$.

Corollary 20 (Two coupled pencil subclasses). *Let H, K satisfy the matrix hypotheses of Theorem 19. Suppose either*

1. *$\det(zH + wK)$ is identically zero as a polynomial in z, w ; or*
2. *there is a common orthogonal decomposition $\mathbb{R}^6 = V \oplus W$, with $\dim V = 4$, $\dim W = 2$, preserved by both matrices, and every restricted block $H|_V, K|_V, H|_W, K|_W$ has trace zero.*

Then, for all $a, b \geq 0$,

$$S_{H,K}(a, b) \leq \frac{120}{3125} h_5(a, b).$$

In particular, a degree-six rank-two stable law whose centered ternary pencil has either property satisfies

$$\|U_5\|_2^2 \leq \frac{1296}{3125} h_5(D) < h_5(D) \quad (D \neq 0).$$

Neither condition requires H and K to commute. The singular-pencil condition concerns the matrix polynomial, not a reduction of the covariance rank below two. The same count-law bound also holds, without a restriction on covariance rank, for $p = p_4 p_2$ where the normalized homogeneous stable count-law factors have degrees four and two and the same mean proportions q .

Proof. Let Z, W be independent standard circular complex Gaussians and set $M = \sqrt{a}ZH + \sqrt{b}WK$. The global orthonormality conditions give

$$\|M\|_F^2 = a|Z|^2 + b|W|^2.$$

In the first case, M has a zero eigenvalue. If it has at least two zero eigenvalues counted algebraically, $e_5(M) = 0$. Otherwise $e_5(M)$ is the product of its other five eigenvalues. Schur triangularization and AM–GM therefore give

$$|e_5(M)|^2 \leq \left(\frac{\|M\|_F^2}{5} \right)^5.$$

This argument does not require normality or a common kernel for the pencil.

In the second case write $M = A \oplus B$ with trace-free complex 4×4 and 2×2 blocks. Characteristic-coefficient multiplication gives $e_5(M) = e_3(A) \det B$. For the four eigenvalues of A , their zero sum permits the Hadamard parameterization

$$\lambda(A) = \frac{1}{2}(u + v + w, u - v - w, -u + v - w, -u - v + w).$$

It satisfies $e_3(A) = uvw$ and $\sum_j |\lambda_j(A)|^2 = |u|^2 + |v|^2 + |w|^2$. Thus, with $X = \|A\|_F^2$ and $Y = \|B\|_F^2$, Schur triangularization and AM–GM imply

$$|e_3(A)|^2 \leq \frac{X^3}{27}, \quad |\det B|^2 \leq \frac{Y^2}{4}.$$

Consequently

$$|e_5(M)|^2 \leq \frac{X^3 Y^2}{108} \leq \frac{(X + Y)^5}{3125} = \frac{\|M\|_F^{10}}{3125}.$$

The middle inequality is weighted AM–GM, with the maximum $X : Y = 3 : 2$ for fixed $X + Y$.

In either case complex monomial orthogonality and exponential moments give

$$S_{H,K}(a, b) = \mathbb{E}|e_5(M)|^2 \leq \frac{\mathbb{E}(a|Z|^2 + b|W|^2)^5}{3125} = \frac{120}{3125} h_5(a, b).$$

The exact factor $54/5$ in Theorem 19 gives the count-law conclusion. Positive $h_5(D)$ makes the final inequality strict whenever $D \neq 0$.

For the last assertion, let B_4, B_2 be the two factors' covariance deficits. Their common $L = q \cdot x$ makes the centered product coefficient exactly $R_5 = R_3^{(4)} R_2^{(2)}$, and covariance additivity gives $B = B_4 + B_2$. On each complex direction w , apply the justified ternary representation separately to each factor. The size-four estimate above and the quadratic top bound give

$$|R_3^{(4)}(w)|^2 \leq \frac{8}{27} (w^* B_4 w)^3, \quad |R_2^{(2)}(w)|^2 \leq (w^* B_2 w)^2.$$

With $X = w^* B_4 w$ and $Y = w^* B_2 w$, weighted AM–GM yields

$$|R_5(w)|^2 \leq \frac{32}{3125} (w^* B w)^5.$$

For total degree six, $D = \frac{1}{3} Q^{-1/2} B Q^{-1/2}$. Substituting $w = Q^{-1/2} Z$ and using the exact fifth-density identity therefore gives

$$\|U_5\|_2^2 \leq \frac{32 \cdot 3^5}{3125 \cdot 6} h_5(D) = \frac{1296}{3125} h_5(D).$$

The separate ternary matrices may depend on w ; no global determinantal model in an arbitrary number of variables is used. $\square$

The second condition retains a genuine common block constraint without imposing simultaneous diagonalization. The first includes the case $R_6 \equiv 0$ in the centered expansion. These are sufficient subclasses; no assertion that all rank-two pencils satisfy either condition is made. The proof uses classical characteristic identities, Schur triangularization and AM–GM, rather than a new general trace-word positivity theorem. The product extension requires matching mean proportions of its two factors; arbitrary products with different means are not covered.

5 A nonfactorizable cubic obstruction to a Frobenius-power extension

The preceding spectral-square bound cannot generally be replaced by the stronger Frobenius power available for permutation-product models. We give actual stable count laws, rather than only a generic tensor obstruction.

Theorem 21. *For every integer $r \geq 5$, there are degree-three real-stable count laws P_r, P_r^0 on $2r$ categories with strictly positive coefficients, identical means and covariance, and nonzero D eigenvalue $2\delta^2/(3r)$ repeated r times, where any fixed $0 < \delta \leq 1/4$ is allowed. With $c = 1/4$,*

$$x = 2\delta^2/3, \quad s_2 = 4\delta^4/(9r), \quad \|U_3(P_r)\|_2^2 \geq \frac{c^2\delta^6}{1000r} = \frac{27c^2}{8000}s_2x.$$

The law P_r^0 has $U_3 = 0$. Consequently no dimension-uniform finite K gives $\|U_3\|_2^2 \leq Ks_2^{3/2}$ on the full stable class, even at fixed degree three and fixed $x > 0$. Both laws are nonfactorizable into nonnegative categorical linear forms.

Norm-one real symmetric cubic tensors with squared Frobenius norm of order r^2 are established tensor-norm phenomena; see Cao, He, Li and Wang [21, Theorem 4.6 and Corollary 4.9]. We include a self-contained probabilistic existence proof with sufficient constants. This is an analytic existence argument, not a numerical search or an algorithm outputting a tensor in every dimension.

Proof. Let $N = \dim \text{Sym}^3(\mathbb{R}^r) = \binom{r+2}{3}$ and choose an isotropic Gaussian G in that Euclidean space, with the ordered Frobenius metric. Then $\|G\|_F^2$ has mean N , variance $2N$, and

$$\Pr(\|G\|_F^2 < N/2) \leq 8/N < 1/4 \quad (r \geq 5).$$

For a fixed unit u , $G(u, u, u)$ is a standard normal because $\|u^{\otimes 3}\|_F = 1$. A sphere $1/6$ -net has at most 13^r points, by disjoint radius- $1/12$ ball volume comparison. Banach's theorem identifies the diagonal supremum M with the full trilinear norm; telescoping gives $|G(u, u, u) - G(v, v, v)| \leq 3M\|u - v\|$, hence $M \leq 2 \max_{\text{net}} |G(v, v, v)|$. The Gaussian tail and union bound give

$$\Pr(M > \sqrt{32r}) \leq 2 \cdot 13^r e^{-4r} < 2e^{-r} < 1/4.$$

Here $\log 13 < 3$ and $e^5 > 8$, both certified by lower Taylor sums. A realization outside both failures has $M > 0$ and, upon normalization $A = G/M$,

$$\sup_{\|u\|=1} |A(u, u, u)| = 1, \quad \|A\|_F^2 \geq N/(64r) \geq r^2/384 \geq r^2/1000. \quad (31)$$

Put $C(u) = A(u, u, u)$ and $F_c(z, u) = z^3 - z\|u\|^2 + cC(u)$, with $c = 1/4$. The depressed-cubic discriminant is

$$4\|u\|^6 - 27c^2C(u)^2 \geq (4 - 27/16)\|u\|^6 \geq 0.$$

Thus F_c is homogeneous and hyperbolic in $e = (1, 0)$ with $F_c(e) = 1$, also at $u = 0$; the same holds for $c = 0$. Let $m = 2r$ and index u_i by $+e_1, -e_1, \dots, +e_r, -e_r$. Then $\sum_i u_i = 0$ and $\sum_i u_i u_i^\top = 2I$. Set $v_i = (1, \delta u_i)$. Every real root of $F_c(z, \delta u_i)$ has modulus less than 2δ : for $|z| \geq 2\delta$, $|z^3 - \delta^2 z| \geq 6\delta^3 > c\delta^3 |C(u_i)|$. Therefore the characteristic roots of v_i relative to e are positive, being one minus these roots. The classical Gårding cone theorem gives an open convex cone of such directions and hyperbolicity in every direction in it [20, Fact 2.7].

Define

$$p_c(X) = F_c\left(\frac{\sum_i X_i}{m}, \frac{\delta}{m} \sum_i X_i u_i\right) = m^{-3} F_c\left(\sum_i X_i v_i\right).$$

It is normalized at $\mathbf{1}$. If all $\Im X_i > 0$, the imaginary vector $b = \sum_i (\Im X_i) v_i$ belongs to that cone. With $a = \sum_i (\Re X_i) v_i$, the polynomial $t \mapsto F_c(a + tb)$ has only real roots and nonzero leading coefficient. Hence $F_c(a + ib) \neq 0$: p_c is real stable. This deduction is the standard hyperbolicity-to-tube argument, not a coordinatewise discriminant shortcut.

Coefficient positivity is separate and explicit. The symmetric polarization at v_i, v_j, v_k equals

$$1 - \frac{\delta^2}{3}(u_i \cdot u_j + u_i \cdot u_k + u_j \cdot u_k) + c\delta^3 A(u_i, u_j, u_k) \geq 1 - \delta^2 - c\delta^3 > 0.$$

Banach bounds the last absolute value by one even with repeated indices. Each count monomial coefficient is a positive multinomial multiplicity times this expression divided by m^3 . Thus p_c and p_0 are genuine strictly positive count laws.

Write $L = \sum X_i/m$ and $U = \sum X_i u_i$. Exactly,

$$p_c = L^3 - \frac{\delta^2}{m^2} \|U\|^2 L + \frac{c\delta^3}{m^3} C(U).$$

Since $U(\mathbf{1}) = 0$, the last term has zero first and second derivatives there. Both laws have $t_i = 3/m$ and $B = (\delta^2/m^2) \text{Gram}(u_i)$, independently of c ; the displayed terms are their centered R_2, R_3 . The Gram spectrum is two repeated r times and zero otherwise. Thus $D = (2m/3)B$ has the asserted spectrum.

Since $q_i = 1/m$ and $\binom{3}{3} = 1$, the exact ordered cubic density identity and $UU^\top = 2I$ give

$$\|U_3\|_2^2 = \frac{c^2 \delta^6}{m^3} 2^3 \|A\|_F^2 = \frac{c^2 \delta^6}{r^3} \|A\|_F^2.$$

Equation (31) proves the lower bound. Since $s_2^{3/2} = 8\delta^6/(27r^{3/2})$, its ratio is at least $27c^2\sqrt{r}/8000$ and diverges.

Finally a product of three normalized categorical forms with vectors w_a and mean row q has $B = \frac{1}{2} \sum_{a=1}^3 (w_a - q)(w_a - q)^\top$ and rank at most two, because the deviations sum to zero. Our B has rank $r \geq 5$ for both laws. Any nonnegative linear factors can be normalized, so neither polynomial admits such a factorization. $\square$

The relative quadratic chi-square increment here is at least $3c^2\delta^2/1000$, since $\chi^2(P_r^0\|M) = a_3 s_2$ and the cubic energy adds orthogonally. It need not vanish as $s_2 \rightarrow 0$ at fixed trace; absolute chi-square still tends to zero by Corollary 9. This does not infer a TV lower bound from L^2 , nor contradict the product-subclass results of Han–Niles–Weed. The higher interior bound $\|U_k\|_2^2 \leq h_k(\lambda(D))$ for general $n \geq 7$ and $5 \leq k < n$ remains open, where h_k is complete homogeneous. Generic real tensors cannot prove that target: $\Re(X + iY)^4 = X^4 - 6X^2Y^2 + Y^4$ has diagonal norm one and ordered squared tensor norm eight, whereas $h_4(1, 1) = 5$. This last tensor is not asserted to arise from a stable law.

6 Coefficient bounds and finite-degree certificates

The coefficient extremum reduction used next is classical: see Riener and Schabert [8, Theorem 2.8 and Corollary 2.10], as well as the real-vector reduction in [5]. We include the needed special proof.

Lemma 22. *For fixed $n \geq 3$ and $3 \leq k \leq n$, the maximum of $|e_k(\lambda)|$ on $\sum_i \lambda_i = 0$, $\sum_i \lambda_i^2 = 2$ is attained at a vector with two distinct values.*

Proof. The slice is nonempty and compact. Maximize either sign of e_k , and among maximizers select one with the largest number d of distinct roots of $f(z) = \prod_i (z - \lambda_i)$. If $3 \leq d < n$, write $f = hg$, where g contains each distinct linear factor once and h retains multiplicity minus one. For either sufficiently small sign of ε , $g + \varepsilon$ has d simple real roots near the old roots. Thus $f + \varepsilon h = h(g + \varepsilon)$ is real-rooted and preserves the first two coefficients, since $\deg h = n - d \leq n - 3$. If the target coefficient of h is nonzero, one sign improves the objective. Otherwise the objective is constant. Every repeated old root remains through h and also supplies a new nearby root through $g + \varepsilon$; no new root equals an old one. Taking disjoint neighborhoods of distinct old roots shows the distinct-root count strictly increases, a contradiction. If $d = n$, perturbing by εz^{n-k} preserves the first two coefficients and real-rootedness for both small signs, but changes the objective. Hence $d \leq 2$. The positive square sum rules out $d = 1$. Apply this argument to both signs to obtain the absolute-value assertion. $\square$

6.1 A uniform real coefficient refinement

The following mass-budget estimate is an elementary rearrangement inequality for the arcsine weight. The contour, binary reduction and enlarged Beta integral that follow are the ingredients of Han and Niles-Weed [5, Appendix B.1]; the change is to use this estimate instead of their endpoint/interior split and to retain the normalization factors.

Lemma 23. *If f is measurable and $0 \leq f \leq 1$ on $[-1, 1]$ and $A = \int_{-1}^1 f(t) dt$, then*

$$\left(\frac{1}{\pi} \int_{-1}^1 \frac{f(t)}{\sqrt{1-t^2}} dt \right)^2 \leq \frac{A}{2}.$$

Proof. The endpoints $A = 0, 2$ are immediate. Otherwise put $b = 1 - A/2$ and $E = [-1, -b] \cup [b, 1]$. Its length is A . For $w(t) = (1 - t^2)^{-1/2}$ one has $(\mathbf{1}_E - f)(w - w(b)) \geq 0$ on both E and its complement. Since $\int (\mathbf{1}_E - f) = 0$, integration gives $\int f w \leq \int_E w = 2 \arccos b$. For $0 \leq u \leq \pi/2$, the chord bound for sine on $[0, \pi/4]$ gives $\sin(u/2) \geq \sqrt{2}u/\pi$, so $1 - \cos u \geq 4u^2/\pi^2$. Apply this with $u = \arccos(1 - A/2)$ to obtain the assertion. $\square$

Theorem 24. *For every centered real vector z with $\sum_i z_i^2 = n$,*

$$|e_k(z)|^2 \leq C_0 \binom{n}{k} \quad (0 \leq k \leq n), \quad C_0 = \frac{\pi e^{1/4}}{\sqrt{2}} < \frac{20}{7}. \quad (32)$$

Proof. The normalized domain is empty at $n = 1$. The orders zero, one and two follow from $e_0 = 1, e_1 = 0, e_2 = -n/2$; order n also follows from AM–GM on $\prod z_i^2$. By Lemma 22 and scaling, the other orders reduce to a vector with r entries $\sqrt{s/r}$ and s entries $-\sqrt{r/s}$, where $r + s = n$. For $1 \leq k \leq n - 1$ put

$$\rho = \sqrt{\frac{k}{n-k}}, \quad v_0 = \frac{k(n-k)}{n}, \quad H_0 = \frac{(1 + \rho^2)^{n/2}}{\rho^k}.$$

Define, for $-1 \leq t \leq 1$,

$$f(t) = \frac{(1 + \rho^2 s/r + 2\rho\sqrt{s/r}t)^{r/2}(1 + \rho^2 r/s - 2\rho\sqrt{r/s}t)^{s/2}}{(1 + \rho^2)^{n/2}}.$$

The affine brackets are nonnegative. Their weighted average with weights $r/n, s/n$ is $1 + \rho^2$, so weighted AM–GM gives $0 \leq f \leq 1$. Cauchy’s coefficient formula, followed by the triangle inequality and $t = \cos \theta$, yields

$$|e_k(z)| \leq \frac{H_0}{\pi} \int_{-1}^1 \frac{f(t)}{\sqrt{1-t^2}} dt.$$

Let t_- and t_+ be the zeros of the first and second brackets. They satisfy $t_- \leq -1$ and $t_+ \geq 1$ by AM–GM. The affine substitution $t = t_- + (t_+ - t_-)u$, extending the nonnegative unweighted integrand to this larger interval, gives

$$A = \int_{-1}^1 f(t) dt \leq \frac{1 + \rho^2}{2\rho} \frac{n^{n/2+1}}{r^{(r+1)/2}s^{(s+1)/2}} \frac{\Gamma(r/2 + 1)\Gamma(s/2 + 1)}{\Gamma(n/2 + 2)}.$$

Here the Beta integral is $\int_0^1 u^{r/2}(1-u)^{s/2} du$. To bound it, use the standard Stirling inequalities, for $x > 0$,

$$\sqrt{2\pi x}(x/e)^x \leq \Gamma(x+1) \leq \sqrt{2\pi x}(x/e)^x e^{1/(12x)}.$$

Apply upper bounds at $r/2, s/2$ and the lower bound at $n/2$, keeping $\Gamma(n/2+2) = (n/2+1)\Gamma(n/2+1)$. Cancellation, and $(1 + \rho^2)/\rho = \sqrt{n/v_0}$, give

$$A \leq \frac{\sqrt{\pi}}{\sqrt{v_0}} \frac{n}{n+2} e^{(1/r+1/s)/6}.$$

Independently, factorial Stirling bounds at $k, n-k, n$ give

$$\frac{H_0^2}{\binom{n}{k}} \leq \sqrt{2\pi v_0} e^{(1/k+1/(n-k))/12}.$$

Combining both estimates with Lemma 23,

$$\frac{|e_k(z)|^2}{\binom{n}{k}} \leq \frac{\pi}{\sqrt{2}} \frac{n}{n+2} \exp\left(\frac{1/r+1/s}{6} + \frac{1/k+1/(n-k)}{12}\right).$$

Each reciprocal sum is at most $n/(n-1)$, since its positive integer arguments sum to n . The exponent is therefore at most $n/[4(n-1)]$. Since

$$\log(1 + 2/n) \geq \frac{2}{n+2} \geq \frac{1}{4(n-1)} \quad (n \geq 2),$$

the last bound is at most C_0 .

For completeness we certify the rational ceiling in (32). The elementary identity

$$\frac{22}{7} - \pi = \int_0^1 \frac{x^4(1-x)^4}{1+x^2} dx > 0$$

follows by division: its integrand equals $x^6 - 4x^5 + 5x^4 - 4x^2 + 4 - 4/(1+x^2)$. Also $\sqrt{2} > 140/99$ since $19600 < 19602$. The exponential series has consecutive-term ratio at most $1/20$ after its order-four term, hence

$$e^{1/4} \leq \sum_{j=0}^3 \frac{4^{-j}}{j!} + \frac{4^{-4}}{4!(1-1/20)} < \frac{257}{200}.$$

Consequently

$$C_0 < \frac{22}{7} \frac{257}{200} \frac{99}{140} = \frac{559746}{196000} < \frac{20}{7}.$$

These rational comparisons are also checked by the accompanying code. $\square$

Scaling the roots in (5) and applying Lemma 5 now gives the all-order estimate

$$\mathbb{E}_M U_k^2 \leq C_0 (2W)^k \leq \frac{20}{7} (2W)^k. \quad (33)$$

It uses no assertion of scalar constant one at arbitrary order.

6.2 Small-degree scalar certificates

Lemma 25. *For $2 \leq n \leq 50$, every centered real vector with squared norm n satisfies $|e_k|^2 \leq \binom{n}{k}$ for every $0 \leq k \leq n$.*

Proof. Orders zero and one are immediate, and $e_2 = -n/2$ gives squared ratio $n/[2(n-1)] \leq 1$. For $k \geq 3$, apply Lemma 22. The raw binary roots s repeated r times and $-r$ repeated s times, $r + s = n$, have squared norm nrs . Their coefficient is

$$F_{n,r,k} = \sum_a \binom{r}{a} \binom{s}{k-a} s^a (-r)^{k-a}.$$

The standalone program checks the integer inequality

$$F_{n,r,k}^2 \leq \binom{n}{k} (rs)^k$$

for every $2 \leq n \leq 50$, every $1 \leq r < n$ and every $2 \leq k \leq n$. All 40,425 comparisons pass. Each label and total count is checked; only exact integers decide the inequalities. Dividing the roots by $\sqrt{rs}$ and applying the same- n extremum lemma proves the full-real statement. This is a direct scalar certificate, not an inference from a sufficient finite-degree TV criterion. $\square$

7 Exceptional profiles and fixed low orders

We normalize centered real root vectors by $\sum_i z_i^2 = n$. If

$$\frac{|e_k(z)|^2}{\binom{n}{k}} \leq b_k, \quad (34)$$

then scaling (5) and applying Lemma 5 gives $\mathbb{E}_M U_k^2 \leq 2^k b_k W^k$. All root extrema below use Lemma 22 one coefficient at a time; their maximizing profiles need not coincide.

7.1 Discrete exceptional profiles

Lemma 26. *If a centered, variance-normalized two-valued vector has either root value with multiplicity one or two, then (34) holds with $b_k = 1$ at every order.*

Proof. Negation and permutation allow us to treat the smaller specified block. For multiplicity one the values are $\sqrt{n-1}$ once and $-1/\sqrt{n-1}$ repeated $n-1$ times. Expansion gives

$$e_k = (-1)^k (1-k) \binom{n}{k} (n-1)^{-k/2}, \quad \frac{e_k^2}{\binom{n}{k}} = \frac{(k-1)^2 \binom{n}{k}}{(n-1)^k}.$$

The ratios at $k=0,1$ are one and zero. At $k=2$ the ratio is $n/[2(n-1)] \leq 1$. For fixed $k \geq 3$, regard the last expression as a function of real $n \geq k$. Its logarithmic derivative is

$$-\frac{1}{n(n-1)} + \sum_{j=2}^{k-1} \frac{j-1}{(n-j)(n-1)} > 0,$$

because the $j=2$ term exceeds the negative term. It therefore increases to $(k-1)^2/k! \leq 2/3$. This last sequence starts at $2/3$ for $k=3$ and decreases thereafter, since its next-term ratio is $k^2/[(k+1)(k-1)^2] < 1$.

For multiplicity two, assume $n \geq 4$; the case $n=3$ is already multiplicity one after reflection. The values are $\sqrt{(n-2)/2}$ twice and $-\sqrt{2/(n-2)}$ repeated $n-2$ times. Direct expansion gives

$$\frac{e_k^2}{\binom{n}{k}} = \binom{n}{k} \left(\frac{2}{n-2} \right)^k F(n, k)^2, \quad F(n, k) = \frac{(k-1)((k-4)n+4)}{4(n-1)}. \quad (35)$$

The orders zero and one are immediate. At order two the general centered identity gives ratio $n/[2(n-1)] \leq 1$. For order three, two-valued extrema have $e_3 = nc/3$, with $c = (s-r)/\sqrt{rs}$, and $c^2 \leq (n-2)^2/(n-1)$. Hence the ratio is at most $\frac{2}{3}n(n-2)/(n-1)^2 < 1$. For order four, the centered fourth-moment bound proved below gives ratio at most $3n(n-2)/[8(n-1)(n-3)] \leq 1$ for $n \geq 4$; the cleared difference $5n^2 - 26n + 24$ is zero at four and increases thereafter.

For $5 \leq k \leq 11$, clearing positive denominators in (35) reduces the assertion to

$$16k!(n-1)^2(n-2)^k - 2^k(n)_k(k-1)^2((k-4)n+4)^2 \geq 0. \quad (36)$$

Put $n = k + x$. The companion exact program expands these seven integer polynomials. Every coefficient is strictly positive; their minimum coefficients, in order $k = 5, \dots, 11$, are

$$1408, 5120, 39168, 444416, 4986880, 55074816, 628633600.$$

The program constructs both products in (36) by integer linear-factor multiplication and checks every coefficient, so this proves the inequality for all real $x \geq 0$, not just a finite list of n .

For $n \geq k \geq 12$, the factor F is nonnegative and is at most $nk(k-1)/[4(n-1)] \leq (3/11)k^2 < k^2/3$. Also $\binom{n}{k} \leq n^k/k!$ and

$$\left(\frac{n}{n-2} \right)^k \leq \left(\frac{k}{k-2} \right)^k < 9.$$

Indeed $t \log(t/(t-2))$ decreases for $t > 2$, and at twelve $6^{12} < 9 \cdot 5^{12}$. Thus the ratio in (35) is at most $2^k k^4/k! < 1$. This holds at twelve because $2^{12} 12^4 < 12!$ and decreases thereafter, since its next-term ratio is $2(k+1)^3/k^4 < 1$. This completes the all-degree exceptional-profile proof. $\square$

7.2 Orders zero through four

Lemma 27. *For every centered real vector with squared norm n , $|e_k|^2 \leq \binom{n}{k}$ for $0 \leq k \leq \min(4, n)$.*

Proof. Orders zero and one are immediate. At order two $e_2 = -n/2$, so its squared ratio is $n/[2(n-1)] \leq 1$ for $n \geq 2$. At order three, Lemma 22 gives a binary extremizer with $c = (s-r)/\sqrt{rs}$ and $e_3 = nc/3$. Since $rs \geq n-1$, $c^2 \leq (n-2)^2/(n-1)$, and the squared ratio is at most $2n(n-2)/[3(n-1)^2] < 1$.

For order four write $p_h = \sum_i z_i^h$. Newton's identity gives $e_4 = (n^2 - 2p_4)/8$. Cauchy on the other $n-1$ coordinates implies $z_i^2 \leq n-1$, whence $n \leq p_4 \leq n(n-1)$ and $|e_4| \leq n(n-2)/8$. Its squared ratio is at most

$$\frac{3n(n-2)}{8(n-1)(n-3)} \leq 1 \quad (n \geq 4),$$

since the cleared difference $5n^2 - 26n + 24$ equals $14y + 5y^2$ at $n = 4 + y$. $\square$

7.3 Fixed orders through twelve on an infinite multiplicity domain

We next prove a scalar coefficient-one estimate uniformly in both integer root multiplicities. It supplies the scalar estimates needed at orders five through twelve. The finite polynomial computations below, including all exceptional rows, are generated by the accompanying standalone code.

Lemma 28. *For every real vector z with $\sum_i z_i = 0$, $\sum_i z_i^2 = n$, and $5 \leq k \leq \min(12, n)$,*

$$|e_k(z)|^2 \leq \binom{n}{k}.$$

Consequently, for every law under consideration,

$$\mathbb{E}_M U_k^2 \leq 2^k W^k, \quad 7 \leq k \leq \min(12, n). \quad (37)$$

Proof. By Lemma 22, it suffices to treat two-valued profiles. Reflect and permute so their multiplicities satisfy $s \geq r$, $n = r + s$. Let $E_k(r, s)$ be the elementary coefficient of the integer root vector consisting of r copies of s and s copies of $-r$. Its squared norm is nrs , so the desired inequality is precisely

$$D_k(r, s) = (n)_k (rs)^k - k! E_k(r, s)^2 \geq 0. \quad (38)$$

For each $k = 5, \dots, 12$, put $a = \lceil k/2 \rceil$, $r = a + u$, $s = a + u + x$. Newton's identities reconstruct the exact polynomial E_k from

$$p_j = rs^j + s(-r)^j, \quad E_0 = 1, \quad E_h = \frac{1}{h} \sum_{j=1}^h (-1)^{j-1} E_{h-j} p_j.$$

The program expands (38) under this substitution and checks that every coefficient in u, x is nonnegative. The numbers of nonzero coefficients, in increasing order of k , are

$$121, 168, 225, 288, 361, 440, 529, 624.$$

Thus (38) holds for the whole real quadrant $u, x \geq 0$, covering every integer $s \geq r \geq \lceil k/2 \rceil$, with no upper cutoff on either multiplicity or on n .

The construction also has an independent exact polynomial-identity check. Since $p_1 = 0$, each Newton expansion term in E_k has at most $\lfloor k/2 \rfloor$ power-sum factors. Its total degree is at most $k + \lfloor k/2 \rfloor$ and its x -degree at most k . Therefore D_k has u -degree at most $3k$ and x -degree at most

2k. On the complete grid $0 \leq u \leq 3k$, $0 \leq x \leq 2k$, the program independently compares the polynomial with the direct formula

$$E_k(r, s) = \sum_j \binom{r}{j} \binom{s}{k-j} s^j (-r)^{k-j}.$$

These 4068 exact comparisons determine the eight polynomials by interpolation. Coefficient non-negativity, rather than those finite evaluations alone, proves the infinite-domain inequality.

It remains to cover $r < \lceil k/2 \rceil$. Multiplicities one and two are already covered by Lemma 26. The remaining pairs are exactly

k	7	8	9	10	11	12
r	3	3	3, 4	3, 4	3, 4, 5	3, 4, 5.

We give the cleared polynomial construction for these twelve cases. Write

$$E_k = (-r)^k \binom{n}{k} \frac{N_{r,k}(n)}{r^r(n)_r},$$

$$N_{r,k}(n) = \sum_{j=0}^{\min(r,k)} (-1)^j \binom{r}{j} n^j r^{r-j} (k)_j (n-j)_{r-j}.$$

To derive this formula, express the generating function as $(1-rT)^n [1+nT/(1-rT)]^r$ and compare coefficients; then use $\binom{n-j}{k-j} / \binom{n}{k} = (k)_j / (n)_j$. Dividing by $\sqrt{rs}$ normalizes the roots to squared norm n . Clearing positive denominators in the scalar inequality gives

$$k! r^{2r} (n)_r^2 (n-r)^k - r^k (n)_k N_{r,k}(n)^2 \geq 0. \quad (39)$$

For every listed pair the code substitutes $n = \max(k, 2r) + x$, expands both products, and checks that all coefficients are nonnegative. Since $x \geq 0$ is exactly the admissible range, this covers all remaining degrees. It independently checks the displayed coefficient factorization against the direct binomial formula on a degree-sized grid. The two-root reduction and Lemma 5 finish the proof. $\square$

7.4 Scalar constant one in a uniform central region

The binary contour modulus has more structure than the area bound alone records: its logarithm is concave. Retaining its interval superlevel sets and splitting sparse from dense multiplicities yields an infinite central scalar-one region without extending a finite-order table.

Lemma 29. *Let $f : [-1, 1] \rightarrow [0, 1]$ be measurable with interval strict superlevel sets up to null sets. Put*

$$A = \int_{-1}^1 f(t) dt, \quad I = \frac{1}{\pi} \int_{-1}^1 \frac{f(t)}{\sqrt{1-t^2}} dt.$$

If $0 < A \leq 3/10$, then $I \leq \Phi(A)$, where $\Phi(a) = \arccos(1-a)/\pi$, and

$$I^2 \leq \frac{2A}{\pi^2} (1 + A/10)^2. \quad (40)$$

The assertions hold trivially when $A = 0$.

Proof. The weight $w(t) = (1 - t^2)^{-1/2}$ is convex, with $w''(t) = (1 + 2t^2)/(1 - t^2)^{5/2} > 0$ in the open interval. Its integral over an interval of fixed length a is convex in the interval's left endpoint. Its maximum is thus attained at either endpoint interval; symmetry gives the same value, $\pi\Phi(a)$, at both. Integrable endpoint singularities follow by limits.

Let $\ell(h)$ be the length of $\{f > h\}$. Layer cake gives $A = \int_0^1 \ell(h) dh$ and $I \leq \int_0^1 \Phi(\ell(h)) dh$. Although Φ is concave only on $[0, 1]$ and convex on $[1, 2]$, its tangent T_A at every $0 < A \leq 3/10$ majorizes it throughout $[0, 2]$. Concavity proves this on $[0, 1]$. On $[1, 2]$, tangent minus Φ is concave, so it suffices to check both endpoints. The endpoint one is already covered. Also

$$T_A(2) = \Phi(A) + (2 - A)\Phi'(A), \quad \frac{d}{dA}T_A(2) = (2 - A)\Phi''(A) < 0.$$

At $A = 3/10$ its value is $(\arccos(7/10) + \sqrt{17/3})/\pi > 1$: $\cos(77/100) \geq 1 - (77/100)^2/2 > 7/10$, $\sqrt{17/3} > 119/50$, and their lower bounds sum to $63/20 > 22/7 > \pi$. Thus $T_A(2) > \Phi(2)$. Integrating the global tangent and using the mean of ℓ proves $I \leq \Phi(A)$ without assuming global concavity.

For $0 \leq v \leq 3/20$,

$$(1 + 3v/5)^2(1 - v) - 1 = v(1/5 - 21v/25 - 9v^2/25) \geq 0,$$

since the decreasing bracket equals $659/10000 > 0$ at $3/20$. Squaring positive factors and integrating gives $\arcsin x \leq x + x^3/5$ for $x^2 \leq 3/20$. Substituting $\arccos(1 - A) = 2 \arcsin \sqrt{A/2}$ proves (40). $\square$

Lemma 30. *For a two-valued normalized centered profile with multiplicities r, s , scalar one holds whenever $k \geq 8 \min(r, s)$.*

Proof. Reflect so $r \leq s$, and put $a = \sqrt{s/r}$, $b = -\sqrt{r/s}$, $n = r + s$. Expanding the r exceptional factors relative to $(1 + bT)^n$ gives

$$e_k = b^k \binom{n}{k} F_r(n, k), \quad F_r(n, k) = \sum_{j=0}^r \binom{r}{j} (-n/r)^j \frac{(k)_j}{(n)_j},$$

where terms with $j > k$ vanish. Because $k \leq n$, each falling-factorial ratio is at most $(k/n)^j$, so $|F_r| \leq (1 + k/r)^r$ and

$$\frac{e_k^2}{\binom{n}{k}} \leq \frac{r^k}{k!} \left(\frac{n}{n-r} \right)^k (1 + k/r)^{2r}.$$

For $n \geq k \geq 2r$, the middle factor is at most 4^r : it decreases in n , and the resulting function $t \log(t/(t-r))$ decreases for $t > r$, so its value at $t = k$ is at most that at $2r$. With $k! \geq (k/e)^k$, $e < 3$ and $\gamma = k/r \geq 8$, the last bound is at most

$$[4(3/\gamma)^\gamma(1 + \gamma)^2]^r.$$

The bracket decreases for $\gamma \geq 8$, since its log derivative is $\log(3/\gamma) - 1 + 2/(1 + \gamma) < 0$. At eight it equals $2125764/16777216 < 1$. This proves the assertion in every degree. $\square$

Theorem 31. *Every centered real vector z with $\sum_i z_i^2 = n$ satisfies*

$$|e_k(z)|^2 \leq \binom{n}{k} \quad \text{if } \min(k, n - k) \geq 80.$$

More precisely, for binary profiles with $r, s \geq 11$ and $v_0 = k(n - k)/n \geq 40$, the squared normalized ratio is at most $3150873/3171875 < 1$.

Proof. For the joint binary assertion, use the modulus f , area A and factor H_0 from Theorem 24. The logarithm of f is a sum of positive multiples of logarithms of positive affine functions on $(-1, 1)$, hence is concave. Therefore f has interval superlevel sets, including cases with endpoint zeros.

The exact Beta-area and factorial bounds already proved give

$$A \leq \sqrt{\frac{\pi}{v_0}} \frac{n}{n+2} e^{E_a}, \quad \frac{H_0^2}{\binom{n}{k}} \leq \sqrt{2\pi v_0} e^{E_b},$$

where $E_a = (1/r + 1/s)/6 \leq 1/33$ and $E_b = 1/(12v_0) \leq 1/480$. Their sum is at most $57/1760 < 1/30$. The estimate $e^x \leq (1-x)^{-1}$ gives $e^{E_a} \leq 33/32$ and $e^{E_a+E_b} < 30/29$. Thus

$$A^2 < \frac{22}{7} \left(\frac{33}{32} \right)^2 \frac{1}{40} = \frac{11979}{143360} < \frac{9}{100}.$$

So $A < 3/10$ and Lemma 29 applies. Multiplying its bound with the two displayed estimates yields

$$\begin{aligned} \frac{|e_k(z)|^2}{\binom{n}{k}} &\leq \frac{2\sqrt{2}}{\pi} \frac{n}{n+2} e^{E_a+E_b} (1 + A/10)^2 \\ &< \frac{792}{875} \frac{30}{29} \left(\frac{103}{100} \right)^2 = \frac{3150873}{3171875} < 1. \end{aligned}$$

Here $\sqrt{2} < 99/70$ and $\pi > 25/8$; the first 64 terms of the alternating arctangent series give a rational lower bound on π exceeding $25/8$. The accompanying code checks all rational comparisons.

Now let $\min(k, n-k) \geq 80$. Then $v_0 \geq \min(k, n-k)/2 \geq 40$ and $k \geq 80$. Apply the same- n extremum reduction of Lemma 22, reflecting its binary profile so $r \leq s$. If $r \leq 10$, $k \geq 80 \geq 8r$ and Lemma 30 applies. If $r \geq 11$, both multiplicities are at least eleven and the joint assertion applies. These cases exhaust all profiles. No reduction of dimension, extrapolation of a finite table or continuous-multiplicity assumption enters the lift. $\square$

8 Complete bottom-half coefficient certificates

We use $(n)_h = n(n-1) \cdots (n-h+1)$, with $(n)_0 = 1$. Every polynomial certificate in this section has nonnegative coefficients in nonnegative variables, so it proves an infinite-domain inequality. Exact determining grids check the polynomial identities; sample values alone never establish positivity.

8.1 Bottom orders thirteen through forty

Lemma 32. *For $13 \leq k \leq 40$, $n \geq 2k$, and every centered real vector of squared norm n ,*

$$|e_k|^2 \leq \binom{n}{k}.$$

Proof. By Lemma 22, use binary multiplicities $r \leq s$, $r + s = n$. Let $F_k(r, s)$ be the elementary coefficient of r copies of s and s copies of $-r$. These raw roots have squared norm nrs , so the desired inequality is exactly

$$F_k(r, s)^2 \leq \binom{n}{k} (rs)^k. \quad (41)$$

The following table specifies a base b_k for each order.

k	13	14	15	16	17	18	19	20	21	22
b_k	7	8	8	9	9	10	10	11	11	12
k	23	24	25	26	27	28	29	30	31	32
b_k	12	14	13	16	15	18	18	21	20	23
k	33	34	35	36	37	38	39	40		
b_k	22	26	25	29	28	32	31	35		

For $r \geq b_k$ put $r = b_k + u$, $s = r + x$ with $u, x \geq 0$. Form the integer power-sum polynomials $p_h = rs^h + s(-r)^h$ and the factorial-scaled Newton polynomials

$$E_0 = 1, \quad E_h = \sum_{a=1}^h (-1)^{a-1} \frac{(h-1)!}{(h-a)!} E_{h-a} p_a.$$

Thus $E_h = h!F_h$. The exact program constructs all twenty-eight polynomials

$$H_k(u, x) = k!(n)_k (rs)^k - E_k^2 \quad (42)$$

and checks every monomial coefficient is nonnegative. Hence $H_k \geq 0$ throughout the quadrant, which is (41) after division by $(k!)^2$.

Here and below the coefficient construction has an additional exact identity check. Since $p_1 = 0$, a Newton term for F_k contains at most $\lfloor k/2 \rfloor$ power-sum factors, so the total degree is at most $k + \lfloor k/2 \rfloor$. Consequently H_k has u -degree at most $3k$ and x -degree at most $2k$. On a complete rectangular grid of one more than each degree, the program compares it with

$$F_k(r, s) = \sum_a \binom{r}{a} \binom{s}{k-a} s^a (-r)^{k-a}. \quad (43)$$

These comparisons determine the full polynomial; its coefficient signs prove positivity on the infinite quadrant.

Multiplicities one and two are covered by Lemma 26. For each remaining $3 \leq r < b_k$, define

$$N_{r,k}(n) = \sum_{a=0}^r (-1)^a \binom{r}{a} r^{r-a} (k)_a n^a (n-a)_{r-a}. \quad (44)$$

The generating identity $(1 + sT)^r (1 - rT)^s = (1 - rT)^n [1 + nT/(1 - rT)]^r$ gives

$$F_k = (-r)^k \binom{n}{k} \frac{N_{r,k}(n)}{r^r (n)_r}.$$

Thus (41) is equivalent, after clearing positive denominators, to

$$G_{r,k}(n) = k! r^{2r} (n)_r^2 (n-r)^k - r^k (n)_k N_{r,k}(n)^2 \geq 0. \quad (45)$$

For $k = 13$ the four rows $r = 3, 4, 5, 6$ use $n = 13 + y$. For $14 \leq k \leq 40$ every row $3 \leq r < b_k$ uses $n = \max(2k, 2r) + y$. All $y \geq 0$ cover the required integer domains, and every cleared factor is positive there. The program checks every power coefficient in these 409 ray polynomials is nonnegative. It also checks the coefficient factorization and the gap identity against (43) at degree-plus-one distinct nonnegative integer nodes in each row. The cases $r \geq b_k$, $3 \leq r < b_k$ and $r = 1, 2$ exhaust all reflected profiles, proving the lemma. $\square$

8.2 Bottom orders forty-one through seventy-nine

Lemma 33. *For $41 \leq k \leq 79$, $n \geq 2k$, and every centered real vector of squared norm n , $|e_k|^2 \leq \binom{n}{k}$.*

Proof. Use the same binary reduction. Multiplicities one and two are already covered. If $3 \leq r \leq 10$ and $k \geq 8r$, apply Lemma 30. The remaining pairs are exactly

$$41 \leq k \leq 79, \quad \lfloor k/8 \rfloor + 1 \leq r \leq 10.$$

There are 115 such pairs. For each pair, construct (44) and (45) at $n = 2k + y$. Every power coefficient is nonnegative, as checked in full by the program, with the same complete univariate identity checks. This proves each entire ray $y \geq 0$.

For $r \geq 11$ the joint assertion of Theorem 31 applies as soon as

$$\frac{k(n-k)}{n} \geq 40, \quad \text{equivalently} \quad n \geq L_k := \left\lceil \frac{k^2}{k-40} \right\rceil.$$

The finite complement is exactly $2k \leq n < L_k$, $11 \leq r \leq \lfloor n/2 \rfloor$. To compute the integer coefficient, logarithmically differentiate $(1+sT)^r(1-rT)^s$. With $d = s - r$ this gives

$$(1+dT-rsT^2)F'(T) = -nrsTF(T),$$

and hence

$$F_0 = 1, \quad F_1 = 0, \quad (h+1)F_{h+1} = -dhF_h - rs(n-h+1)F_{h-1}. \quad (46)$$

The program checks (41) with exact integers at every profile using this recurrence. All 39 labels and 1,231,692 profiles pass. Each per-label count is also checked against $\sum_{n=2k}^{L_k-1} (\lfloor n/2 \rfloor - 10)$. The largest cutoff is $L_{41} = 1681$, so the finite interval ends at 1680, while its next endpoint is analytically covered. There is no extrapolation in n and no coefficient-reversal symmetry. $\square$

Corollary 34. *For every centered real vector of squared norm n and every $0 \leq k \leq \lfloor n/2 \rfloor$, $|e_k|^2 \leq \binom{n}{k}$.*

Proof. Lemmas 27 and 28 cover $k \leq 12$. Lemmas 32 and 33 cover the remaining $13 \leq k \leq 79$ in their exact bottom-half domains. If $k \geq 80$ and $n \geq 2k$, then $n - k \geq k \geq 80$, so the full-vector assertion of Theorem 31 applies. This partition covers every bottom order at the same n . $\square$

9 Top boundary bridges

9.1 Exact reciprocal normalization

For a reflected binary profile put $d = s - r \geq 0$. Its normalized coordinates are $\sqrt{s/r}$ repeated r times and $-\sqrt{r/s}$ repeated s times. Their product has squared modulus $(r/s)^d$ and their reciprocals are $r/\sqrt{rs}$ repeated r times and $-s/\sqrt{rs}$ repeated s times. Define integers T_h by

$$F(T) = (1+rT)^r(1-sT)^s = \sum_{h=0}^n T_h T^h.$$

Elementary reciprocal duality gives

$$\begin{aligned} |e_{n-j}|^2 &= \left(\frac{r}{s}\right)^d \frac{T_j^2}{(rs)^j}, \\ |e_{n-j}|^2 \leq \binom{n}{j} &\iff T_j^2 r^d \leq \binom{n}{j} (rs)^j s^d. \end{aligned} \quad (47)$$

In particular $d = 0$ causes no exceptional convention beyond $r^0 = s^0 = 1$. The differential identity

$$(1 - dT - rsT^2)F'(T) = -n(d + rsT)F(T)$$

yields the exact reverse recurrence

$$T_0 = 1, \quad T_1 = -nd, \quad (h+1)T_{h+1} = -d(n-h)T_h - rs(n-h+1)T_{h-1}. \quad (48)$$

These identities apply only after reduction to nonzero binary roots. The original vector may have zeros; Lemma 22 preserves its original n .

Lemma 35. *For $41 \leq j \leq 79$, $n \geq 2j$, and every centered real vector of squared norm n , $|e_{n-j}|^2 \leq \binom{n}{j}$.*

Proof. Set $k = n - j$ and

$$L_j = \max \left\{ \left\lceil \frac{j^2}{j-40} \right\rceil, j+80 \right\} = \left\lceil \frac{j^2}{j-40} \right\rceil.$$

The equality holds because $j^2/(j-40) - (j+80) = 40(80-j)/(j-40) > 0$. If $n \geq L_j$ and $r \geq 11$, then $j(n-j)/n \geq 40$ and the joint central theorem applies. If $r \leq 10$, then $k \geq 80 \geq 8r$ and the sparse lemma applies.

For $2j \leq n < L_j$ every $r \leq \lfloor (n-j)/8 \rfloor$ is also sparse. The program checks the exact integer inequality in (47) for every remaining

$$\lfloor (n-j)/8 \rfloor + 1 \leq r \leq \lfloor n/2 \rfloor.$$

All 39 labels and 1,006,861 profiles pass. Exact per-label counts are compared with $\sum_{n=2j}^{L_j-1} (\lfloor n/2 \rfloor - \lfloor (n-j)/8 \rfloor)$. Again $L_{41} = 1681$ is the maximum cutoff and its endpoint is analytically covered. The coefficient order $n-j \geq j \geq 41$ permits the same- n extremum lift, completing the full-real assertion. $\square$

Lemma 36. *For $51 \leq n \leq 599$, every centered real vector of squared norm n satisfies $|e_k|^2 \leq \binom{n}{k}$ at every order $0 \leq k \leq n$.*

Proof. For $k \leq n/2$ use Corollary 34. Otherwise put $j = n - k < n/2$. At $j \geq 80$ both $k, j \geq 80$, so Theorem 31 applies; at $41 \leq j \leq 79$ use Lemma 35. The only remaining labels are $0 \leq j \leq 40$ with $n \geq 2j$.

The program checks (47) for every one of these forty-one labels and every reflected multiplicity, in exactly the domain

$$0 \leq j \leq 40, \quad \max(51, 2j) \leq n \leq 599, \quad 1 \leq r \leq \lfloor n/2 \rfloor.$$

There is no pruning. The reverse recurrence shares its coefficient computation across all j at each (n, r) ; the zero-order coefficient is $T_0 = 1$. All 3,645,330 scalar comparisons pass across 89,075 shared profiles. For each j the exact count is checked against $\sum_{n=\max(51, 2j)}^{599} \lfloor n/2 \rfloor$. The coefficient order $k = n - j \geq \lfloor n/2 \rfloor \geq 26$ is in Lemma 22's range. Exact arithmetic and that lemma give the stated full-vector bounds. $\square$

10 An exact reciprocal strip in every degree at least six hundred

Theorem 37. *For $n \geq 600$, $1 \leq j \leq 19$, and every centered real vector of squared norm n ,*

$$|e_{n-j}|^2 \leq \binom{n}{j}.$$

Proof. Use Lemma 22 at order $k = n - j \geq 581$ and the binary notation of (47). Put

$$d = s - r, \quad u = r/s, \quad \eta = 1/n, \quad y = d^2/n, \quad a = 1 - y\eta = 4rs/n^2.$$

Define $A_h = h!2^h T_h/n^{3h/2}$. The reverse recurrence gives exactly

$$\begin{aligned} A_0 &= 1, & A_1 &= -2\sqrt{y}, \\ A_{h+1} &= -2\sqrt{y}(1 - h\eta)A_h - h(1 - y\eta)(1 - (h-1)\eta)A_{h-1}. \end{aligned} \tag{49}$$

There is no asymptotic error term. With

$$D_j(y, \eta) = (1 - y\eta)^j \prod_{i=0}^{j-1} (1 - i\eta),$$

the squared ratio in (47) is

$$\frac{|e_{n-j}|^2}{\binom{n}{j}} = \frac{u^d A_j^2}{j! D_j(y, \eta)}. \tag{50}$$

All denominator factors are positive. For $v = d/n \in [0, 1)$, $\log((1-v)/(1+v)) \leq -2v$, so $u^d \leq e^{-2y}$; at $d = 0$ equality holds with $u^0 = 1$. It therefore suffices to bound A_j^2 by $j! D_j e^{2y}$.

10.1 The compact finite-n polynomial certificate

Parity gives $A_h = y^{(h \bmod 2)/2} B_h(y, \eta)$, where $B_0 = 1, B_1 = -2$ and

$$B_{h+1} = -2y^{h \bmod 2} (1 - h\eta)B_h - h(1 - y\eta)(1 - (h-1)\eta)B_{h-1}. \tag{51}$$

Let $N_j = 4j + 40$ and $S_N(2y) = \sum_{h=0}^N (2y)^h / h!$. The integer polynomial

$$P_j = N_j! \left[j! D_j(y, \eta) S_{N_j}(2y) - y^{j \bmod 2} B_j(y, \eta)^2 \right] \tag{52}$$

has nonnegative complete tensor-product Bernstein arrays on each of

$$\left[0, \frac{25}{4}\right], \quad \left[\frac{25}{4}, \frac{25}{2}\right], \quad \left[\frac{25}{2}, 25\right], \quad [25, 50], \quad \eta \in [0, 1/600]. \tag{53}$$

All 19 polynomials and 76 rectangles pass. The degrees are at most $N_j + j$ and $2j - 1$ in y, η , respectively, with maxima 135 and 37. The program reconstructs every basis entry and accepts a rectangle only if all its entries are nonnegative. Throughout this enlarged continuous rectangle $1 - y\eta \geq 11/12$, so every denominator factor remains positive. As $e^{2y} \geq S_N(2y)$, the certificate proves the desired ratio bound for every $0 \leq y \leq 50$.

Here is the exact basis conversion and coverage argument. If M, Z are the two degrees, substitute $y = 50t, \eta = w/600$ and multiply P_j by 600^Z to get integer power coefficients c_{bc} . Set $l_y = \text{lcm}_b \left(\frac{M}{b} \right)$

and $l_\eta = \text{lcm}_c \binom{Z}{c}$. The tensor-product Bernstein coefficient at (i, h) , multiplied by the positive integer $l_y l_\eta$, is

$$\sum_{b \leq i, c \leq h} c_{bc} \binom{i}{b} \frac{l_y}{\binom{M}{b}} \binom{h}{c} \frac{l_\eta}{\binom{Z}{c}}. \quad (54)$$

This follows by applying $t^b = \sum_{i=b}^M \binom{i}{b} \binom{M}{b}^{-1} \binom{M}{i} t^i (1-t)^{M-i}$ in both variables. All basis functions are nonnegative. Exact midpoint de Casteljau subdivision uses sums and a common positive denominator; common positive gcd factors can be removed. Both children are retained, and an already-positive interval can be refined further to obtain the common partition (53). The code explicitly checks those four intervals, the full η interval and total area $1/12$. Thus the finite computation covers the full rectangle, not a set of sample points. Optional output identifies each reconstructed full array by its stream hash; its sign proof is the checked entries themselves, not the hash.

10.2 An analytic tail with translated polynomial certificates

If $r \leq k/8$, Lemma 30 already proves the result at the original order $k = n - j$. Otherwise

$$\frac{d}{n} < \frac{3 + j/n}{4} \leq \frac{1819}{2400} < \frac{19}{25}.$$

Consequently $a > 264/625 > 2/5$, and the elementary product inequality gives

$$\prod_{i=0}^{j-1} (1 - i/n) \geq 1 - \frac{j(j-1)}{2n} \geq \frac{143}{200} > \frac{7}{10}.$$

Let $C_0(X) = 1$, $C_1(X) = X$ and $C_{h+1}(X) = XC_h(X) + hC_{h-1}(X)$. These are the absolute-coefficient Hermite polynomials, explicitly

$$C_j(X) = \sum_{l=0}^{\lfloor j/2 \rfloor} \frac{j! X^{j-2l}}{2^l l! (j-2l)!}.$$

In (49) every coefficient factor apart from $2\sqrt{y}$ and h lies in $[0, 1]$ on the actual domain. Induction yields $|A_j| \leq C_j(2\sqrt{y})$. Thus (50) is at most

$$\frac{10}{7} \left(\frac{5}{2}\right)^j e^{-2y} \frac{C_j(2\sqrt{y})^2}{j!}. \quad (55)$$

Parity makes $C_j(2\sqrt{y})^2$ an integer polynomial in y . Define

$$Q_j(y) = S_{N_j}(2y) - \frac{10}{7} \left(\frac{5}{2}\right)^j \frac{C_j(2\sqrt{y})^2}{j!}.$$

After multiplication by $7 \cdot 2^j j! N_j!$, the program checks that every power coefficient of $Q_j(50 + t)$ is nonnegative, for every $1 \leq j \leq 19$. More explicitly the unshifted coefficient of y^h is

$$7 \cdot 2^j j! 2^h \frac{N_j!}{h!} - 10 \cdot 5^j N_j! [y^h] C_j(2\sqrt{y})^2,$$

with the second term zero above degree j . Translation uses $[t^b]P(50 + t) = \sum_{h \geq b} [y^h]P(y) \binom{h}{b} 50^{h-b}$. All 19 complete translated arrays pass. Hence $Q_j(y) \geq 0$ for every $y \geq 50$, and $e^{2y} \geq S_N(2y)$ makes (55) at most one on that entire half-line.

The sparse cases and the compact and tail regions exhaust all binary profiles, including balance at $y = 0$ and the shared boundary $y = 50$. Lemma 22 completes the full-real conclusion without reciprocal operations on original zero coordinates. The proof controls finite n exactly and does not use a limiting Hermite assertion. $\square$

11 A global affine contour budget for top codimensions nineteen through forty

Theorem 38. *For $n \geq 600$, $19 \leq j \leq 40$, and every centered real vector of squared norm n , $|e_{n-j}|^2 \leq \binom{n}{j}$.*

Proof. Use the binary Cauchy modulus f , entropy factor H_0 and area A in the proof of Theorem 24; put $k = n - j$, $v = k(n - k)/n$. The interval rearrangement and layer-cake argument in Lemma 29 gives, without an area restriction,

$$I := \frac{1}{\pi} \int_{-1}^1 \frac{f(t)}{\sqrt{1-t^2}} dt \leq \int_0^1 \Phi(\ell(h)) dh, \quad \Phi(a) = \frac{\arccos(1-a)}{\pi}, \quad \int_0^1 \ell(h) dh = A.$$

The tangent at $a_0 = 3/10$ is $T(a) = \alpha + \beta a$, where

$$\beta = \frac{1}{\pi\sqrt{51/100}}, \quad \alpha = \frac{\arccos(7/10)}{\pi} - \frac{3}{10}\beta.$$

It is a global majorant of Φ on $[0, 2]$. Indeed Φ is concave on $[0, 1]$, while $T - \Phi$ is concave on $[1, 2]$, whose minimum is at an endpoint. Endpoint one is covered by the tangent inequality, and at two

$$\pi T(2) = \arccos(7/10) + \sqrt{17/3} > \frac{77}{100} + \frac{119}{50} = \frac{63}{20} > \frac{22}{7} > \pi.$$

The first bound follows from $\cos(77/100) \geq 1 - (77/100)^2/2 > 7/10$ and $(119/50)^2 < 17/3$. Therefore

$$I \leq \alpha + \beta A \quad (0 \leq A \leq 2). \quad (56)$$

Both coefficients are positive. The following conservative rational upper bounds are certified exactly:

$$\alpha < a := \frac{11947}{100000}, \quad \beta < b_0 := \frac{44573}{100000}, \quad \sqrt{\pi} < p_0 := \frac{177246}{100000}, \quad \sqrt{2\pi} < h_0 := \frac{250663}{100000}.$$

For completeness, Machin's identity

$$\pi = 16 \arctan(1/5) - 4 \arctan(1/239)$$

follows from $\tan(4 \arctan(1/5)) = 120/119$ and the tangent subtraction formula on the positive principal branch. Alternating partial sums with eight/nine terms at $1/5$ and three/four terms at $1/239$ give

$$\frac{314159}{100000} < \pi < \frac{3141593}{1000000}.$$

Also $71414/100000 < \sqrt{51/100} < 71415/100000$, and the alternating degree-eight Taylor upper bound for $\cos(7954/10000)$ is below $7/10$, so $\arccos(7/10) < 7954/10000$. These bounds give the upper β by lower bounding both denominator factors, and the upper α by upper bounding its

positive numerator and lower bounding π . The numerator is positive, for instance from $\arccos(.7) > .77$ and $.3/\sqrt{.51} < .3/.7$. Squaring checks the two radical upper bounds. The program verifies all these rational comparisons; no decimal transcendental approximation decides acceptance.

If $r \leq k/8$, use Lemma 30. Otherwise

$$r > k/8 \geq (n - 40)/8 \geq 7n/60, \quad rs > 371n^2/3600.$$

The Beta and Stirling bounds proved earlier are

$$A \leq \sqrt{\pi/v} \frac{n}{n+2} e^{E_a}, \quad \frac{H_0^2}{\binom{n}{k}} \leq \sqrt{2\pi v} e^{E_b}, \quad E_a = \frac{n}{6rs}, \quad E_b = \frac{1}{12v}.$$

Here $E_a < 1/371$, while $11039/600 \leq v \leq 40$ gives $E_b \leq 50/11039$. The lower endpoint follows because $j(1 - j/600)$ increases on $19 \leq j \leq 40$. Since $e^t \leq (1 - t)^{-1}$ for $0 \leq t < 1$, $e^{E_a} \leq 371/370$ and $e^{E_b} \leq 11039/10989$.

Drop the favorable $n/(n+2)$ and put

$$b = b_0 p_0 \frac{371}{370}, \quad C = h_0 \frac{11039}{10989}.$$

Equations (56) and the coefficient contour bound yield

$$\frac{|e_k|^2}{\binom{n}{k}} \leq C \left(a^2 t + 2ab + \frac{b^2}{t} \right), \quad t = \sqrt{v}.$$

The right side is convex in $t > 0$, as its second derivative is $2Cb^2/t^3 > 0$. At the two endpoints use

$$\frac{42893}{10000} < \sqrt{11039/600} < \frac{42894}{10000}, \quad \frac{63245}{10000} < \sqrt{40} < \frac{63246}{10000}.$$

The positive linear term uses the upper enclosure and the reciprocal term the lower. Both resulting rational upper bounds are strictly below one:

$$C \left(a^2 \frac{42894}{10000} + 2ab + b^2 \frac{10000}{42893} \right) < 1, \\ C \left(a^2 \frac{63246}{10000} + 2ab + b^2 \frac{10000}{63245} \right) < 1.$$

These are exact fraction comparisons (approximately .99918 and .95378). Convexity proves the entire nonsparse binary domain; the sparse lemma and same- n extrema finish the theorem. $\square$

12 All scalar orders and the covariance certificate

Proof of Theorem 1. First normalize a positive squared norm to $\sum z_i^2 = n$; the centered normalized slice is empty at $n = 1$. For $2 \leq n \leq 50$, use Lemma 25. For $51 \leq n \leq 599$, use Lemma 36. For $n \geq 600$, every $k \leq \lfloor n/2 \rfloor$ follows from Corollary 34. In the top half put $j = n - k < n/2$. At $j = 0$, product AM-GM gives $|e_n|^2 \leq 1$. At $1 \leq j \leq 19$, use Theorem 37; at $19 \leq j \leq 40$, use Theorem 38. The overlap is harmless. At $41 \leq j \leq 79$, Lemma 35 applies since $n \geq 2j$. At $j \geq 80$, both $j, k \geq 80$ and Theorem 31 applies. This exhausts every coefficient and degree at the same n , including the $n = 50/51$ and $599/600$ endpoints and equality $k = n/2$.

Homogeneity of e_k now gives (2) for any positive squared norm. At zero norm every coordinate vanishes, so all $e_k = 0$ for $k \geq 1$, while $e_0 = 1$ separately; no 0^0 convention is needed. For even n , the vector with $n/2$ copies of one and $n/2$ copies of minus one has squared norm n and $|e_n|^2 = 1 = \binom{n}{n/2}$. Thus the universal scalar coefficient one is optimal. Original zeros, repeated coordinates and all balanced or unbalanced profiles are included by the same- n reductions. $\square$

13 Equality and endpoint stability of the scalar bound

Theorem 39. *On the normalized real slice $\sum_i z_i = 0$, $\sum_i z_i^2 = n$ with $n \geq 2$, equality in $|e_k(z)|^2 \leq \binom{n}{k}$ holds exactly at $k = 0$, or at $k = n$ with even n and balanced entries $+1, -1$ up to permutation. Every interior order $1 \leq k < n$ is strict.*

Proof. For each fixed $n, k \geq 3$, Lemma 22 supplies some binary maximizer at that SAME n . It therefore suffices to prove strictness for every binary interior profile; it is not necessary that every maximizer itself be binary. Orders one and two are immediate from $e_1 = 0$ and the ratio $n/[2(n-1)] < 1$ for $n > 2$. At $k = n$, product AM–GM on z_i^2 gives equality only when all $|z_i| = 1$; centering forces even n and balanced signs.

Here are complete strictness gates on the proof’s infinite domains. Multiplicity one has ratio at most $2/3$ for $k \geq 3$. Multiplicity two has strict cubic and quartic bounds except the balanced $n = k = 4$ endpoint; its seven cleared polynomials have strictly positive coefficients, and its factorial tail is strict. The general cubic bound is strict, and the quartic bound is strict for $n > 4$.

For each fixed order $5 \leq k \leq 12$, the accepted quadrant gap arrays have nonnegative coefficients, with strictly positive coefficients at both u and x . At odd k the constant is positive. At even k the only zero on the nonnegative quadrant is therefore $u = x = 0$, namely $n = k$, $r = s = k/2$. Every smaller-multiplicity exception ray has positive constant. All 28 bottom13–40 quadrant gaps, all 409 corresponding rays and all 115 bottom41–79 rays have positive constant and nonnegative coefficients. Their cleared denominators are positive. Thus these entire quadrants and rays are strict. The central and sparse bounds have ratios strictly below one, as do both global-tangent endpoint guards.

For the reciprocal strip, if $d = s - r > 0$, then $y > 0$ and $e^{2y} > S_N(2y)$ strictly. The compact and translated-tail certificates, with their positive denominator, therefore give strict ratio below one even if a polynomial gap vanishes. At $d = 0$, the normalized generating polynomial is $(1 - T^2)^{n/2}$. Odd coefficients vanish. For $k = 2l$ interior, Vandermonde gives

$$\binom{n}{2l} = \sum_a \binom{n/2}{a} \binom{n/2}{2l-a} > \binom{n/2}{l}^2,$$

because besides $a = l$, the term $a = l - 1$ is positive for $1 \leq l \leq n/2 - 1$.

The remaining domains are finite and explicit. The companion verifier checks all 40,425 small scalar labels, 1,231,692 bottom bridge profiles, 1,006,861 top bridge profiles and 3,645,330 finite endpoint labels for STRICT inequality, permitting equality only at original order $k = n$ with $r = s$. It also checks all 572 coefficient strictness/zero-set gates just described on the reconstructed complete arrays. These are the full finite complements of the analytic partition, not a cutoff extrapolation. The partition in the proof of Theorem 1 now proves binary strictness at every interior order; Lemma 22 completes the full-real assertion. $\square$

The classification is a strictness completion of Theorem 1, rather than a new extremal method. Its elementary endpoint consequences are recorded for clarity. Put $\varrho = |e_n(z)|^2 > 0$. The classical logarithmic AM–GM inequality gives

$$\sum_i (|z_i| - 1)^2 \leq -\log \varrho. \tag{57}$$

Indeed $\log u \leq 2(\sqrt{u} - 1)$ and $\sum_i z_i^2 = n$ imply $\log \varrho \leq 2 \sum_i (|z_i| - 1) = -\sum_i (|z_i| - 1)^2$. The square-root variance refinement of AM–GM in Aldaz [19, Theorem 0.1] also gives the stronger intermediate

formula

$$\varrho^{1/n} \leq \left(1 - \frac{1}{2n} \sum_i (|z_i| - 1)^2\right)^2.$$

Thus no new general quantitative AM–GM principle is claimed. If $\varrho \geq 1 - \epsilon > 0$, let $\sigma_i = \text{sign}(z_i)$. Centering and Cauchy yield

$$|\sum_i \sigma_i| \leq \sqrt{n \sum_i (|z_i| - 1)^2}.$$

If $-\log(1 - \epsilon) < 1/n$, the integer sign imbalance is less than one and hence zero. Then n is even, σ is an exact balanced equality vector, and $\|z - \sigma\|_2^2 \leq -\log(1 - \epsilon)$. This is endpoint stability only; no uniform interior-order gap is asserted.

For odd $n \geq 3$, classical two-moment product optimization gives the exact endpoint maximum

$$\max |e_n(z)|^2 = \frac{n-1}{n+1}. \quad (58)$$

For completeness, a maximizer has nonzero product since a binary competitor does. Lagrange multipliers for $\sum \log z_i^2$ are valid: the constraint gradients are independent. The equation $2/z_i = \alpha + 2\beta z_i$, summed after multiplying by z_i , gives $\beta = 1$. All coordinates therefore take at most two values, necessarily of opposite sign. Their reflected multiplicities $r \leq s$ have squared product $(r/s)^{s-r}$. Writing $d = s - r$, its logarithm $d \log((n-d)/(n+d))$ strictly decreases for $d > 0$. Odd n makes the smallest positive d equal one, proving (58), with exactly the nearest-balanced two-value profiles up to sign and permutation. This elementary endpoint optimization is not claimed as a new phenomenon.

14 Completion of the count-law certificate

Proof of Theorem 2. For $n \geq 2$, scaling Theorem 1 with (5) gives

$$|R_k(y)|^2 \leq \binom{n}{k} (2/n)^k (y^\top B y)^k.$$

The quotient tensor transfer therefore yields $\mathbb{E}_M U_k^2 \leq (2W)^k$. Orthogonality proves (3)’s chi-square bound for every $W \geq 0$; the sum is finite, so no geometric convergence hypothesis is required. At $2W = 1$ it is $n - 1$, and for $2W < 1$ it is at most $(2W)^2/(1 - 2W)$.

For the TV conclusion let $x = 2W \in (0, 1)$ and $\delta = \text{TV}(P, M)$. Then $v := \chi^2(P \| M) \leq \sum_{k=2}^n x^k \leq x^2/(1 - x)$. If $x \leq 1/2$, Cauchy gives $\delta \leq \sqrt{v}/2 \leq x$. If $x \geq 1/2$ and $\delta \leq 1/2$ the result is immediate; otherwise $\delta > x$ contradicts Lemma 6, since $v \geq \delta/(1 - \delta) > x/(1 - x) \geq v$. At $W = 0$ use rigidity and at $W \geq 1/2$ use $\text{TV} \leq 1$. Degree one is exactly multinomial, giving an empty chi-square sum; one retained category is deterministic, and mean-zero categories vanish almost surely and can be deleted.

For optimality, take even n and $p(x, y) = x^{n/2} y^{n/2}$, the classical balanced deterministic/binomial example [9, Remark 2.2(a)]. It has $W = 1/2$ and $\text{TV}(P, \text{Mult}(n, 1/2)) = 1 - 2^{-n} \binom{n}{n/2} \rightarrow 1$. Thus no universal linear coefficient below two is possible. $\square$

14.1 The equivalent own-mean Krawtchouk bound

In the conventional normalization [16, equation 18.23.3 and Table 18.19.1],

$$(1 - [(1 - p)/p]t)^x (1 + t)^{n-x} = \sum_{k=0}^n \binom{n}{k} K_k(x; p, n) t^k.$$

The squared norm under the binomial probability weight $w_p(x) = \binom{n}{x} p^x (1-p)^{n-x}$ is $[(1-p)/p]^k / \binom{n}{k}$. Thus the binomial-orthonormal polynomial is

$$\hat{K}_k(x; p, n) = \sqrt{\binom{n}{k}} [p/(1-p)]^{k/2} K_k(x; p, n).$$

Put $s = np$, $r = n - s$ with positive integer r, s , and substitute $t = sT$ in the generating function. The raw coefficient of $(1 + sT)^r (1 - rT)^s$ is $s^k \binom{n}{k} K_k(s; s/n, n)$. After division by $(rs)^{k/2} \sqrt{\binom{n}{k}}$, this yields

$$\frac{e_k}{\sqrt{\binom{n}{k}}} = \hat{K}_k(np; p, n).$$

Consequently Theorem 1 is equivalent, through the same- n two-value reduction and elementary low orders, to

$$|\hat{K}_k(np; p, n)| \leq 1 \quad (0 < p < 1, np \in \mathbb{Z}, 0 \leq k \leq n).$$

The integer-mean condition is substantive; no continuous-bias relaxation at a noninteger evaluation is claimed. This normalization is distinct from the counting-measure orthonormal function $\sqrt{w_p(x)} \hat{K}_k(x)$. Entrywise orthogonal-matrix bounds for that function only give $|\hat{K}_k(x)| \leq 1/\sqrt{w_p(x)}$, which does not imply the own-mean bound one. The generating function and norm are classical inputs, not new constructions.

14.2 Reproducibility

Run `verify_spectral.py` with its shipped helpers `verify_scalar_one.py`, `verify_sharp_two.py` and `verify_refined.py`; the option `--output`, followed by a filename, saves exact certificate data. The program checks the 40,425 small scalar comparisons, all multiplicity-two and fixed-order certificate arrays, the complete bottom and top boundary domains, all 3,645,330 finite endpoint comparisons through 599, the 19 reciprocal-strip polynomials with 76 full compact basis reconstructions and 19 translated tails, and every global-tangent rational guard. Every accepted compact basis entry is regenerated and sign-checked. Saved array hashes identify those computations and do not replace their signs. Infinite polynomial quadrants/rays and finite exhaustive domains have distinct, explicit coverage arguments. All acceptance gates use exact integers/rationals and remain active under Python optimization. The entry point also verifies all 572 equality strictness gates, the complete finite strictness domains and exact small-law spectral/example identities. These finite checks do not certify the Gaussian random-tensor existence proof, the analytic cone argument or any limit theorem; those are manuscript proofs.

15 Gaussian mutual-TV limits along powers of a fixed law

We now fix one law P throughout, including its dimension, support and probabilities. The following result concerns mutual TV of two discrete convolution laws; it does not assert ordinary TV convergence of a lattice law to a continuous Gaussian.

Theorem 40. *Let $C = \text{Cov}(P)$, $S = \text{diag}(t) - tt^\top/n$, and $H = \{x : \mathbf{1}^\top x = 0\}$. For $P_N = P^{*N}$ and $M_N = M^{*N} = \text{Mult}(Nn, t/n)$,*

$$\lim_{N \rightarrow \infty} \text{TV}(P_N, M_N) = \text{TV}(\mathcal{N}_H(0, C), \mathcal{N}_H(0, S)). \quad (59)$$

Degenerate C is allowed. If $0 < W < 1/2$, the right side is strictly below $2W$. Consequently every fixed P with $0 < W < 1/2$ has a threshold $N_0(P)$ such that $\text{TV}(P_N, M_N) < 2W$ for every integer $N \geq N_0(P)$. At $W = 0$ all powers are exactly multinomial; at $W \geq 1/2$ the inequality holds at every power.

The support input is classical: Brändén's jump-system theorem [11, Theorem 3.2] extends earlier homogeneous support results of Choe, Oxley, Sokal and Wagner [12]. The finite-support Fourier local limit used below is also classical; see the background and lattice-period discussion in Randles [13]. We give the needed argument, including the 11 passage and singular case, to expose the exact hypotheses. No new local-limit or Gaussian-comparison theorem is claimed.

Proof of Theorem 40. **15.1 Covariance coordinates and support lattice**

For p^N , the mean, covariance, S and B are multiplied by N , so $W_N = W$. Products preserve stability, homogeneity and normalization. If $m = 1$ all laws are identical point masses. Suppose $m \geq 2$, put $d = m - 1$, and delete the final coordinate. The omitted count is determined by the total, so this is a bijection on each support and preserves TV. Denote the principal covariance blocks by C', S', B' , and the common projected block mean by $\mu = (t_1, \dots, t_d)$.

The matrix S' is positive definite and

$$(S')^{-1} = \text{diag}(1/t_1, \dots, 1/t_d) + \frac{1}{t_m} \mathbf{1}\mathbf{1}^\top.$$

Since $B\mathbf{1} = 0$, $\sum_{i,j \leq d} B_{ij} = B_{mm}$, hence

$$\text{Tr}((S')^{-1}B') = W. \tag{60}$$

When $0 < W < 1/2$, the matrix $D = 2(S')^{-1/2}B'(S')^{-1/2}$ is positive semidefinite with trace $2W < 1$. Thus every eigenvalue is below one and $C' = (S')^{1/2}(I - D)(S')^{1/2}$ is positive definite. In particular C is then nondegenerate on H . More generally, C is positive definite on H exactly when C' is positive definite, since centered count vectors are obtained from their first d coordinates by the linear isomorphism $x' \mapsto (x', -\mathbf{1}^\top x')$ onto H .

Assume first that C is positive definite on H . The support $\mathcal{A}$ of p obeys unit exchange: if $a, b \in \mathcal{A}$ and $a_i > b_i$, some j with $a_j < b_j$ has $a - e_i + e_j \in \mathcal{A}$. Indeed the jump-system two-step axiom applies to the step $-e_i$ toward b ; its first step leaves the fixed-total support, and homogeneity forces the second step to be $+e_j$ toward a deficient coordinate. Repeated exchange decreases the distance to b by two and connects every pair of support points.

Make a graph on categories by joining i, j whenever two support points differ by $e_i - e_j$. If it were disconnected, each graph-component total would be constant throughout the support: every exchange path stays within its components. Subtracting a multiple of $\mathbf{1}$ from a proper component indicator would give a nonzero vector in H with zero variance, contradicting positive definiteness of C there. The graph is therefore connected. Its edge differences generate the full root lattice $\{z \in \mathbb{Z}^m : \mathbf{1}^\top z = 0\}$, and all support differences already lie in this lattice. Consequently the projected support-difference group is exactly $\mathbb{Z}^d$.

For a characteristic function $\phi(\theta)$, modulus one occurs precisely when all support phases agree. The full difference group therefore implies $|\phi(\theta)| < 1$ away from the origin of the Fourier torus. The projected multinomial block law has the same aperiodicity: its support contains zero and each unit vector, because all categorical probabilities are positive. Thus both block laws have positive covariance, full difference group and the same mean μ . Positive covariance alone would not exclude lattice periods for an arbitrary discrete law; the stable-support argument is essential here.

15.2 Uniform local limit and its l1 consequence

Let X be a fixed finite-support $\mathbb{Z}^d$ law with mean μ , positive definite covariance Σ and full support-difference group. Write $f_N(k) = \Pr(X_1 + \dots + X_N = k)$ and g_Σ for the centered Gaussian density. Its centered characteristic function satisfies

$$\psi(\theta) = e^{-i\mu \cdot \theta} \phi(\theta) = 1 - \frac{1}{2} \theta^\top \Sigma \theta + O(\|\theta\|^3).$$

Positive covariance gives $|\psi(\theta)| \leq e^{-c\|\theta\|^2}$ near zero for some fixed $c > 0$. On the compact complement of that neighborhood in the torus, aperiodicity gives $|\phi| \leq r < 1$.

Fourier inversion and $\theta = u/\sqrt{N}$ show

$$\sup_{k \in \mathbb{Z}^d} \left| N^{d/2} f_N(k) - g_\Sigma \left(\frac{k - N\mu}{\sqrt{N}} \right) \right| \rightarrow 0. \quad (61)$$

Here is the uniform error argument. On bounded u -sets, $\psi(u/\sqrt{N})^N$ converges uniformly to $e^{-u^\top \Sigma u/2}$. On the expanding small- θ region it is dominated by $e^{-c\|u\|^2}$, so dominated convergence controls the integral error there. The complementary scaled Fourier integral is bounded by a constant times $N^{d/2} r^N$, and the Gaussian integral outside the expanding region also tends to zero. The phase $e^{-i(k - N\mu) \cdot u/\sqrt{N}}$ has modulus one, making the error estimate uniform in k . This proves (61) with the usual Gaussian Fourier normalization.

It follows that

$$\sum_{k \in \mathbb{Z}^d} \left| f_N(k) - N^{-d/2} g_\Sigma \left(\frac{k - N\mu}{\sqrt{N}} \right) \right| \rightarrow 0. \quad (62)$$

For fixed R , the box $\|k - N\mu\|_\infty \leq R\sqrt{N}$ has $O_R(N^{d/2})$ sites, so (61) sums to $o_R(1)$ there. Outside, Chebyshev bounds the probability mass by $\text{Tr } \Sigma / R^2$. Shifted-grid Gaussian sums converge to the corresponding integrals: on compact sets this follows from uniform continuity, and Gaussian decay controls the tails uniformly in the grid shift. Thus their tail also vanishes after first taking $N \rightarrow \infty$ and then $R \rightarrow \infty$. This proves (62); no discrete-to-continuous TV assertion is involved.

Apply this to the projected P and M blocks. Their two l1 errors imply

$$\text{TV}(P_N, M_N) - \frac{1}{2} N^{-d/2} \sum_{k \in \mathbb{Z}^d} \left| g_{C'} \left(\frac{k - N\mu}{\sqrt{N}} \right) - g_{S'} \left(\frac{k - N\mu}{\sqrt{N}} \right) \right| \rightarrow 0.$$

Another shifted-grid Riemann sum gives the Gaussian TV on the right side of (59). The coordinate isomorphism onto H preserves TV, proving the claimed coordinate-independent limit when C is nonsingular there.

15.3 Singular covariances and the strict Gaussian gap

If C is singular on H , choose nonzero $v \in H$ with $v^\top C v = 0$. Then $v \cdot (A - t) = 0$ almost surely. Thus P_N is supported on $E_N = \{a : v \cdot (a - Nt) = 0\}$. Under M_N , the scalar projection $v \cdot (A - Nt)/\sqrt{N}$ converges by the ordinary scalar CLT to a nondegenerate normal, since $v^\top S v > 0$. For every $\eta > 0$, Portmanteau on the closed interval $[-\eta, \eta]$ gives

$$\limsup_N M_N(E_N) \leq \Pr(|\mathcal{N}(0, v^\top S v)| \leq \eta).$$

Letting $\eta \downarrow 0$ shows $M_N(E_N) \rightarrow 0$. Hence $\text{TV}(P_N, M_N) \geq 1 - M_N(E_N) \rightarrow 1$. The degenerate Gaussian with covariance C is supported on the corresponding hyperplane, to which the Gaussian with covariance S assigns zero mass; their TV is one. This proves (59) also in the singular case.

Finally suppose $0 < W < 1/2$, so the projected covariances are positive definite. By (60), $D \succeq 0$ and $\text{Tr } D = 2W < 1$. For the two centered Gaussian densities,

$$\frac{g_{S'}(x)}{g_{C'}(x)} = \sqrt{\det(I - D)} e^{x^\top ((C')^{-1} - (S')^{-1})x/2} \geq \sqrt{\det(I - D)}.$$

The quadratic exponent is nonnegative by covariance ordering. Integrating the minimum of the densities bounds their TV by $1 - \sqrt{\det(I - D)}$. If λ_j are the eigenvalues of D , then

$$\det(I - D) = \prod_j (1 - \lambda_j) \geq 1 - \sum_j \lambda_j = 1 - 2W.$$

This elementary product inequality follows by induction. Since $\sqrt{a} > a$ for $0 < a < 1$, the Gaussian TV is strictly below $2W$. Convergence in (59) leaves a fixed positive gap, yielding $N_0(P)$ for all subsequent integers. Rigidity handles $W = 0$, while $W \geq 1/2$ is trivial. This completes the proof. $\square$

The coefficient two in this eventual statement is optimal uniformly over fixed base laws. For $p(x, y) = xy$, one has $W = 1/2$ and

$$\text{TV}(P_N, M_N) = 1 - \binom{2N}{N} / 4^N \rightarrow 1.$$

Theorem 2 already bounds every power, because $W_N = W$. The limit theorem additionally identifies the limiting value and its strict Gaussian gap; it provides no uniform threshold or rate. Total variation under powers need not be monotone: $(x/4 + 3y/4)(3x/4 + y/4)$ has TV $1/8$ from its degree-two reference, while its square has TV $11/128$; xy has TV $1/2$, while its square has TV $5/8$. In each example W is unchanged under powers. These exact examples prevent pulling the limiting bound back by an assumed monotonicity.

16 Scope and remaining questions

The optimal real scalar coefficient is one, and the optimal universal linear TV coefficient is two. The covariance-spectrum refinement supplies a quadratic likelihood approximation, diffuse-spectrum convergence and a cubic obstruction to a stronger product-class bound. The sharp complex (6, 5) theorem gives the optimal spectral factor $3/4$ at degree six in every covariance rank; together with the lower and top orders it closes the factor-one target throughout that degree. The exact rank-two realization and the stronger factor $1296/3125$ for structured pencils and matching-mean degree-four times degree-two stable products are retained. For higher degrees the order-five bounds include factor 2, factor $10/9$ for every $n \geq 256$, two factor-one spectral regions and the improved derivative branch. The unrestricted higher-interior target in degrees $n \geq 7$ remains open. The credited complex transfer gives factor 24 in every order and an inverse-determinant chi-square envelope when $\rho(D) < 1$. The all-degree real scalar proof uses a complete finite/infinite domain partition, while the new complex theorem concerns a fixed dimension and order. Neither a general complex coefficient-one theorem nor all-degree spectral closure is asserted. The fixed-law Gaussian limit is a separate asymptotic conclusion.

The mean vector and covariance are treated as known inputs. We prove no sample-complexity result for estimating W , practical testing guarantee, sampler construction or computational speedup. Stability is a substantive hypothesis; matching a few moments without it does not certify the full law. Singular covariance deficit, repeated roots and zero coefficients are included. The positivity lemma is applied only to the complete density P/M , since individual centered-expansion terms are signed.

Computational and AI assistance

The computer-assisted proof is reproducible from the five supplied Python files using only the standard library. The main verifier reconstructs all exact arrays and exhaustive finite domains described above; no private module or external numerical package is required. Seeded rational diagnostic fixtures are checked exactly and are not used as proof of an untested domain. The sharp complex-six bound additionally uses the twenty exact rational Bernstein interval certificates reconstructed by `verify_complex_six.py`; its weighted-median and affine-root reductions and the stable-law transfer are supplied as proofs. The other fifth-order scalar and spectral comparisons and the positive-polynomial high-degree certificate are analytic proofs. The support-exchange, Fourier local-limit, singular-covariance and Gaussian arguments are also analytic proofs; no numerical calculation is claimed to verify a limit theorem or its unspecified threshold. Generative AI systems assisted with mathematical exploration, literature discovery, proof drafting, exact-code development and independent adversarial proof checks. These are internal AI-assisted checks, not external human refereeing. The author retains responsibility for the manuscript. Classical inputs and the close Han–Niles-Weed methodology are explicitly attributed; the proofs and complete computational domains are provided for independent assessment.

References

- [1] S. Banach, Über homogene Polynome in L^2 , *Studia Mathematica* **7** (1938), 36–44. doi:10.4064/sm-7-1-36-44.
- [2] D. Carando and J. T. Rodríguez, Symmetric multilinear forms on Hilbert spaces: where do they attain their norm? (2018). arXiv:1810.09373.
- [3] W. Ehm, Binomial approximation to the Poisson binomial distribution, *Statistics & Probability Letters* **11**(1) (1991), 7–16. doi:10.1016/0167-7152(91)90170-V.
- [4] A. Guntuboyina, S. Saha and G. Schiebinger, Sharp inequalities for f -divergences (2013). arXiv:1302.0336.
- [5] Y. Han and J. Niles-Weed, Approximate independence of permutation mixtures, *The Annals of Statistics* **54**(3) (2026), 1594–1621. doi:10.1214/26-AOS2619. arXiv:2408.09341; the coefficient result used here is Theorem 4.3(2) in version 3.
- [6] Y. Han and J. Niles-Weed, Elementary symmetric polynomial inequalities for centered vectors and matrices (2026). arXiv:2607.23836.
- [7] W.-L. Loh, Stein’s method and multinomial approximation, *The Annals of Applied Probability* **2**(3) (1992), 536–554. doi:10.1214/aoap/1177005648.
- [8] C. Riener and R. Schabert, Linear slices of hyperbolic polynomials and positivity of symmetric polynomial functions (2023). arXiv:2203.08727, version 2.
- [9] B. Roos, Closeness of convolutions of probability measures, *Bernoulli* **16**(1) (2010), 23–50. doi:10.3150/08-BEJ171. arXiv:0811.0622.
- [10] B. Roos, On Bobkov’s approximate de Finetti representation via approximation of permanents of complex rectangular matrices (2014). arXiv:1203.2871, version 2.
- [11] P. Brändén, Polynomials with the half-plane property and matroid theory, *Advances in Mathematics* **216** (2007), 302–320. arXiv:math/0605678, version 4.

- [12] Y.-B. Choe, J. G. Oxley, A. D. Sokal and D. G. Wagner, Homogeneous multivariate polynomials with the half-plane property, *Advances in Applied Mathematics* **32** (2004), 88–187. doi:10.1016/S0196-8858(03)00078-2. arXiv:math/0202034.
- [13] E. Randles, Local limit theorems for complex functions on $\mathbb{Z}^d$, *Journal of Mathematical Analysis and Applications* **519** (2023), 126832. doi:10.1016/j.jmaa.2022.126832. arXiv:2201.01319, version 2.
- [14] H. Derksen, P. Ivanov, C. H. Lee and E. Viola, Pseudorandomness, Symmetry, Smoothing: I, in *39th Computational Complexity Conference*, LIPIcs **300** (2024), 18:1–18:27. doi:10.4230/LIPIcs.CCC.2024.18.
- [15] J.-T. Hsieh, D. M. Kane, P. K. Kothari, J. Li, S. Mohanty and S. Tiegel, Rigorous Implications of the Low-Degree Heuristic (2026). arXiv:2601.05850.
- [16] National Institute of Standards and Technology, *Digital Library of Mathematical Functions*, §18.19 and §18.23, Table 18.19.1 and equation 18.23.3.
- [17] B. Roos, Multinomial and Krawtchouk approximations to the generalized multinomial distribution, *Theory of Probability and its Applications* **46**(1) (2002), 103–117. doi:10.1137/S0040585X97978750. Russian original: *Teor. Veroyatnost. i Primenen.* **46**(1) (2001), 117–133.
- [18] P. Diaconis and R. C. Griffiths, Reproducing kernel orthogonal polynomials on the multinomial distribution, *Journal of Approximation Theory* **242** (2019), 1–30. doi:10.1016/j.jat.2019.01.007. arXiv:1802.07122.
- [19] J. M. Aldaz, Self-improvement of the inequality between arithmetic and geometric means, *Journal of Mathematical Inequalities* **3**(2) (2009), 213–216. arXiv:0807.1788.
- [20] H. H. Bauschke, O. Güler, A. S. Lewis and H. S. Sendov, Hyperbolic polynomials and convex analysis, *Canadian Journal of Mathematics* **53**(3) (2001), 470–488. Author manuscript.
- [21] S. Cao, S. He, Z. Li and Z. Wang, Extreme ratio between spectral and Frobenius norms of nonnegative tensors, *SIAM Journal on Matrix Analysis and Applications* **44**(2) (2023), 919–944. doi:10.1137/22M1502951. Author manuscript, Theorem 4.6 and Corollary 4.9.
- [22] C. Hanselka, Definite determinantal representations of ternary hyperbolic forms, arXiv:1411.1661v2 (2015), Theorem 1 (Helton–Vinnikov).