

Technical Supplement to The Riemann Hypothesis: Arithmetic Covariance Bounds and Complete Signed Criteria

Scott Schubert
Luminary Oddities

October 29, 2026

This document accompanies the main paper of the same title. It retains the alternative hard-cutoff and finite-source methods and their numerical applications. Definitions and principal results are in the main text; original-domain proofs and genuine-null lemmas are retained in the main paper’s appendices; unprefixed section references to those arguments refer to that PDF. All local sections and equations carry an S prefix. The complete source/error derivations for the numerical application are included here. The final sections preserve the earlier M20 certificate, the full fixed-support ridge comparison and the alternative growing-floor criterion; their verification roles are summarized in the main paper’s applications section and appendix guide. The results originate in the author’s earlier weighted-Weil and signed-frequency work [14, 12, 15, 16, 13]; the references preserve attribution. These supplementary methods complement the main finite-verifiable criterion and averaged-covariance theorem. Literal data and reproducible code are described in the package’s supplementary-material index.

Contents

S1	A cofinal finite-response criterion for RH	2
S2	Relative Green approximation and exact-response criteria	3
S3	Local inverse covariance for the averaged reference	8
S4	Explicit reference origin bounds	9
S5	Feasibility and complete exterior control	10
S6	Source-aware positive receiving	15
S7	Complete observation under moving arithmetic cutoffs	21
S8	The hard-cutoff mean and intrinsic receiver rank	24
S9	The full residual covariance and the signed return	26
S10	A complete conditioned construction	28
S11	The necessary fixed-source dimension	31
S12	Complete joint sources and stable row elimination	32
S13	A complete 4473-column source certificate	34
S14	Complete source bounds with uniform null cost	37
S15	Protected critical representers and their spectral floor	38
S16	Origin compression and fixed-window separation	39
S17	Source and error contracts for the finite-reference application	40
S18	Arithmetic bounds and complete signed-deficit coverage	49

S19	Actual-prime discrepancy and physical packet energy	53
S20	Verification of the complete supported example	57
S21	Verification of the second moving-window certificate	61
S22	Arithmetic certificate transfers in original profile coordinates	63
S23	The earlier complete supported certificate	74
S24	Complete fixed-support ridge continuation	75
S25	An alternative growing-floor arithmetic matrix criterion	76

S1 A cofinal finite-response criterion for RH

The complete covariance criterion admits a second, explicit supported-form reduction. It keeps the original polar condition and physical norm while replacing the infinite family of positive Green responses by logarithmically many features. The receiving complement remains infinite and has a proved floor; its exact inverse occurs in the finite Schur matrix below.

Use the odd-half unitary identification with $L^2(0, L)$ and put

$$p(x) = \sinh(x/2), \quad \mathcal{F}_L = p^\perp, \quad e_q(x) = \sqrt{2/L} \sin(k_q x), \quad k_q = (q + 1/2)\pi/L.$$

Let $D_L e_q = a(k_q) e_q$. The complete supported archimedean form is $D_L + R_L$, where

$$R_L = \sum_{j \geq 0} r_j \otimes r_j, \quad r_j(x) = \frac{e^{-c_j(L-x)} - e^{-c_j(L+x)}}{\sqrt{1 + e^{-2c_j L}}}, \quad c_j = 2j + \tfrac{1}{2}.$$

The Green identity and original-domain closure are proved in the main paper, Appendix E. Section S2 below proves the relative-response and joint estimates used here. Write $R_{L,J}$ for the first J responses and $V_L = m_0 - C_L^{\text{ar}}$ for the complete bounded arithmetic operator on the supported interval, including every active prime power. Thus $\|V_L\| \leq B_L := 6 + 8(\log 2)e^L$ by (10). Restrictions of these forms to $\mathcal{F}_L$ are understood throughout. Precisely, if $I = (-L, L)$ and $\mathcal{U}_L$ is the odd-half unitary map,

$$C_L^{\text{ar}} = \mathcal{U}_L \left[1_I \sum_{2 \leq n < e^{2L}} \frac{\Lambda(n)}{\sqrt{n}} (S_{\log n} + S_{-\log n}) 1_I \right] \mathcal{U}_L^*.$$

This transport retains the reflected channel as well as same-side shifts; the endpoint overlap has measure zero. Consequently the complete supported profile form $D_L + R_L + V_L + M^{-1}s^{-2}$ is exactly the original $H + M^{-1}I$ under $f = sh$ and this unitary map.

Theorem S1.1 (Cofinal finite-response criterion). *For integers $M \geq 3$ define*

$$\begin{aligned} L_M &= \tfrac{1}{2} \log(10 \log M / \pi) + 2, & \kappa_M &= M^{-1}, \\ B_M^{\text{ar}} &= 6 + 8(\log 2)e^{L_M}, & J_M &= \lceil \tfrac{1}{2} \log M \rceil, \\ N_M &= \lceil L_M e^{2B_M^{\text{ar}}} / \pi \rceil. \end{aligned}$$

Let U_M be a profile isometry onto $P_{\mathcal{F}_{L_M}} \text{span}(e_0, \dots, e_{N_M-1})$, and let Q_M be its orthogonal complement in $\mathcal{F}_{L_M}$. Form the complete finite-response operator

$$S_{0,M} = D_{L_M} + R_{L_M, J_M} + V_{L_M} + M^{-1}s^{-2}.$$

Set $T_{0,M} = S_{0,M}|_{Q_M}$ and

$$\mathbf{K}_{0,M} = U_M^* S_{0,M} U_M - (Q_M S_{0,M} U_M)^* T_{0,M}^{-1} (Q_M S_{0,M} U_M). \quad (\text{S1})$$

Then $T_{0,M} \geq \frac{1}{3}A_{Q_M} > 0$, where A_{Q_M} is the form restriction of $D_{L_M} + B_M^{\text{ar}} + M^{-1}s^{-2}$. The Riemann hypothesis is equivalent to $\mathbf{K}_{0,M} \succeq 0$ along some unbounded sequence of integers M . Under RH these matrices are strictly positive for every integer $M \geq 2^{20}$. Moreover $N_M = M^{o(1)}$ and $J_M = O(\log M)$.

Proof. Orthogonality to the projected modes implies orthogonality to the original first N_M DN modes. Since $a(k) \geq \log k$ for $k \geq 1$, the diagonal form on this complement is at least $2B_M^{\text{ar}}$. The signed arithmetic is bounded below by $-B_M^{\text{ar}}$ and $R_{L_M, J_M} \succeq 0$. Comparing coefficients gives $T_{0,M} \geq \frac{1}{3}A_{Q_M}$, with every receiving direction retained. The complete form Schur identity therefore makes $\mathbf{K}_{0,M} \succeq 0$ equivalent to $S_{0,M} \succeq 0$.

The omitted responses are positive, so this matrix inequality implies $q_H[h] + M^{-1}\|h\|_2^2 \geq 0$ for every original polar-free test supported in $(-L_M, L_M)$. Since $L_M \rightarrow \infty$, every smooth compact test is admitted eventually. Taking the unbounded sequence and then the original form closure proves Weil positivity and hence RH by the odd converse.

Conversely, the complete relative-response estimate in Section S2 gives $0 \leq R_{L_M} - R_{L_M, J_M} \leq M^{-2}(D_{L_M} + B_M^{\text{ar}} + M^{-1}s^{-2})$. Its full physical form sandwich, with $\|s\|_\infty < 3$, yields under RH

$$sS_{0,M}s \geq \left[\frac{1 - M^{-2}}{M} - \frac{18B_M^{\text{ar}}}{M^2} \right] I > 0$$

For an explicit admission, $B_M^{\text{ar}} < 6 + 120\sqrt{\log M}$ by $\log 2 < 1$, $e^4 < 64$ and $\pi > 3$. This upper divided by M decreases for $M \geq e$. At $M = 2^{20}$ it is below $606/2^{20}$, and $2^{-40} + 10908/2^{20} < 1/2$. The displayed lower is consequently at least $1/(2M)$ for every $M \geq 2^{20}$, proving strict positivity of the Schur matrix. Finally $B_M^{\text{ar}} = O(\sqrt{\log M})$, so $\log N_M = O(\sqrt{\log M})$ and the stated resource bounds follow. $\square$

The matrix in (S1) retains the *exact inverse of the full receiving operator*; it is not a principal compression. Theorem S1.1 isolates its sign as the remaining condition in this route. Section S2 also bounds the complete core-mixed-receiving change caused by the response reduction by $M^{-2+o(1)}$ in physical core coordinates, and preserves original-norm $o(M^{-3/2})$ allowances directly at the full-form level. The core coordinates do not make the transported receiving complement physically orthogonal; quantitative allowance tests shift the complete physical form before taking its Schur complement. The finite-sine alternative in Proposition S2.5 retains the same cofinal RH equivalence, uses only finite DN combinations, and has raw profile Gram condition below $4L_M = O(\log \log M)$. For that core the joint Schur error is $O(\sqrt{\log M}/M^2)$ in physical core coordinates.

S2 Relative Green approximation and exact-response criteria

S2.1 A whole relative-response bound

Let $R_{L,J} = \sum_{j < J} r_j \otimes r_j$ and $E_J = R_L - R_{L,J}$. First on ambient $L^2(0, L)$, for $B \geq 1$, $\kappa > 0$ define the closed positive reference form

$$A = D_L + B + \kappa s^{-2}.$$

Proposition S2.1 (Complete relative tail). *For $0 < \eta < L$, let $\sigma_\eta^2 = \text{ess sup}_{L-\eta < x < L} s(x)^2$. Then $0 \leq E_J \leq \epsilon A$, where*

$$\epsilon = \left[\sqrt{C_L \sigma_\eta^2 / \kappa} + \sqrt{\frac{e^{-2c_J \eta}}{2Bc_J(1 - e^{-4\eta})}} \right]^2. \quad (\text{S2})$$

The inequality includes every omitted response and survives restriction to the original polar-free subspace.

Proof. Write $P_e = 1_{(L-\eta, L)}$ and $P_i = I - P_e$. Inverse order gives $A^{-1} \preceq (B + \kappa s^{-2})^{-1}$, so

$$\|E_J^{1/2} P_e A^{-1/2}\|^2 \leq C_L \|P_e A^{-1} P_e\| \leq C_L \sigma_\eta^2 / \kappa.$$

For $d = L - x \geq \eta$, the exact feature obeys $0 \leq r_j(x) \leq e^{-c_j d}$. Tonelli and positivity give

$$\text{Tr}(P_i E_J P_i) \leq \sum_{j \geq J} \frac{e^{-2c_j \eta}}{2c_j} \leq \frac{e^{-2c_J \eta}}{2c_J(1 - e^{-4\eta})}.$$

Using $\|A^{-1/2}\| \leq B^{-1/2}$ bounds the interior factor. Add $E_J^{1/2} P_e A^{-1/2}$ and $E_J^{1/2} P_i A^{-1/2}$ and square their norm bound. This proves (S2) as a form inequality. The discontinuous spatial multipliers are used only as bounded L^2 operators on inverse square roots; they are not assumed to preserve the form domain. Apply the positive-series arguments first to partial sums and pass monotonically. No unweighted whole-tail norm convergence is used. $\square$

Corollary S2.2 (Explicit logarithmic response schedule). *Let $U = \log M$, $M \geq e$, and*

$$L = \frac{1}{2} \log(10U/\pi) + 2, \quad \kappa = M^{-1}, \quad J = \max(1, \lceil U/2 \rceil).$$

Then $0 \leq E_J \leq M^{-2} A$ for every $B \geq 1$.

Proof. Use $\eta = 1$ and the main paper's theta envelope $s(u)^2 \leq 169e^{4u - \pi e^{2u}}$. The envelope decreases on $u \geq 0$; since $L > 1$,

$$\sigma_1^2 \leq 169(10e^2/\pi)^2 U^2 M^{-10e^2}.$$

For $U \geq 1$, $C_L \leq 5U$, $e^2 < 8$, $e^2 > 7$ and $\pi > 3$. The square of the first term in (S2), divided by M^{-2} , is therefore at most

$$845(80/3)^2 U^3 e^{-67U} \leq 845(80/3)^2 e^{-67} < 2^{-47} < \frac{1}{4}.$$

Here $U^3 e^{-67U}$ decreases for $U \geq 1$, its constant is below 2^{20} , and $e > 2$. For the second squared term use $4J \geq 2U$, $c_J \geq 5/2$ and

$$[5(1 - e^{-4})]^{-1} < \frac{1}{4}.$$

Both squared contributions are below $M^{-2}/4$; their square-root sum is below M^{-1} . All constants are elementary admissions, not sampled floating inequalities. $\square$

S2.2 Preserving the original-norm allowance

On the original polar-free supported space write

$$S_f = D_L + R_L + V_L + \kappa s^{-2}, \quad \|V_L\| \leq B, \quad S_{f,J} = S_f - E_J.$$

Since $A \preceq S_f + 2BI$, the complete relative bound implies

$$(1 - \epsilon)S_f - 2\epsilon BI \preceq S_{f,J} \preceq S_f. \tag{S3}$$

Congruence by the same s , with $\|s\|_\infty < 3$, gives

$$(1 - \epsilon)S_h - 18\epsilon BI \preceq S_{h,J} \preceq S_h. \tag{S4}$$

At the displayed growing window the elementary arithmetic budget is

$$B_M = 6 + 8(\log 2)\sqrt{(10e^4/\pi)\log M} = O(\sqrt{\log M}).$$

Thus $18B_MM^{-2} = o(M^{-3/2})$. Equation (S4) proves that existence of original physical allowances $S_h \succeq -\delta_M I$, $\delta_M = o(M^{-3/2})$, is equivalent to existence of such allowances for $S_{h,J}$: take $\delta'_M = (1 - M^{-2})\delta_M + 18B_MM^{-2}$ in one direction and use $S_h \succeq S_{h,J}$ in the other. This uses the full form, not merely a core matrix. Nonvanishing extra finite corrections require a complete two-sided budget; the same power-two deduction then needs that total budget to be $o(\sqrt{M})$.

S2.3 An explicit paid core and its energy

Let $P_{\mathcal{F}}$ be the projection onto $p^\perp$ and take

$$N = \lceil Le^{B+\beta}/\pi \rceil, \quad \text{ran } U = P_{\mathcal{F}} \text{span}(e_0, \dots, e_{N-1}), \quad U^*U = I,$$

where $\beta > 0$. A vector in the complementary subspace is orthogonal to each original e_q , $q < N$. Since a is increasing and $a(k) \geq \log k$ for $k \geq 1$, its diagonal energy is at least $B + \beta$. Put $Q = I - UU^*$ on $\mathcal{F}$, and let A_Q denote the form restriction of A to Q . For both the complete and partial-response receiving forms,

$$T \succeq cA_Q, \quad T_0 \succeq cA_Q, \quad c = \frac{\beta}{\beta + 2B}. \quad (\text{S5})$$

Indeed $(1 - c)D_L + R_{L,J} + (1 - c)\kappa s^{-2} + V_L - cB$ is bounded below by $(1 - c)\beta - 2cB = 0$. Every response and receiving direction is accounted for.

Lemma S2.3 (Projected-polar core control). *For $L \geq 1$, the preceding core has dimension N and*

$$D_{\text{core}} := \|U^*D_LU\| \leq 4L(17k_N^2 + 68)(1/4 + k_N^2)^2. \quad (\text{S6})$$

Its projected-column Gram has condition number at most $2L(1/4 + k_N^2)^2$.

Proof. Direct integration gives

$$p_q = \sqrt{2/L}(-1)^q \frac{\frac{1}{2} \cosh(L/2)}{1/4 + k_q^2}, \quad \|p\|^2 = (\sinh L - L)/2.$$

Consequently $p \in \text{Dom } D_L$. With $\bar{p} = p/\|p\|$ and $\delta_N = \|(I - P_N)\bar{p}\|^2$,

$$\delta_N \geq |\langle \bar{p}, e_N \rangle|^2 \geq [2L(1/4 + k_N^2)^2]^{-1}.$$

The raw projected-column Gram is $I - vv^*$, $v = P_N\bar{p}$ in DN coordinates; its eigenvalues are 1 and $\delta_N > 0$. This proves the dimension and Gram assertions and gives an exact rank-one orthonormalization formula if desired.

The positive digamma series yields $a(k) \leq 2k^2 \sum c_j^{-3} < 17k^2$: its first coefficient is 16 and the rest is at most $\frac{1}{4} \sum_{j \geq 1} j^{-3} \leq 3/8$. The DN Laplacian form identity gives

$$D_L[\bar{p}] \leq 17 \frac{\|p'\|^2}{\|p\|^2} = 17 \frac{\sinh L + L}{4(\sinh L - L)} < 68 \quad (L \geq 1).$$

The last quotient is below 4 because $L/\sinh L$ decreases and $\sinh 1 > 7/6$. For $w \in \text{ran } P_N$, $v = P_{\mathcal{F}}w$ satisfies

$$\|v\|^2 \geq \delta_N \|w\|^2, \quad D_L[v] \leq 2(17k_N^2 + 68)\|w\|^2.$$

Combining proves (S6). In particular the projected columns are legal operator-domain vectors. This is the projected DN core, not the smaller intersection of an unprojected finite prefix with $p^\perp$. $\square$

S2.4 Joint core, mixed, and receiving error

Let $S = S_0 + E$, where S_0 retains the first J responses, and $0 \preceq E \preceq \epsilon A$. All operators here act on the same polar-free profile space. Their form domains equal $\text{Dom } A^{1/2}$. Put $d = 1 + C_L/B$. The bounds $-A \preceq S_0 \preceq dA$ and (S5) hold without a smallness assumption on ϵ . Let

$$H_0 = U^* S_0 U, \quad C_0 = Q S_0 U, \quad Z_0 = T_0^{-1} C_0, \quad K_0 = H_0 - C_0^* T_0^{-1} C_0,$$

and define K analogously for S . Inverse pairings are form-dual pairings when needed; A_Q^{-1} is the inverse of the restricted form, not $Q A^{-1} Q$.

Proposition S2.4 (Complete joint Schur budget). *With the preceding objects,*

$$0 \preceq K - K_0 \preceq \epsilon(1 + d/c)^2 U^* A U. \quad (\text{S7})$$

If $W = U^ s^{-2} U$ is the complete physical core Gram, then*

$$\|W^{-1/2}(K - K_0)W^{-1/2}\| \leq \epsilon(1 + d/c)^2 [9(D_{\text{core}} + B) + \kappa]. \quad (\text{S8})$$

On the growing window, $\beta = \omega B$ with fixed $\omega > 0$ gives $c = \omega/(\omega+2)$, $N = M^{o(1)}$ and $D_{\text{core}} = M^{o(1)}$. Hence with the explicit response schedule the right side is $M^{-2+o(1)} = o(M^{-3/2})$.

Proof. The relative form norm of S_0 is at most d . Thus $C_0^* A_Q^{-1} C_0 \preceq d^2 U^* A U$. Coercivity of T_0 and form Cauchy–Schwarz applied to $T_0 Z_0 = C_0$ give $Z_0^* A_Q Z_0 \preceq (d/c)^2 U^* A U$. The triangle inequality in the A energy norm, in every coefficient direction, therefore gives

$$(U - Z_0)^* A (U - Z_0) \preceq (1 + d/c)^2 U^* A U.$$

Variational minimization over the entire tail yields $K \succeq K_0$. Testing S at the exact S_0 minimizer $-Z_0$ gives $K \preceq K_0 + (U - Z_0)^* E (U - Z_0)$, proving (S7). The square retains all core–response, mixed, and receiving terms together.

Since $W \succeq I/9$,

$$W^{-1/2} U^* A U W^{-1/2} \preceq [9(D_{\text{core}} + B) + \kappa] I.$$

The ridge becomes exactly κI here, with no factor $\max s^{-2}$. At the declared window $B = O(\sqrt{\log M})$, $k_N = \exp(O(\sqrt{\log M}))$ and $L = O(\log \log M)$. Lemma S2.3 proves the asserted growth, completing the quantitative comparison. $\square$

The Gram W normalizes physical images of the core columns only. The profile-orthogonal receiving complement generally transports to a nonorthogonal physical complement. Thus (S8) is a finite-coordinate shorted-matrix bound, not a whole-physical-operator norm bound. For a quantitative global allowance, shift the complete physical form first, or pay the corresponding lift/projection stability. Exact nonnegativity transfers by the form Schur identity; the global allowance comparison (S4) does not require a core-to-tail lift inference.

S2.5 A finite-sine alternative with a sharper joint budget

The following alternative combines the same relative-tail estimate with a finite pole-free intersection core. It avoids a pole-profile input action and improves the joint energy bound.

Proposition S2.5 (Stable finite-sine core). *Suppose $L \geq 3$, $B = 6 + 8(\log 2)e^L$, and take*

$$N' = \lceil Le^{2B+1}/\pi \rceil, \quad \text{ran } U' = \text{span}(e_0, \dots, e_{N'-1}) \cap p^\perp, \quad U'^* U' = I.$$

In this proposition Q, K, K_0, W and all receiving and mixed blocks are rebuilt from U' ; no block from the preceding projected core is reused. Both full and finite-response receiving forms on this complement satisfy $T, T_0 \succeq A_Q/3$. A finite DN generating list has Gram condition number below $4L$, and $D_{\text{core}} \leq 2B + 7$. On the growing support, for every integer $M \geq 2^{20}$, the joint bound (S8) therefore improves to

$$\|W^{-1/2}(K - K_0)W^{-1/2}\| \leq \frac{49[27B + 63 + M^{-1}]}{M^2} = O\left(\frac{\sqrt{\log M}}{M^2}\right). \quad (\text{S9})$$

The cofinal RH criterion also holds with this $(N' - 1)$ -dimensional core.

Proof. Let $\delta_n = \|(I - P_n)\bar{p}\|^2$. The exact pole coefficients and $\cosh(L/2)^2/(\sinh L - L) \leq 2$ for $L \geq 3$ give

$$\delta_n \leq \frac{2L^3}{\pi^4(n + 1/2)^3}, \quad n \geq 1.$$

For the ratio bound use $\sinh L - L \geq e^L/4$ and $\cosh(L/2)^2 \leq e^L/2$; the former holds at 3 and its difference increases. For the series bound, the decreasing summands are bounded by their first term plus their integral. A vector $f \perp \text{ran } U'$ in $p^\perp$ has $P_{N'}f$ proportional to $P_{N'}p$. The exact pole constraint and Cauchy–Schwarz imply $\|P_{N'}f\|^2 \leq \delta_{N'}\|f\|^2$. Hence

$$D_L[f] \geq \{a(k_{N'})(1 - \delta_{N'}) + a(k_0)\delta_{N'}\}\|f\|^2.$$

Put $X = 2B + 1 \geq 2$. Then $e^X \leq k_{N'} < e^X + 2$, while the main logarithmic symbol bounds give $X \leq a(k_{N'}) < X + 6$. Also $\delta_{N'} \leq (2/\pi)e^{-3X}$, so its diagonal leakage is below

$$\frac{2}{3}(X + 6)e^{-3X} \leq 16/(3e^6) < 1/12 < 1.$$

It follows that $D_L|_Q \geq 2B$ and therefore $T, T_0 \geq A_Q/3$. On the core itself $D_{\text{core}} \leq a(k_{N'}) < 2B + 7$.

Let $v_q = \langle e_q, p \rangle$, with $v_0 > 0$, and use the finite list $w_j = v_0 e_j - v_j e_0$, $1 \leq j < N'$. Its Gram is $v_0^2 I + v_{\text{tail}} v_{\text{tail}}^*$; thus its condition number is at most

$$\frac{\|p\|^2}{v_0^2} < 2L(1/4 + \pi^2/(4L^2))^2 < 4L.$$

These columns require only finite primitive DN input actions. The complete output polar projection and physical Gram are still retained. Finally $C_L < B$ for this elementary budget, so $d < 2$ and $(1 + d/c)^2 < 49$. Insert $D_{\text{core}} \leq 2B + 7$ in (S8). The specified onset ensures $L_M \geq 3$. The RH equivalence follows by the same exact Schur and full-form argument, with this paid complement. $\square$

S2.6 A finite-feature residual test with all mixed terms

For completeness, a trial Z into the full receiving form domain gives an alternative sufficient certificate. Let $T = S|Q \succeq cA_Q$, $H = U^*SU$, $C = QSU$, and set

$$\begin{aligned} L_Z &= C^*Z + Z^*C - Z^*TZ, & L_0 &= C_0^*Z + Z^*C_0 - Z^*T_0Z, \\ r_0 &= C_0 - T_0Z, & K_r &= r_0^*A_Q^{-1}r_0, \\ G &= (U - Z)^*A(U - Z). \end{aligned}$$

For every $\tau > 0$,

$$H - C^*T^{-1}C \succeq H_0 - L_0 - c^{-1}\{(1 + \tau)K_r + (1 + \tau^{-1})\epsilon^2G\}. \quad (\text{S10})$$

Indeed the full residual is $r = r_0 + QE(U - Z)$. Relative positivity of E gives the joint A_Q -dual Gram of the second term at most ϵ^2G . Weighted Young and inverse order bound $r^*T^{-1}r$ by the displayed error. The exact residual identity and

$$(H - L_Z) - (H_0 - L_0) = (U - Z)^*E(U - Z) \succeq 0$$

then prove (S10). This is a joint matrix inequality, not a collection of diagonal tests. A numerical use must enclose the complete K_r , G , constraints and reference inverse; its cofinal budget includes the actual growth of G/c . The theorem does not assert that this finite-feature signed test has succeeded, nor that the full receiving inverse has become a finite principal compression.

S3 Local inverse covariance for the averaged reference

For $k = 1$ the continuous prime integral is exactly

$$\int_1^\infty t^{-1/2+i\xi} \min(1, M/t) dt = \frac{M^{1/2+i\xi}}{1/4 + \xi^2} - \frac{1}{1/2 + i\xi}.$$

Its phase-mean negative receiver, divided by $\sqrt{M}$, is

$$\mathcal{C}_Q = \alpha Q M_s L_0^{-1} M_s Q, \quad \alpha = 8/\pi, \quad L_0 = 1 - 4d^2/du^2. \quad (\text{S11})$$

Here Q is an original orthogonal projection on the odd space, with a finite constraint space $Z = \ker Q$. Write $\|g\|_{1,4}^2 = \int(|g|^2 + 4|g'|^2)$.

Theorem S3.1 (Full inverse-form source covariance). *For $h \in \text{ran } Q$,*

$$\|\mathcal{C}_Q^{-1/2}h\|^2 = \frac{\pi}{8} \inf_{z \in Z: (h+z)/s \in H^1} \|(h+z)/s\|_{1,4}^2, \quad (\text{S12})$$

with value infinity when the set is empty. For any finite odd H^1 profile family g_i and $e_i = Q(sg_i)$ this yields the simultaneous Gram inequality

$$[\langle \mathcal{C}_Q^{-1/2}e_i, \mathcal{C}_Q^{-1/2}e_j \rangle] \preceq \frac{\pi}{8} [\langle g_i, g_j \rangle + 4\langle g'_i, g'_j \rangle].$$

Equality holds when Z is any finite span of the original polar row and canonical derivative-null rows. All statements allow arbitrary finite source and row counts.

Proof. Factor $\mathcal{C}_Q = \alpha AA^*$ with $A = Q M_s L_0^{-1/2}$. The actual-range identity $\text{ran}(AA^*)^{1/2} = \text{ran } A$ and polar decomposition identify the inverse form with the minimum squared preimage norm [6]. Writing $g = L_0^{-1/2}a$ translates $Aa = h$ to $sg - h \in Z$, and $\|a\|^2 = \|g\|_{1,4}^2$. The chosen profiles give the matrix upper on every common coefficient vector.

More exactly, project the profiles orthogonally in H^1 , with this norm, off $\{g \in H^1 : sg \in Z\}$. Their projected Gram gives equality in (S12). This kernel is zero for the canonical rows. Indeed $b/s = \sinh(u/2)$, and for $z_{2j+1} = (\psi^{(2j+3)} - \psi^{(2j+1)})/4/s$,

$$\frac{z_{2j+1}(u)}{s(u)} \sim \frac{(-2\pi)^{2j+3}}{2} e^{(4j+13/2)u} \quad (u \rightarrow +\infty).$$

To obtain this, the first theta atom derivative polynomial has leading coefficient $4(-2)^m$ at order m , by the recurrence $P_{m+1} = (1/2 - 2r)P_m + 2rP'_m$. At fixed order the higher atoms contribute a polynomial times e^{-3r} relative to the first. The exponents of these rows and b/s are distinct, so a nonzero finite combination is not in L^2 . This proves the stated equality. $\square$

If the full profile Gram is bounded by KD and the full profile-error Gram by $\eta^2 D$ in this same H^1 metric, the resulting inverse-form Grams differ in either order by at most $(\pi/8)(2\sqrt{K}\eta + \eta^2)D$. Expand each common coefficient combination and apply Cauchy–Schwarz before passing to coordinates. The exact profile projection in the proof is a contraction, so the same error transport applies to every finite constraint space.

Lemma S3.2 (Regularized preimage identity). *For $a, \kappa > 0$ and $e = Q(sg)$, $g \in H_{\text{odd}}^1$,*

$$\langle e, (a\mathcal{C}_Q + \kappa I)^{-1}e \rangle = \inf_{f \in H_{\text{odd}}^1} \left\{ \frac{\|f\|_{1,4}^2}{a\alpha} + \frac{\|Q[s(g-f)]\|^2}{\kappa} \right\}. \quad (\text{S13})$$

Proof. For $B = \sqrt{a\alpha}QM_sL_0^{-1/2}$, minimize $\|z\|^2 + \kappa^{-1}\|e - Bz\|^2$. Its value is $\langle e, (BB^* + \kappa I)^{-1}e \rangle$. Substitute $f = \sqrt{a\alpha}L_0^{-1/2}z$. This is the classical approximate-preimage regularization identity in the present weighted and projected space [17]. $\square$

S4 Explicit reference origin bounds

Let $U = \log M$, $\Omega = L/U$, and use the actual smoothed negative symbol $n_{M,1}$. On the *entire* profile space $\mathcal{Y}_{M,L}^{\text{org}} = L^2([0, \Omega], n_{M,1}(\xi) d\xi/\pi)$ put

$$g_\phi(u) = \int_0^\Omega n_{M,1}(\xi)\phi(\xi)\sin(\xi u)\frac{d\xi}{\pi}, \quad E\phi = Q(sg_\phi), \quad D_\phi = \|\phi\|_{\mathcal{Y}_{M,L}^{\text{org}}}^2.$$

The reference $\mathcal{C}_Q$ is exactly (S11).

Theorem S4.1 (Uniform ridged origin cap). *For every $L > 0$, every integer $M \geq 64$ and every finite original constraint space,*

$$E^*(\sqrt{M}\mathcal{C}_Q + M^{-1}I)^{-1}E \preceq K(M, L)I_{\mathcal{Y}_{M,L}^{\text{org}}}, \quad (\text{S14})$$

where, with $R = \frac{1}{2}\log(2\log M)$,

$$K(M, L) = \frac{160}{3}L^3 \left(\frac{R}{\log M} \right)^3 + \frac{4096\pi^2}{3}L^3M^{3/2-2\pi}.$$

In particular $K(M, L) = O_L((\log \log M / \log M)^3)$, and $K(M, 1) < 1/10$ for every integer $M \geq 2^{20}$.

Proof. The elementary prime bound and $\mathbb{E}e^{v/2} = 2$ give $n_{M,1} \leq 6 + 16(\log 2)\sqrt{M} < 16\sqrt{M}$. Thus, for $A_M = \int_0^\Omega n_{M,1}(\xi)\xi^2 d\xi/\pi$,

$$A_M \leq \frac{16\sqrt{M}L^3}{3\pi U^3}, \quad |g_\phi(u)|^2 \leq D_\phi A_M u^2, \quad |g'_\phi(u)|^2 \leq D_\phi A_M.$$

These common-profile estimates are Cauchy–Schwarz and $|\sin(\xi u)| \leq |\xi u|$. The finite frequency band puts g_ϕ in H^1 . Choose even χ_R equal to one on $[-R, R]$, zero outside $[-2R, 2R]$, and linear on each intervening interval. Then $|\chi'_R| \leq 1/R$, and direct integration gives

$$\|\chi_R g_\phi\|_{1,4}^2 \leq 80D_\phi A_M R^3, \quad \|Q[s(g_\phi - \chi_R g_\phi)]\|^2 \leq D_\phi A_M T_R, \quad T_R = \int_{|u|>R} s(u)^2 u^2 du.$$

Indeed $\int_{-2R}^{2R} u^2 du = 16R^3/3$ and the derivative square is at most $(56/3)D_\phi A_M R$; multiplication by four gives the constant 80 for $R \geq 1$.

The complete theta estimate $s^2 \leq 16r^2 e^{-r}$, $r = \pi e^{2|u|}$, and the two-arm Jacobian $du = dr/r$ give

$$T_R \leq 16 \int_{r_R}^{\infty} r^3 e^{-r} dr \leq 32r_R^3 e^{-r_R}, \quad r_R = \pi e^{2R} \geq 6.$$

The last integral has logarithmic slope at most $-1/2$. Use the trial $f = \chi_R g_\phi$ in (S13), with $a = \sqrt{M}$, $\kappa = M^{-1}$ and $r_R = 2\pi U$. The two costs are exactly bounded by the two terms of $K(M, L)D_\phi$. This proves the inequality on every profile, hence as a full observation operator and for every common finite source combination. Projection contraction makes it uniform in all finite original constraints. Both terms decrease on $M \geq 64$. Directed scalar evaluation gives $K(2^{20}, 1) < 0.091767 < 1/10$, completing the explicit threshold. $\square$

Corollary S4.2 (Complete positive continuum-model cap). *Define the positive continuum receiver*

$$P_M^{\text{cont}} = Q M_s \left[\frac{8\sqrt{M}}{1 + 4D^2} (-\cos(UD))_+ \right] M_s Q.$$

For the same actual origin observation E ,

$$E^*(P_M^{\text{cont}} + M^{-1}I)^{-1}E \preceq 2K(M, L)I.$$

For $L = 1$ this cap is below $1/5$ for every $M \geq 2^{20}$.

Proof. Lemma 5.1 gives error $e_M = 12M^{5/2}e^{-\pi(M-1)} \leq 1/(2M)$ on $M \geq 64$. The gate holds at 64 using $\pi > 3$, $e > 2$, and the majorant decreases thereafter. Hence $P_M^{\text{cont}} + M^{-1}I \succeq \frac{1}{2}(\sqrt{M}C_Q + M^{-1}I)$. Inverse order and Theorem S4.1 finish the proof. $\square$

S5 Feasibility and complete exterior control

The preceding certificates retain exact signed information. This section identifies the costs of two simpler bounds and locates the earlier scalar complementary criterion within the complete construction.

Proposition S5.1 (Cost of the positive majorant). *For the explicit E_0 in the proof above, with $m \sim b \log M / \log 10$,*

$$\|E_0^* R_M (M^{-b}) E_0\| \geq \frac{c_b \sqrt{m}}{\sqrt{M} (\log m)^3}$$

for all sufficiently large M . Thus this positive-majorant inverse cost is not $o(M^{-1/2})$.

Proof. Choose compact smooth odd f with $\int f(u) \sinh(u/2) du = 0$ but $\int u f(u) du \neq 0$. Two odd bump pairs construct such an f , since u and $\sinh(u/2)$ are linearly independent on a positive interval. Then $y = f/s$ is legal, $|\hat{f}(\xi)| \geq c|\xi|$ near zero, and for large M , $E_0^* y = T_m f$. On $|\xi| \leq 1/(2 \log m)$ every $\cos(\xi \log n) > 1/2$ for $n \leq m$. The prime number theorem gives $\sum_{n \leq m} \Lambda(n)/\sqrt{n} \geq c\sqrt{m}$; boundedness of a near zero therefore yields $v_m(\xi) \geq c\sqrt{m}$ on this interval. Plancherel gives $\|E_0^* y\|^2 \geq c\sqrt{m}/(\log m)^3$. The full bound $\|N_M\| \leq 144\sqrt{M}$ and the signed identity imply $P_M[y] + M^{-b}\|y\|^2 = O_y(\sqrt{M})$. The inverse variational quotient proves the assertion. $\square$

This concerns the particular positive majorant, not the signed covariance excess. It explains why the next construction retains the exact null remainder rather than trying to sharpen this majorant by constants alone.

The technical supplement's arithmetic-gain section supplies a sharper compact arithmetic price, a complete residual optimizer, and matrix tests for whether the resulting improvement covers a signed certificate deficit. These retain the same receiving core and every mixed term.

S5.1 The complete exterior in the actual inverse metric

The next estimate identifies source directions that a constructive null head must capture globally. Put $R_M = (P_M + \kappa I)^{-1}$ and $G_M = E_M^* R_M E_M$ on the base original polar recipient.

Proposition S5.2 (Actual exterior payments). *Let $F = F_M \rightarrow \infty$ satisfy*

$$F^4/\sqrt{M} \longrightarrow 0, \quad F^3 \bar{e}_M \longrightarrow 0,$$

and let Π_F be source multiplication by $1_{[F, 2F]}$. Uniformly for $0 < \kappa \leq 1$, for all sufficiently large M ,

$$\text{Tr}(E_M \Pi_F E_M^*) \geq c\sqrt{M}/F, \tag{S15}$$

$$\text{Tr}(\Pi_F G_M \Pi_F) \geq c/F, \quad \|\Pi_F G_M \Pi_F\| \geq c/F^2. \tag{S16}$$

These conclusions hold, in particular, for $F = \exp(A\sqrt{\log M})$ with any fixed $A > 0$.

Proof. Write $U = \log M$, $B(\xi) = 8/(1 + 4\xi^2)$ and $e_\xi = Q_b[s \sin(\xi \cdot)]$. The complete Abel formula and $a(\xi) \leq 17\xi^2$ give on $[F, 2F]$

$$\sup \left| \frac{m_M(\xi)}{\sqrt{M}B(\xi)} + \cos(U\xi) \right| \leq C(F^4/\sqrt{M} + F^3 \bar{e}_M) = o(1).$$

Thus $n_M \leq C\sqrt{M}/F^2$ there, and $n_M \geq c\sqrt{M}/F^2$ on its subset where $\cos(U\xi) \geq 1/2$. That subset occupies at least a quarter of each fixed-length interval for large M , and at least $F/4$ of $[F, 2F]$.

The exact projected source Gram is

$$\begin{aligned} \langle e_\xi, e_\zeta \rangle &= \frac{1}{2} \int s^2 \cos((\xi - \zeta)u) du - \frac{1}{2} \int s^2 \cos((\xi + \zeta)u) du \\ &\quad - \frac{\langle s \sin(\xi \cdot), b \rangle \langle b, s \sin(\zeta \cdot) \rangle}{\|b\|^2}. \end{aligned}$$

Riemann–Lebesgue therefore gives $\|e_\xi\|^2 \rightarrow \|s\|_2^2/2 > 0$. Differentiating se_ξ shows $\|(se_\xi)'\|^2 \leq C(1 + \xi^2)$. The fixed hard-cutoff decomposition of the original form is $H = A + B_0$ with B_0 bounded and $A[h] = \int a(\xi) |\widehat{sh}(\xi)|^2 d\xi/(2\pi)$. Consequently these sources belong to the original form domain and $q_H[e_\xi] \leq C(1 + \xi^2)$. Since $P_M = N_M + H - W_M$ and $\|N_M\| \leq 144\sqrt{M}$,

$$P_M[e_\xi] + \kappa\|e_\xi\|^2 \leq C\sqrt{M}, \quad \langle e_\xi, R_M e_\xi \rangle \geq \frac{\|e_\xi\|^4}{P_M[e_\xi] + \kappa\|e_\xi\|^2} \geq c/\sqrt{M}.$$

Here $F^2 \leq \sqrt{M}$ eventually. Integrating these estimates against $n_M(\xi)d\xi/\pi$ proves the raw and inverse-weighted trace bounds.

The norm bound requires a separate coherent source. Choose fixed $\ell > 0$ so small that $\int s^2 \cos(tu) du \geq 3\|s\|^2/4$ for $|t| \leq \ell$. The other two Gram terms tend uniformly to zero on

$[F, F + \ell]^2$, so $\langle e_\xi, e_\zeta \rangle \geq c_0 > 0$ there for large M . Let $Z = [F, F + \ell] \cap \{\cos(U\xi) \geq 1/2\}$, with $\mu(Z) = \int_Z d\xi/\pi$, and use the unit source $f = 1_Z/\sqrt{\mu(Z)}$. Its coefficients are nonnegative and $\mu(Z)$ is bounded above and below by positive constants. For $v = E_M f$, the full Gram and differentiated Bochner integral give

$$c\sqrt{M}/F^2 \leq \|v\|^2 \leq C\sqrt{M}/F^2, \quad \|(sv)'\| \leq CM^{1/4}.$$

In particular v belongs to the original form domain and $q_H[v] \leq C\sqrt{M}$. The complete positive energy satisfies

$$P_M[v] + \kappa\|v\|^2 \leq 144\sqrt{M}\|v\|^2 + C\sqrt{M} + (\delta_M + \kappa)\|v\|^2 \leq CM/F^2.$$

The inverse variational test now yields $\langle f, G_M f \rangle \geq \|v\|^4/(P_M[v] + \kappa\|v\|^2) \geq c/F^2$. No trace-to-norm inference is used. Finally Theorem 7.1 proves $\bar{e}_M = O_D(e^{-D\sqrt{U}})$ for every fixed $D > 0$. For fixed A , choose $D > 3A$ to admit $F = e^{A\sqrt{U}}$; $F^4/\sqrt{M} \rightarrow 0$ is immediate. $\square$

If a source head is supported below F , its orthogonal complement acts identically on $[F, 2F]$, so its complete inverse remainder has norm at least c/F^2 and trace at least c/F . The same holds for the remainder of a positive-contraction head supported there. In the displayed stretched-exponential range these bounds exceed $M^{-1/2}$ asymptotically. Thus a low-band head cannot pay its entire positive remainder as $o(M^{-1/2})$, even in the actual inverse metric. Global null-polynomial heads and the directional Schur criterion remain available: these are lower bounds on a positive complement, not on $\|G_M\| - 1$.

S5.2 The scalar mixed-band criterion

Proposition S5.3 (Complete mixed-band cap). *For the same positive inverse $R_P = (P + \kappa I)^{-1}$, split the exact observation source as $E = [E_o, E_c]$. If its two diagonal caps are $E_o^* R_P E_o \preceq aI$ and $E_c^* R_P E_c \preceq bI$, and the full mixed block has norm at most ζ , then*

$$\|E^* R_P E\| \leq \frac{1}{2}[a + b + \sqrt{(a - b)^2 + 4\zeta^2}] \leq a + b \quad \text{when } \zeta \leq \sqrt{ab}.$$

Proof. Positive-form Cauchy–Schwarz pays the mixed block by $\sqrt{ab}$. More generally bound each two-vector quadratic form by the real matrix $\begin{bmatrix} a & \zeta \\ \zeta & b \end{bmatrix}$ and compute its largest eigenvalue. All mixed observation terms are retained. $\square$

Corollary S5.4 (A complementary-band cofinal criterion for RH). *Let a_M be the bound in (24) at $\Omega = 1/\log M$, and use any finite genuine-null recipient. If, along an unbounded sequence, the complete complementary observation for the same $\bar{P}_M + M^{-1}I$ has cap*

$$b_M \leq 1 - a_M + M^{-1},$$

then on the original domain

$$q_H[h] \geq - \left[d_{M,1} + \frac{1}{M} + \frac{B_{M,1}}{M+1} \right] \|h\|^2,$$

and RH holds. A paid mixed block can instead use the sharper condition in Proposition S5.3 at target $1 + 1/M$.

Proof. The proposition gives the complete cap $1 + 1/M$ including all mixed terms. Apply the bounded-negative absorption of Corollary 4.2 with $k = 1$, $\tau = 0$, $\eta = \kappa = 1/M$. Here $B_{M,1} = O(\sqrt{M})$, so the displayed allowance tends to zero; Theorem 2.2 applies. $\square$

S5.3 Why bounded null rank does not fund the scalar criterion

The sufficient scalar comparison in Corollary S5.4 loses directional covariance information. The following result gives its precise rank limitation at the displayed schedule.

Proposition S5.5 (Fixed-rank scalar obstruction). *Put $\Omega = 1/\log M$, $\kappa = M^{-1}$ and split the complete source into E_o on $[0, \Omega]$ and its entire complement E_c . For the same-form recipient obtained by removing a genuine-null space K , put*

$$a = \|E_o^* R_M E_o\|, \quad b = \|E_c^* R_M E_c\|, \quad \chi = \|E_o^* R_M E_c\|.$$

For every fixed integer $D \geq 0$ there are positive constants c_D, C_D and a finite threshold, uniform over all K with $\dim K \leq D$, such that

$$\chi^2 \geq c_D \Omega^{4D+3}, \quad b \geq 1 - C_D \Omega^{4D+7} - C_D M^{-3/2}.$$

Consequently the scalar matrix $\begin{bmatrix} a & \chi \\ \chi & b \end{bmatrix}$ is not bounded by $(1+1/M)I$ for all sufficiently large M . Any cofinal realization of that scalar target requires the null dimension to eventually exceed every fixed D .

Proof. Let $L_D = \text{span}\{z_0, \dots, z_D\}$, with the genuine nulls of Lemma G.1. Dimension gives a unit $z = \sum_{i=0}^D \alpha_i z_i \in L_D \cap K^\perp$. The derivative Fourier rule and $\hat{\psi}(0) > 0$ show, near zero,

$$|\widehat{sz}(\xi)| \geq c\xi \left| \sum_{i=0}^D (-1)^i \alpha_i \xi^{2i} \right|.$$

The complete Abel formula gives $n_{M,1} \geq c\sqrt{M}$ on $[0, \Omega]$ for large M , since $\cos((\log M)\xi) \geq \cos 1 > 0$. The positive moment matrix $G_D = (1/(2i+2j+3))_{0 \leq i,j \leq D}$ gives

$$\int_0^\Omega \xi^2 \left| \sum_{i=0}^D (-1)^i \alpha_i \xi^{2i} \right|^2 d\xi \geq \lambda_{\min}(G_D) \Omega^{4D+3} \|\alpha\|^2.$$

The fixed physical Gram of L_D bounds $\|\alpha\|$ below. Therefore $N_o[z] \geq c_D \sqrt{M} \Omega^{4D+3}$, uniformly in the moving K . All constants here are for fixed D ; no uniform conditioning in D is used.

Write $A_o = E_o^* R_M E_o$ and $C_{oc} = E_o^* R_M E_c$. The exact null identity gives

$$C_{oc} E_c^* z = (I - A_o) E_o^* z + E_o^* R_M (W_M - \kappa I) z.$$

Since $R_M^2 \preceq \kappa^{-1} R_M$,

$$\chi \sqrt{N_c[z]} \geq (1-a) \sqrt{N_o[z]} - (d_{M,1} + \kappa) \sqrt{a/\kappa}.$$

The actual origin upper is $a \leq C\Omega^3(1+\log \log M)^3$. Dividing its error term by the lower bound for $\sqrt{N_o[z]}$ gives at most $C_D M^{-3/4} \Omega^{-2D} (1+\log \log M)^{3/2} + o(1)$, which tends to zero. The complete bound $N_c[z] \leq 144\sqrt{M}$ then proves the stated lower for χ^2 . If $N_c[z] = 0$, the displayed identity already contradicts its positive right side for large M .

For the complementary lower choose a unit $z' \in \text{span}\{z_{D+1}, \dots, z_{2D+1}\} \cap K^\perp$. Its transform vanishes to order at least $2D+3$ at zero, so $N_o[z'] \leq C_D \sqrt{M} \Omega^{4D+7}$. The averaged quadratic mean proved in Theorem 7.1 is positive definite on this fixed finite-dimensional span. Polarization makes its convergence uniform on that span's unit sphere; hence $N_M[z'] \geq c'_D \sqrt{M}$ for every allowed moving choice. The exact null quotient yields

$$b \geq \frac{N_c[z']}{P_M[z'] + \kappa} \geq 1 - C_D \Omega^{4D+7} - C_D M^{-3/2}.$$

Finally put $t = 1 + 1/M$. If $b > t$ the scalar target fails immediately. Otherwise its necessary determinant inequality would imply

$$\chi^2 \leq (t - a)(t - b) \leq 2\{M^{-1} + C_D\Omega^{4D+7} + C_DM^{-3/2}\} = o(\Omega^{4D+3}),$$

contradicting the proved lower. $\square$

The proposition concerns scalarization of the blocks. It neither rejects the operator-valued Schur criterion nor establishes sufficiency of growing null rank. The regularized whole-source construction retains the signed correlations that this scalar comparison discards.

S5.4 Spectral tests for a growing genuine-null head

The directional construction has an actual compressed-spectrum test that retains the physical lift cost. It does not replace that spectrum by the unrestricted mean operator.

Proposition S5.6 (Null-compressed spectral and lift tests). *Let $\mathcal{K}$ be the closed physical span of the genuine nulls and $N_{\mathcal{K}}$ the compression of N_M to $\mathcal{K}$. For a rank- r null-valued map V with $V^*N_MV = I$, put*

$$J = E_M^*V, \quad Q = I - JJ^*, \quad \ell = \|V^*V\|, \quad b_c = \|QG_MQ\|, \quad g = \|G_M\|.$$

If $b_c \leq q < 1$, then

$$\lambda_{r+1}(N_{\mathcal{K}}) \leq \frac{q(\kappa + \delta_M)}{1 - q}. \quad (\text{S17})$$

For every eigenvalue $\lambda > 0$ of $N_{\mathcal{K}}$,

$$b_c \geq \left[\sqrt{(1 - (\kappa + \delta_M)/\lambda)_+} - \sqrt{g\lambda\ell} \right]_+^2. \quad (\text{S18})$$

Proof. Closedness of H implies $\mathcal{K} \subset \text{Dom } H$ and $H|_{\mathcal{K}} = 0$. For $z \in \mathcal{K}$ with $E_M^*z \perp \text{ran } J$, the inverse variational test at z gives

$$\langle E_M^*z, G_ME_M^*z \rangle = \langle N_Mz, R_MN_Mz \rangle \geq \frac{N_M[z]^2}{P_M[z] + \kappa\|z\|^2}.$$

Genuine nullity gives $P_M[z] = N_M[z] - \langle z, W_Mz \rangle$. Putting $t = N_M[z]/\|z\|^2$, we obtain $b_c \geq t/(t + \kappa + \delta_M)$ whenever $t > 0$. There are at most r source-orthogonality conditions on z . Compact min-max on $N_{\mathcal{K}}$ proves (S17), with eigenvalue multiplicities retained.

For the second bound choose a unit compressed eigenvector h at λ and put $f = E_M^*h$, so $\|f\|^2 = \lambda$. Since V is $\mathcal{K}$ -valued, the compressed eigenidentity gives $V^*N_Mh = \lambda V^*h$, without assuming invariance of $\mathcal{K}$ under N_M . Hence

$$\|JJ^*f\|^2 \leq \lambda^2\ell, \quad \langle f, G_Mf \rangle \geq \frac{\lambda^2}{\lambda + \kappa + \delta_M} \geq \lambda - (\kappa + \delta_M).$$

The triangle inequality in the positive G_M seminorm yields

$$\sqrt{\langle f, G_Mf \rangle} \leq \sqrt{g} \lambda \sqrt{\ell} + \sqrt{b_c \lambda}.$$

Division by $\sqrt{\lambda}$ and positive parts prove (S18). $\square$

At $\kappa = M^{-1}$, a fixed complementary gap requires the head to capture every null-compressed eigenvalue above a constant multiple of M^{-1} . If actual eigenvalues satisfy $(\kappa + \delta_M)/\lambda_M \rightarrow 0$ and $\lambda_M\ell_M \rightarrow 0$, then Theorem 7.1 and (S18) give $\liminf b_c \geq 1$. In particular, an actual eigenvalue comparable to $M^{-1/2}$ would exclude a fixed gap with $\ell_M = o(\sqrt{M})$. The spectral and lift tests are evaluated jointly with the signed complement estimate in the complete source metric.

S6 Source-aware positive receiving

The polynomial construction controls complete positive complements in well-conditioned coordinates. A second, compatible finite reduction retains the actual positive and negative parts of the signed symbol. It prices the error on its finite source rather than on the whole inverse.

Fix $M \geq 2$, put $p_M = (m_M)_+$ and $n_M = (-m_M)_+$, and let X_0 be the odd Hilbert space with the polar row and any specified finite exact rows imposed. On this same recipient define the closed positive form

$$P[v] = \int_0^\infty p_M(\xi) |\widehat{sv}(\xi)|^2 \frac{d\xi}{\pi}, \quad N[v] = \int_0^\infty n_M(\xi) |\widehat{sv}(\xi)|^2 \frac{d\xi}{\pi}.$$

Thus $H_M = P - N$ as forms. In this section P denotes this positive operator, rather than the range projection of Section S9. Every inverse and observation below uses the original Hilbert norm and the same exact-row recipient.

Theorem S6.1 (Complete inverse exhaustion and finite-source convergence). *Let $\kappa > 0$, $R = (P + \kappa I)^{-1}$, and let P_F restrict the complete positive observation to $[0, F]$. Then $R_F = (P_F + \kappa I)^{-1}$ converges strongly to R , while*

$$\|R_F - R\| = \kappa^{-1}$$

for every finite F . The same norm equality holds for every finite-rank positive lower $P_f \preceq P$. For each fixed finite column map E , however,

$$\|E^*(R_F - R)E\| \longrightarrow 0.$$

Proof. The original graph closure is $\{v : sv \in H_{\log}\}$, with the exact rows imposed, where $H_{\log}$ has Fourier weight $1 + \log(2 + |\xi|)$. Indeed the digamma series makes $1 + a$ comparable to this weight, and $p_M - a$ is bounded. The square root of the weight is submultiplicative up to a constant. Weighted Young's inequality therefore bounds multiplication by a smooth cutoff by its weighted Fourier L^1 norm. Dilated even cutoffs have uniformly bounded such norms. Schwartz density, followed by cutoff convergence in H^1 , proves cutoff convergence in $H_{\log}$. On each compact interval division by the smooth positive s , then even mollification, supplies odd smooth compact graph approximants. Multiplication back by s is bounded in $H_{\log}$ because s and its derivatives decrease rapidly. Correct the finitely many row moments by fixed compact odd dual functions; their coefficients tend to zero in the graph norm. This proves the asserted minimal-domain identification. Increasing closed-form convergence now gives strong resolvent convergence [1].

For the norm assertion first use the ambient odd space. Choose a nonzero even smooth compactly supported χ and unit odd packets $h_k = \chi \sin(ku) / \|\chi \sin(k \cdot)\|_2$. These are weakly zero. The finite-band observation is Hilbert–Schmidt, since its squared norm is bounded by $\|s\|_2^2 \int_0^F p_M d\xi / \pi$; hence P_F is compact. Consequently $R_F h_k - \kappa^{-1} h_k \rightarrow 0$. The digamma asymptotic [11] gives $p_M(\xi) = \log |\xi| + O_M(1)$. Translating a Schwartz Fourier transform and using the integrable logarithmic majorant shows, for each smooth compact g ,

$$\|(\log k)^{-1} p_M(D)(ge^{iku}) - ge^{iku}\|_2 \rightarrow 0.$$

Apply this to both sine arms of $v_k = (\log k)^{-1} s^{-2} h_k$. These compact smooth functions lie in the operator domain and satisfy $v_k \rightarrow 0$ and $(P + \kappa)v_k - h_k \rightarrow 0$. Thus $Rh_k \rightarrow 0$. Inverse order gives $0 \preceq R_F - R \preceq \kappa^{-1}I$, establishing equality. A finite-rank P_f has the same compactness argument.

For independent exact rows Z_0 , the constrained inverse extended by zero is

$$R_X = R_Y - R_Y Z_0 (Z_0^* R_Y Z_0)^{-1} Z_0^* R_Y.$$

Its correction is finite rank. Projecting the weakly zero packets onto X_0 changes them by $o(1)$, so the norm argument survives. Redundant rows are first removed. Finally strong convergence is uniform on the unit sphere of a fixed finite-dimensional source range, proving the last assertion. $\square$

S6.1 A full mixed observation residual

Let $P_f \preceq P$ be a bounded finite positive lower, $R_f = (P_f + \kappa I)^{-1}$, and $V = R_f E$. Its actual trial columns are assumed in the original positive form domain. The variational principle gives the full matrix estimate

$$0 \preceq E^* R_f E - E^* R E \preceq V^* (P - P_f) V. \quad (\text{S19})$$

To prove the upper, evaluate the complete variational formula for $\langle Ez, REz \rangle$ at Vz ; its value is $\langle Ez, R_f Ez \rangle - (P - P_f)[Vz]$. This proves the matrix order on every coefficient vector.

Here is a computable price including all cross terms. Write $P = T^* T$, where $Tv(\xi) = \sqrt{p_M(\xi)} \widehat{sv}(\xi)$ in $L^2((0, \infty), d\xi/\pi)$. Suppose

$$P_f = T^* Z (D + \rho I)^{-1} Z^* T, \quad D = Z^* Z, \quad \rho > 0,$$

and Z is supported in $[0, F_0]$. For $F \geq \max(F_0, 1)$, a trial coefficient matrix B , and $r \geq 1$, define

$$A_{\text{in}} = \int_0^F (TV - ZB)^* (TV - ZB) \frac{d\xi}{\pi}, \quad D_r = [\langle D^r(sv_i), D^r(sv_j) \rangle],$$

and $C_M = 4 + 2 \sum_{n \leq M} \Lambda(n)/\sqrt{n}$. Then

$$V^* (P - P_f) V \preceq A_{\text{in}} + \rho B^* B + \frac{C_M + \log(3F)}{F^{2r}} D_r. \quad (\text{S20})$$

Ridge-square completion gives the first two terms and the full exterior $\int_F^\infty (TV)^* TV d\xi/\pi$. The digamma series, compared with its integral, gives $a(\xi) \leq 4 + \frac{1}{2} \log(1 + 4\xi^2)$, hence $p_M(\xi) \leq C_M + \log(3\xi)$ for $\xi \geq 1$. The quotient by ξ^{2r} decreases there. Parseval proves the exterior bound with the displayed complete derivative Gram. Actual finite-band columns and row corrections have finite D_r : their weighted profiles are s^2 times bounded-band integrals, complete theta derivatives, and $s^2 \sinh(u/2)$.

The finite inverse is explicit. For $P_f = W D_p^{-1} W^*$, with its actual metric $D_p \succ 0$, set $L = W^* W + \kappa D_p$. Then

$$V = \kappa^{-1} \{E - W L^{-1} W^* E\}. \quad (\text{S21})$$

On an exactly constrained recipient use the finite inverse-row correction in the proof of Theorem S6.1. This retains the row/source, row/positive and all mixed covariances.

Theorem S6.2 (Paid source-aware cofinal reduction). *For each j let $\kappa_j > 0$ tend to zero and let $\tau_j, \delta_j \geq 0$. Take the recipient to be the original polar-free space with a finite span of genuine operator nulls removed, and on this recipient let*

$$N_{*,j} = E_j D_{n,j}^{-1} E_j^*, \quad 0 \preceq N_j - N_{*,j} \preceq \tau_j I, \quad D_{n,j} \succ 0.$$

Suppose $P_{f,j} \preceq P_j$ is a finite positive lower and, for $0 < \gamma_j < 1/2$,

$$E_j^* (P_j + \kappa_j)^{-1} E_j \preceq (1 - 2\gamma_j) D_{n,j}, \quad V_j^* (P_j - P_{f,j}) V_j \preceq \gamma_j D_{n,j}.$$

The second inequality may be certified by (S20). If the complete raw arithmetic error is δ_j and $\tau_j + \delta_j \rightarrow 0$, then

$$q_H[h] \geq -(\kappa_j + \tau_j + \delta_j) \|h\|_2^2$$

on the corresponding genuine-null complement. These certificates, with genuine operator nulls restored, imply RH.

Proof. Equation (S19) gives $E_j^*(P_{f,j} + \kappa_j)^{-1}E_j \preceq (1 - \gamma_j)D_{n,j} \preceq D_{n,j}$. The positive Schur complement is equivalent to $N_{*,j} \preceq P_{f,j} + \kappa_j I$. Add the negative approximation and complete raw errors. The original closed-form domain and exact null transport in Theorem 2.2 prove the conclusion. $\square$

Corollary S6.3 (Polynomial excess above one). *On the same original polar-free recipient with finitely many genuine operator nulls removed, suppose instead that $\eta_M, \tau_M, \delta_M \geq 0$, $\kappa_M > 0$, $0 \preceq P_{f,M} \preceq P_M$, and the full finite-source cap satisfies*

$$E_M^*(P_{f,M} + \kappa_M I)^{-1}E_M \preceq (1 + \eta_M)D_{n,M}, \quad 0 \preceq N_M - E_M D_{n,M}^{-1} E_M^* \preceq \tau_M I.$$

Let B_M be an upper bound for $\|s\|_\infty^2(|m_0| + 2\sum_{n \leq M} \Lambda(n)/\sqrt{n})$ and let δ_M pay the complete raw tail. Then, on the entire original domain,

$$q_H[h] \geq - \left\{ \delta_M + \kappa_M + \frac{\tau_M + \eta_M B_M}{1 + \eta_M} \right\} \|h\|_2^2. \quad (\text{S22})$$

In particular complete caps on a cofinal sequence with $\kappa_M = \eta_M = \tau_M = 1/M$ give original allowances $O(M^{-1/2})$ tending to zero, and hence RH.

Proof. The cap is equivalent to $N_* \preceq (1 + \eta)(P_f + \kappa I)$, hence $N \preceq (1 + \eta)P + tI$, where $t = (1 + \eta)\kappa + \tau$. The exact symbol gives $0 \preceq N \preceq BI$. Writing $q = P - N + W$, $W \succeq -\delta I$, the following identity of forms proves the absorption without bounding the unbounded P :

$$(1 + \eta)(q + \varepsilon I) = [(1 + \eta)P + tI - N] + \eta[B I - N] + (1 + \eta)[W + \delta I], \quad \varepsilon = \delta + \frac{t + \eta B}{1 + \eta}.$$

Each summand is nonnegative. Genuine-null transport returns the bound to the whole original domain. The elementary prime-power estimate (10) and $\|s\|_\infty < 3$ permit the choice $B_M = 9R_M = O(\sqrt{M})$, with R_M from (S26). Substitution yields (S22) and its stated rate. All finite-source, ridge and continuous errors are included in the cap or in τ_M . $\square$

The cap in this alternative may lie above one; its excess is paid by the complete bounded negative part. It uses a different regularization/cutoff schedule from Theorem S10.3. The source-residual enclosure (S19) can fund the finite positive receiver against either target.

S6.2 Complete strip price and integration collar

For $f_i = sv_i$ put $A_\eta = [\langle f_i(\cdot - i\eta), f_j(\cdot - i\eta) \rangle]$, with $\eta = 1/8$. For actual finite observation and genuine-row columns these analytic weighted profiles give

$$\int_F^\infty (TV)^* TV \frac{d\xi}{\pi} \preceq 2[C_M + \log(3F)]e^{-F/4}A_\eta, \quad F \geq 4. \quad (\text{S23})$$

Contour shifting gives $\widehat{f(\cdot - i\eta)} = e^{\eta\xi}\widehat{f}$; full Plancherel and the positive-half measure pay the factor two. The scalar multiplier $[C_M + \log(3\xi)]e^{-\xi/4}$ decreases for $\xi \geq 4$, proving the full matrix order.

For completeness, usable whole-source constants are

$$\|s^2(\cdot \pm i/8)\|_2 < 20, \quad \|\psi^{(m)}(\cdot \pm i/8)\|_2 \leq 32 \cdot 16^m m!, \quad \|s^2(\cdot \pm i/8) \sinh((\cdot \pm i/8)/2)\|_2 \leq 32.$$

Here is a direct complete-tail proof. Put $x = |\Re z|$, $r = \pi e^{2x}$. On $|\Im z| \leq 1/4$, the theta atom series and reflection give $|\psi(z)| \leq 7e^{x/2}r^2e^{-7r/8}$: each atom is at most $6e^{x/2}(\ell^2 r)^2e^{-7\ell^2 r/8}$ and successive ratios are below $1/8$. On the $1/8$ line, the sharper cosine bound $31/32$ gives $|s^2(z)| \leq$

$28r^2e^{-31r/32}$. The squared two-arm integral is at most $4704(16/31)^4 < 400$. Cauchy's circle of radius $1/16$ bounds the derivative by $1416^m m! e^{x/2} r^2 e^{-3r/4}$; its squared two-arm integral is at most $196 \int_0^\infty r^4 e^{-3r/2} dr < 1024$. The polar bound follows from $|\sinh(z/2)| \leq e^{x/2}$ and the same integral comparison.

An observation feature $e = s \int_0^B a(\xi) \sin(\xi u) d\xi / \pi$ therefore has weighted strip norm at most $20e^{B/8} \int_0^B |a| d\xi / \pi$. A genuine derivative-null feature has weighted profile $\psi^{(2i+3)} - \psi^{(2i+1)}/4$, so the displayed derivative bounds apply. If $sV = gA$ uses b such features and their strip norms are bounded by k_ℓ , then

$$A_\eta \preceq bA^* \text{diag}(k_\ell^2)A.$$

This finite matrix includes the actual inverse metrics and row coefficients. Under the explicit hypothesis $A_\eta \preceq e^{B_j/4+L_j} D_{n,j}$, the exterior is at most

$$2[C_{M_j} + \log(3F_j)]e^{-(F_j-B_j)/4+L_j} D_{n,j}.$$

For $\gamma_j = c\kappa_j^\rho$, $0 < \rho < 1$, it suffices that

$$\frac{F_j - B_j}{4} \geq L_j + \rho \log(1/\kappa_j) + \log \frac{8[C_{M_j} + \log(3F_j)]}{c}.$$

If $L_j = O(U_j)$, $\log B_j = O(U_j)$, $B_j \geq 1$ and $\log C_{M_j} = O(\log U_j)$, with $U_j = \log(1/\kappa_j)$, a sufficiently large fixed C permits $F_j = B_j + CU_j$. This is an integration collar for the chosen receiver. The interior residual in (S20) and the complete cap in Theorem S6.2 retain their stated hypotheses.

S6.3 Null leakage in the same positive metric

Let T now be an isometry onto finitely many genuine operator nulls inside a base recipient X_0 , before their removal; write $Q_T = I - TT^*$, $C = P + \kappa I$, $S = T^*CT$. This use of T is distinct from the observation map above. Suppose $N_* = ED_n^{-1}E^*$, $0 \preceq N - N_* \preceq \tau I$, and the raw tail has norm at most $\delta < \kappa$. Put

$$A_0 = D_n^{-1}E^*T, \quad B_* = Q_T E A_0, \quad s_0 = \kappa - \delta + \lambda_{\min}(T^*N_*T) > 0.$$

For a finite positive lower $P_f = WD_p^{-1}W^* \preceq P$ on the *base* recipient, define

$$J_B = B_*^*B_* - B_*^*W(W^*W + \kappa D_p)^{-1}W^*B_* \succeq 0.$$

Then, for every $t > 0$,

$$\kappa \|T^*C^{-1}T - S^{-1}\| \leq s_0^{-2} \|(1+t)J_B + (1+1/t)(\tau + \delta)^2 I\|. \quad (\text{S24})$$

Indeed the exact null identity $HT = 0$ gives $PT = NT - W_{\text{tail}}T$. Thus $B = Q_T PT$ differs from B_* by at most $\tau + \delta$, and $S \succeq s_0 I$. The exact trial-residual identity is

$$T^*C^{-1}T - S^{-1} = S^{-1}B^*C^{-1}BS^{-1}.$$

Inverse order, quadratic Young's inequality and (S21) give $\kappa B^*C^{-1}B \preceq \tilde{J}$, where $\tilde{J} = (1+t)J_B + (1+1/t)(\tau + \delta)^2 I$. This proves (S24). More sharply the matrix-relative target $\tilde{J} \preceq \eta S^2$ yields debit at most η by congruence with S^{-1} ; no commutation is assumed. The finite lower must be certified before removing T , so all its mixed row covariances remain present.

These null-source tests are compatible with the full-source cap but have distinct recipients. For example $P = \text{diag}(1, 1)$, $N = \text{diag}(1, 2)$, $T = e_1$ gives $HT = 0$ and zero null debit for every κ ,

while the second direction has negative signed form. The complete cap in Theorem S6.2 keeps that direction in its test.

Finally a trial inverse from a different positive reference has a paid exact bridge. For $A \succeq 0$, bounded self-adjoint $K = P - A$, $V_A = (A + \kappa)^{-1}Z$ and $G_A = Z^*V_A$, square completion gives

$$Z^*(P + \kappa)^{-1}Z \preceq G_A - V_A^*KV_A + \kappa^{-1}V_A^*K^2V_A.$$

The identity uses residual $(P + \kappa)V_A - Z = KV_A$, followed by $(P + \kappa)^{-1} \preceq \kappa^{-1}I$. It preserves the full covariance of the changed reference.

S6.4 A complete 547-source inverse-action example

The source-aware viewpoint also gives a quantitative action certificate for a fixed finite positive reference. We use the cutoff-thirteen head tuple comprising its first 46 negative observations of the signed-frequency companion [16]: 547 negative columns E , 1680 positive columns W , and the original polar row with eleven genuine complete-operator null rows. Every column is projected through the full original twelve-row Gram. Let $D_n, D_p \succ 0$ be the actual midpoint frequency metrics plus their rational ridges and put

$$\kappa = 5 \cdot 10^{-19}, \quad B = \kappa I + WD_p^{-1}W^*, \quad C = E^*W, \quad L = W^*W + \kappa D_p.$$

The full frequency quadrature uncertainty is funded within half each ridge, so $B \preceq \kappa I + P_{13}$ in the original metric. Use the exact 1680-by-547 dyadic proposal F in the included head data and set

$$R = C^* - LF, \quad Y = \kappa^{-1}(E - WF), \quad Q_F = \kappa^{-1}(E^*E - CF - F^*C^* + F^*W^*WF + \kappa F^*D_pF).$$

Proposition S6.4 (Full finite-reference covariance and action errors). *For this complete tuple there is a saved full mixed covariance upper $H_R \succeq R^*D_p^{-1}R$ satisfying $H_R \prec 9.36 \cdot 10^{-96}D_n$. In the same coefficient metric the following strict bounds hold:*

$$\begin{aligned} 0 \preceq Q_F - E^*B^{-1}E &\prec 3.745 \cdot 10^{-59}D_n, \\ (Y - B^{-1}E)^*B(Y - B^{-1}E) &\prec 3.745 \cdot 10^{-59}D_n, \\ (BY - E)^*(BY - E) &\prec 1.685 \cdot 10^{-57}D_n, \\ (Y - B^{-1}E)^*(Y - B^{-1}E) &\prec 7.490 \cdot 10^{-41}D_n. \end{aligned}$$

The reference is the stated bounded finite B ; inverse order gives $E^*(P_{13} + \kappa)^{-1}E \preceq E^*B^{-1}E$.

Proof. Woodbury square completion, with the actual rectangular F , gives

$$Q_F - E^*B^{-1}E = \kappa^{-1}R^*L^{-1}R \preceq \kappa^{-2}R^*D_p^{-1}R, \quad BY - E = \kappa^{-1}WD_p^{-1}R.$$

Moreover $W^*B^{-1}W \preceq D_p$. Thus, for $e = Y - B^{-1}E$,

$$e^*Be \preceq \kappa^{-2}R^*D_p^{-1}R, \quad e^*e \preceq \kappa^{-3}R^*D_p^{-1}R.$$

The complete positive-frame trace bounds $\|D_p^{-1/2}W^*WD_p^{-1/2}\| < 44.991$, which pays the action residual separately.

Here is the complete numerical error contract. Reconstruct all 2227 ambient sources and all 2227-by-12 original row crosses, then short by the actual enclosed 12-by-12 row Gram. The source construction and complete error derivation in Section S17, adapted from [16], give a joint error $\delta < 4.129 \cdot 10^{-58}$ in metric $\text{diag}(D_n, D_p)$, including both sine images and the complete kernel, root,

reflected and local errors. Let R_0 be the modeled residual and $H_0 = R_0^* D_p^{-1} R_0$. Applying that joint error to the full coefficient map $[I; -F]$ proves

$$H_R = 2H_0 + 2\delta^2(D_n + F^* D_p F) \succeq R^* D_p^{-1} R.$$

All mixed entries are formed before taking the positive normalized trace. The public saved matrix and actual D_n blocks give, by directed arithmetic,

$$\text{Tr}(D_n^{-1} H_R) < 9.36 \cdot 10^{-96}.$$

Since $H_R \succeq 0$, this proves its full relative upper. The preceding identities, the frame trace and the exact powers of κ prove the table.

The source reconstruction used 768-bit directed arithmetic; its separate sufficient head-cap congruence used 2048 bits. The new source reconstruction and residual assembly took approximately 448 seconds. The reader's saved-array verification reparses all 598418 entries of the two stored 547-square matrices, verifies finite symmetric enclosures and recomputes the actual D_n -normalized residual trace; it is a saved-array replay, rather than another source reconstruction. Its slightly widened durable decimal bounds are those stated above. The public supplement contains the full arrays, true frequency-metric blocks, portable readback and exact algebra controls. Full source regeneration uses the included scientific input subset, analytic source/error supplement and portable producer. $\square$

The distinction between the finite reference and the complete positive observation is retained throughout. A two-sided complete-positive inverse comparison uses the additional source-observation residual of (S19); the finite-reference action estimates above supply their own bounded application of the method.

S6.5 A necessary full-cap margin

The full cap also has a quantitative necessary test independent of null-source conditioning. Let $\mathcal{L}$ be a $(D+1)$ -dimensional genuine operator-null space, and let the exact recipient have codimension at most D in the polar space. Suppose $N[v] \geq b_D \|v\|^2$ on $\mathcal{L}$, with $b_D > 0$. If

$$N_* \preceq (1 - \gamma)(P + \kappa I), \quad 0 \preceq N - N_* \preceq \tau I,$$

on that recipient, and the complete raw tail has norm at most δ , then necessarily

$$\gamma \leq \frac{\kappa + \tau + \delta}{b_D + \kappa + \delta}. \quad (\text{S25})$$

Indeed dimension gives a unit $v \in \mathcal{L}$ in the recipient. The exact null identity gives $P[v] \leq N[v] + \delta$. Together with $N_* \succeq N - \tau I$, the cap gives

$$N[v] - \tau \leq (1 - \gamma)(N[v] + \delta + \kappa),$$

which proves (S25). The lower observation floors developed for genuine derivative nulls in [14] can be inserted without changing this argument. The estimate concerns the entire signed cap, whereas (S24) prices the inverse error on its specified null source.

S7 Complete observation under moving arithmetic cutoffs

The finite-source criteria above admit a complementary reduction on the *entire* negative observation space. Its cutoff is an operator energy, rather than a Fourier window or a finite-rank projection. The useful new input is a complete-positive source estimate uniform in the arithmetic cutoff, the number of source profiles and the number of genuine null rows.

Let X_M be the original odd polar space with any finite subspace K_M of $\ker H$ removed. All projections and norms in this section are original Hilbert ones. Let $P = P_{M,K_M}$ and $N = N_{M,K_M}$ be the restrictions of the complete positive and negative forms to this recipient. Put

$$\begin{aligned} R_M &= 6 + 8(\log 2)\sqrt{M}, & B_M &= 9R_M, \\ J_M &= 3(84R_M + 310 + 2^{27}), & C_M^{\text{src}} &= R_M J_M^2. \end{aligned} \quad (\text{S26})$$

Here $\|s\|_\infty < 3$, $0 \leq n_M \leq R_M$ and $\|N\| \leq B_M$. Section S14 proves these constants from the complete theta series and an elementary prime-power bound.

Theorem S7.1 (Uniform complete-positive source moment). *Let $\phi_1, \dots, \phi_r$ be any finite family in*

$$\mathcal{Y}_M = L^2(\{\xi > 0 : n_M(\xi) > 0\}, n_M(\xi) d\xi/\pi),$$

and let $D \succ 0$ dominate their true joint Gram matrix in this space. Writing Q_M for the original orthogonal projection onto X_M , define

$$g_i(u) = \int_0^\infty n_M(\xi) \phi_i(\xi) \sin(\xi u) \frac{d\xi}{\pi}, \quad E_i = Q_M(sg_i).$$

Then every E_i belongs to $\text{Dom } P$, and

$$E^* E \preceq B_M D, \quad (PE)^*(PE) \preceq C_M^{\text{src}} D. \quad (\text{S27})$$

The constants are independent of r and $\dim K_M$. In particular, for $\kappa, \Lambda > 0$,

$$0 \preceq E^* 1_{(\Lambda, \infty)}(P)(P + \kappa)^{-1} E \preceq \frac{C_M^{\text{src}}}{\Lambda^2(\Lambda + \kappa)} D. \quad (\text{S28})$$

Proof. Section S14 proves, for an odd F -bandlimited g ,

$$\|Q_M(sg)\| \leq 3\|g\|, \quad \|PQ_M(sg)\| \leq 3[51R_M + 33\log(2+F) + 210 + 2^{27} + \delta_M]\|g\|.$$

The complete negative support lies in $[-\Omega_M, \Omega_M]$, with $\Omega_M = e^{R_M+1}$. For every common coefficient vector z , Plancherel gives $\|g_z\|^2 \leq R_M z^* D z$ and $\log(2+\Omega_M) < R_M + 3$. Since $\delta_M < 1$, the displayed graph constant is below J_M . This proves both full matrix inequalities, including all mixed terms. The spectral theorem gives

$$\frac{1_{\lambda > \Lambda}}{\lambda + \kappa} \leq \frac{\lambda^2}{\Lambda^2(\Lambda + \kappa)},$$

which proves (S28). The second moment uses $E_i \in \text{Dom } P$ and does not require $E_i \in \text{Dom } P^2$. $\square$

Define the complete observation $T_M : X_M \rightarrow \mathcal{Y}_M$ by

$$(T_M h)(\xi) = \int_{\mathbb{R}} s(u) h(u) \sin(\xi u) du.$$

Thus $T_M^* \phi = Q_M(sg_\phi)$ and $N = T_M^* T_M$. Let

$$\kappa_M = M^{-1}, \quad \Lambda_M = (4MC_M^{\text{src}})^{1/3}, \quad L_M = T_M 1_{[0, \Lambda_M]}(P)(P + \kappa_M)^{-1} T_M^*. \quad (\text{S29})$$

Theorem S7.2 (Complete-observation energy criterion). *For every $M \geq 2$ and every such finite genuine-null choice,*

$$0 \preceq T_M 1_{(\Lambda_M, \infty)}(P)(P + \kappa_M)^{-1} T_M^* \preceq \frac{1}{4M} I_{\mathcal{Y}_M}. \quad (\text{S30})$$

If on an unbounded sequence of integers M the complete low-energy cap

$$L_M \preceq \left(1 + \frac{3}{4M}\right) I_{\mathcal{Y}_M} \quad (\text{S31})$$

holds, then on the entire original domain

$$q_H[h] \geq - \left[\delta_M + \frac{1}{M} + \frac{B_M}{M+1} \right] \|h\|_2^2. \quad (\text{S32})$$

Conversely, $H \succeq 0$ implies (S31) for every $M \geq 2$ and every finite genuine-null choice. The cofinal energy criterion is therefore equivalent to positivity of this original H and, by Theorem 2.2, to RH . Its original allowance is $O(M^{-1/2})$, and $\Lambda_M = O(M^{5/6})$.

Proof. Theorem S7.1 applies to every finite tuple of profiles. On a dense profile core it bounds both $T_M^* \phi$ and $PT_M^* \phi$ by fixed multiples of $\|\phi\|_{\mathcal{Y}_M}$. For a convergent profile sequence, both images are Cauchy. Closedness of P gives graph-norm completion to every $\phi \in \mathcal{Y}_M$, with the same bounds. Thus (S28) extends to the entire observation space. The choice of Λ_M gives (S30); its high and low pieces are the exact functional-calculus decomposition of the same resolvent.

Adding (S31) gives $T_M(P + \kappa_M)^{-1} T_M^* \preceq c_M I$, $c_M = 1 + 1/M$. The equivalent form inequality is $N \preceq c_M(P + \kappa_M I)$. Rearrange using the bounded N , rather than introducing an adverse multiple of unbounded P :

$$P - N \succeq -\kappa_M I - (1 - c_M^{-1})N \succeq - \left[\frac{1}{M} + \frac{B_M}{M+1} \right] I.$$

Add the complete original-norm raw tail. Exact genuine-null transport from Theorem 2.2 proves (S32) on all original V . Its allowance tends to zero.

For the converse, $H = P - N + W_M \succeq 0$ gives $N \preceq P + \delta_M I$. The bound (8) satisfies $\delta_M < 1/M$ for all $M \geq 2$. A rational admission at $M = 2$ is

$$2 \cdot 451 \cdot 9(10/27)^9(10/13) < 1,$$

using $\pi > 31/10$, $e > 27/10$ and $e^{3/10} > 13/10$. The consecutive majorant ratio is at most $(8/3)e^{-\pi} < (8/3)(3/8)^3 < 1$. Hence the complete observation cap is at most I , and so is its low-energy summand. $\square$

The spectral space $1_{[0, \Lambda_M]}(P)X_M$ may be infinite dimensional. Theorem S7.2 specifies a complete operator comparison; the finite-source Schur criterion and source-residual enclosures of Section S6 supply separate constructive interfaces. The energy scale does not replace the Fourier support Ω_M or the rank of a chosen positive receiving frame.

Corollary S7.3 (A finite cutoff interval from the M16 certificate). *If a complete original-domain certificate gives $H \succeq -aI$ with $a = 5.289 \cdot 10^{-19}$, as in the signed-frequency companion [16], then for every integer $2 \leq M \leq 10^{18}$ and every finite genuine-null choice,*

$$T_M(P + M^{-1}I)^{-1} T_M^* \preceq I, \quad L_M \preceq I.$$

Proof. At every arithmetic cutoff the same original bound gives $N_M \preceq P_M + (a + \delta_M)I$. It suffices that $M(a + \delta_M) < 1$. For $2 \leq M \leq 15$, the decreasing majorant used above gives $M\delta_M < 819/1000$, while $Ma < 1/1000$. For $M \geq 16$,

$$M\delta_M \leq 451 \cdot 16 \cdot 17^2(3/8)^{51} < 1/4, \quad Ma \leq 5289/10000 \quad (M \leq 10^{18}).$$

These rational bounds prove the assertion. The bridge is the complete original-domain inequality, rather than an identification of the M16 finite matrices with matrices at another cutoff. $\square$

This finite-interval consequence and the unbounded-sequence hypothesis in Theorem S7.2 have different quantifiers. The former follows from one fixed global allowance; the latter is the complete low-energy comparison that would make the allowance vanish.

S7.1 True observation compression and a paid omitted trace

The complete energy criterion has a finite observation-space interface. It retains the true source columns in its finite head and pays the entire remaining covariance, including mixed blocks, by positivity. Keep $\mathcal{Y}_M, T_M, P, \kappa = M^{-1}$ and $\Omega = e^{R_M+1}$ from Section S7, and write $E = T_M^*$ and $\nu = n_M(\xi) d\xi/\pi$ on $[0, \Omega]$. Put

$$W = R_M\Omega/\pi, \quad a_* = 31/32, \quad t(L) = 28e^{-a_*r_L}(r_L/a_* + a_*^{-2}), \quad r_L = \pi e^{2L}.$$

The complete two-arm bound in Section S14 gives

$$\nu([0, \Omega]) \leq W, \quad \|s\|_2^2 \leq 28/a_*^2 < 32, \quad \int_{|u|>L} s(u)^2 du \leq t(L).$$

The last integral follows by substituting $r = \pi e^{2|u|}$ in $s^2 \leq 28r^2e^{-a_*r}$; both arms together have Jacobian dr/r .

Theorem S7.4 (True finite head and complete observation return). *Choose $L \geq 1$ with $Wt(L) \leq 1/(8M^2)$ and an integer $J \geq 1$ such that*

$$N_J = 2J + 1 \geq 2e\Omega L, \quad N_J \geq \log_2(16M\sqrt{W}). \quad (\text{S33})$$

Let Π be the true orthogonal projection in $\mathcal{Y}_M$ onto $\text{span}\{\xi, \xi^3, \dots, \xi^{2J-1}\}$, and let $U : \mathbb{C}^J \rightarrow \mathcal{Y}_M$ be an isometry onto that span. Then

$$\|E(I - \Pi)\|_{\text{HS}}^2 \leq \frac{1}{4M^2}. \quad (\text{S34})$$

For $R_{\text{low}} = 1_{[0, \Lambda_M]}(P)(P + \kappa)^{-1}$ define the true finite matrix $G_M = (EU)^ R_{\text{low}}(EU)$. The finite-head condition*

$$G_M \preceq \left(1 + \frac{1}{2M}\right) I_J \quad (\text{S35})$$

implies the complete low-energy cap (S31). Its validity along an unbounded sequence of admitted cutoffs and finite genuine-null complements is equivalent to original H positivity, with allowance (S32). Sufficient choices have $J = O(\Omega \log M) = \exp(O(\sqrt{M}))$.

Proof. The ambient sine-source kernel is Hilbert–Schmidt, with square at most $W\|s\|_2^2$. Applying the original physical orthogonal projections preserves this property. Replace its sine on $|u| \leq L$

by $p_J(\xi u) = \sum_{j=0}^{J-1} (-1)^j (\xi u)^{2j+1} / (2j+1)!$ and set the approximant to zero outside. The resulting source map E_J has rank at most J and

$$\|E - E_J\|_{\text{HS}}^2 \leq W \left[t(L) + 32 \left\{ \frac{(\Omega L)^{N_J}}{N_J!} \right\}^2 \right]. \quad (\text{S36})$$

Indeed p_J is also the sine Taylor polynomial through degree $2J$, so its real remainder is bounded by $|\xi u|^{N_J} / N_J!$. Orthogonal polar and genuine-null projections contract the whole error. The observation dependence of E_J lies in $\text{ran } \Pi$; thus $E_J(I - \Pi) = 0$. The elementary bound $N! \geq (N/e)^N$ and (S33) make the remainder at most $2^{-N_J} \leq 1/(16M\sqrt{W})$. Equation (S36) is then at most $1/(4M^2)$, proving (S34).

For any positive $0 \preceq R \preceq \kappa^{-1}I$, the operator $\mathcal{L} = E^*RE$ is positive trace class. Positivity bounds its mixed quadratic term between $x \in \text{ran } \Pi$ and $y \in \ker \Pi$ by the geometric mean of its diagonal terms. Scalar Cauchy–Schwarz consequently gives

$$\|\mathcal{L}\| \leq \|\Pi \mathcal{L} \Pi\| + \text{tr}((I - \Pi)\mathcal{L}(I - \Pi)) \leq \|\Pi \mathcal{L} \Pi\| + \kappa^{-1} \|E(I - \Pi)\|_{\text{HS}}^2. \quad (\text{S37})$$

This pays all mixed blocks without declaring them zero. Use $R = R_{\text{low}}$: the omitted trace is at most $1/(4M)$, and (S35) gives $\|\mathcal{L}\| \leq 1 + 3/(4M)$. Theorem S7.2 now returns the original allowance. Conversely $H \succeq 0$ gives complete observation cap at most I , so every such true compression satisfies (S35).

The choices terminate because $t(L) \rightarrow 0$. Since $\log W = O(\sqrt{M})$, a polynomially growing r_L suffices, giving $L = O(\log M)$; the two requirements on N_J then give the asserted sufficient rank. The monomial Gram is positive definite because $m_M(0) < 0$ and hence ν is positive on an interval near zero. This proves existence of U ; numerical evaluation must certify its actual observation metric. $\square$

The finite matrix uses EU , not the Taylor-surrogate E_JU . This true compression is what permits a quadratic omitted-error payment. The rank bound concerns the observation compression and makes no rank assertion about $1_{[0, \Lambda_M]}(P)$. For upper certification one may instead bound the larger full-resolvent head $V^*(P + \kappa)^{-1}V$, $V = EU$. If trial columns Y lie in $\text{Dom } P$ and $\mathcal{A} = V - (P + \kappa)Y$ is their *complete* action residual, then

$$\|V^*(P + \kappa)^{-1}V - \text{Herm}(V^*Y)\| \leq \kappa^{-1} \|V\| \|\mathcal{A}\|.$$

This follows from $(P + \kappa)^{-1}V - Y = (P + \kappa)^{-1}\mathcal{A}$. It supplies a paid upper in the true source coordinates. A residual $V - BY$ for a different finite reference still leaves $(P + \kappa - B)Y$ to be enclosed; all sources, norms and operators in the complete head estimate must match.

S8 The hard-cutoff mean and intrinsic receiver rank

The original hard-cutoff receivers have a useful independent description. On the polar space set

$$\mathcal{C}_0 = \frac{4}{\pi} Q_b M_s \beta(D) M_s Q_b, \quad \beta(\xi) = (1 + 4\xi^2)^{-1/2}, \quad \mathcal{R} = (\ker H)^\perp.$$

The two references $\mathcal{C}_0$ and $\mathcal{C}_Q$ have Fourier orders one and two, respectively. The next estimates concern the hard cutoff. For the right-continuous prime-power sum $\Psi_\Lambda(x) = \sum_{n \leq x} \Lambda(n)$ write $E_\Lambda(x) = \Psi_\Lambda(x) - x$ and

$$e_\Lambda(M) = \frac{|E_\Lambda(M)|}{M} + \frac{1}{\sqrt{M}} \int_1^M |E_\Lambda(x)| x^{-3/2} dx.$$

Theorem S8.1 (Complete common mean and one-sided rate). *The operators $N_M/\sqrt{M}$ converge strongly to $\mathcal{C}_0$. On $\text{Dom } H$, $P_M h/\sqrt{M} \rightarrow \mathcal{C}_0 h$, and the selfadjoint positive operators have the same strong-resolvent limit. For $F \geq 1$, $0 < \eta < 1/2$ and $M^\eta \geq \sqrt{(1+2\eta)/(1-2\eta)}$,*

$$\frac{N_M}{\sqrt{M}} \succeq \mathcal{C}_0 - \rho_M I, \quad (\text{S38})$$

$$\rho_M = 18(1+F)e_\Lambda(M) + \frac{9(12 + \log(3F))}{\sqrt{M}} + \frac{36}{\sqrt{1+4F^2}} + \frac{30M^{-\eta}}{1-2\eta}.$$

The same one-sided estimate compresses to every finite row complement. Classical PNT permits $F \rightarrow \infty$ with $\rho_M \rightarrow 0$.

Proof. Put $z = 1/2 + i\xi$, $U = \log M$. Abel summation, with its lower endpoint retained, gives

$$\begin{aligned} \frac{1}{\sqrt{M}} \sum_{n \leq M} \Lambda(n) n^{-1/2+i\xi} &= \frac{e^{i\xi U}}{z} + \frac{z-1}{z\sqrt{M}} + e^{i\xi U} \frac{E_\Lambda(M)}{M} \\ &\quad - \frac{z-1}{\sqrt{M}} \int_1^M E_\Lambda(x) x^{-3/2+i\xi} dx. \end{aligned}$$

As $|z-1| = |z|$, the difference from $e^{i\xi U}/z$ on $|\xi| \leq F$ is at most $M^{-1/2} + (1+F)e_\Lambda(M)$. The positive-part map is 1-Lipschitz. Using $|m_0| < 6$ and $a(\xi) \leq 4 + \log(3F)$, the normalized negative symbol differs on this interval by at most $d = 2(1+F)e_\Lambda(M) + (12 + \log(3F))/\sqrt{M}$ from

$$v_M(\xi) = 4\beta(\xi)(\cos(\xi U - \arctan(2\xi)))_+.$$

The absolutely convergent Fourier series of $\cos_+$ has constant coefficient $1/\pi$ and nonconstant complex coefficient sum $1/2 + 1/\pi$. Riemann–Lebesgue gives the local weak multiplier mean $4\beta/\pi$. The normalized symbols are uniformly bounded by (S26); their local weak means therefore extend against every L^1 frequency test. For fixed input, truncate its Fourier profile $\widehat{sh}$ to $[-F, F]$. Multiplication by s after a finite frequency projection is Hilbert–Schmidt, so it converts the weak convergence on that band to strong convergence. The input exterior tends to zero uniformly in M . This proves the negative strong limit.

For a quantitative continuum bound, the positive l th harmonic continues analytically on $|\Im z| < 1/2$ as

$$e^{iUz}(1-2iz)^{(l-1)/2}(1+2iz)^{-(l+1)/2}.$$

Shift to $\Im z = \eta$ on compact smooth physical tests. The multiplier bound is $(1-4\eta^2)^{-1/2}q^l$, $q = M^{-\eta}\sqrt{(1+2\eta)/(1-2\eta)} \leq 1$. The two physical factors acquire opposite exponential weights and $\|e^{\eta u}s\|_\infty \|e^{-\eta u}s\|_\infty < 9$. Indeed the complete envelope bounds their squares by $169x^{2+\eta}e^{-\pi x} \leq 169e^{-\pi} < 9$, $x = e^{2|u|} \geq 1$. Vertical contour ends vanish by smooth compact integration by parts; density extends the resulting bounded operators to all inputs. Summing the harmonics costs at most $30M^{-\eta}/(1-2\eta)$. Discard the actual positive negative-symbol exterior beyond F and pay the omitted continuum by $36/\sqrt{1+4F^2}$. The retained symbol error costs $9d$. This proves (S38).

Finally $P_M = N_M + H - W_M$ on $\text{Dom } H$ gives the positive limit there. The resolvent identity at $(\mathcal{C}_0 + i)h$ and the uniform resolvent norm bound extend it by density to strong resolvents. PNT makes $e_\Lambda(M) \rightarrow 0$ by splitting its integral at a fixed large threshold; for example take $F = \min(M^{1/4}, e_\Lambda(M)^{-1/2})$. $\square$

Theorem S8.2 (Intrinsic hard-cutoff receiver rank). *There is $c_* > 0$ such that*

$$\lambda_j(P_{\mathcal{R}}\mathcal{C}_0|_{\mathcal{R}}) \geq c_*/j, \quad j \geq 1.$$

Consequently, if a rank- J positive receiver B_M on the complement of any finite genuine-null space K_M satisfies $N_{M,K_M} \preceq B_M + \kappa_M I$, then

$$J + 1 \geq \frac{c_*}{\rho_M + \kappa_M / \sqrt{M}}. \quad (\text{S39})$$

With a classical PNT remainder one may choose $\rho_M = O(e^{-c\sqrt{\log M}})$. Along $\kappa_M \rightarrow 0$, the required receiver rank is therefore at least $c_1 e^{c_2 \sqrt{\log M}}$ for sufficiently large M .

Proof. The operator $\mathcal{C}_0$ is compact: its finite-frequency sandwiches are Hilbert–Schmidt, and the omitted multiplier has norm at most $(36/\pi) \sup_{|\xi| > F} \beta(\xi) \rightarrow 0$. Section S15 proves the eigenvalue floor using the original-domain critical-zero representers and complete Schwartz-weight Gram estimates. On its $(J + 1)$ -dimensional protected span choose a unit vector in $\ker B_M$. It belongs to every genuine-null complement. Combining the floor with (S38) gives (S39). The classical error $|E_\Lambda(x)| \leq A x e^{-c\sqrt{\log x}}$ gives $e_\Lambda(M) = O(e^{-c_0 \sqrt{\log M}})$ by splitting at $\sqrt{M}$. Choose $F = e^{c_0 \sqrt{\log M}/2}$ and fixed $\eta = 1/4$. $\square$

For arbitrary finite, possibly nongenuine row spaces of dimension d , an elementary version is also useful. If $\sigma_R = \min_{|u| \leq R} s^2$ and $a_R = 4\sigma_R / [\pi \sqrt{1 + 4\pi^2/R^2}]$, then $\lambda_j(\mathcal{C}_0) \geq a_R/(j + 1)$ and $J + d + 2 \geq a_R/(\rho_M + \kappa_M/\sqrt{M})$. Take the first $j + 1$ odd Dirichlet sine modes f on $[-R, R]$, set $h = f/s$ and impose the single polar moment. Fourier Jensen gives

$$\langle f, (1 + 4D^2)^{-1/2} f \rangle \geq \frac{\|f\|^2}{\sqrt{1 + 4\|f'\|^2/\|f\|^2}},$$

and $\|f'\| \leq \pi(j + 1)\|f\|/R$. Minmax and codimension interlacing give the displayed assertions. These are representation costs for the hard-cutoff split; the smoothed reference of Section S3 has its own source geometry.

S9 The full residual covariance and the signed return

The finite receiving gate must retain the source covariance between the phase space, the exact rows and the complementary carriers. The following exact Schur identity gives the interface. It is classical finite-rank elimination specialized to the original constrained recipient [18, 7].

Theorem S9.1 (Residual-covariance signed gate). *Let $Z : \mathbb{C}^d \rightarrow \mathcal{H}$ have finite-dimensional range containing the exact row map $\mathcal{R}$. Let P be its range projection and $Q = I - P$. For a finite carrier map U , define*

$$G = Z^* Z, \quad F = U^* Z, \quad J = U^* Q U.$$

Let $C = C^$ and $a > \|J\|$. Then*

$$aI - ZCZ^* - UU^* \succeq 0 \quad \text{on } \ker \mathcal{R}^* \quad (\text{S40})$$

if and only if

$$aG - GCG - aF^*(aI - J)^{-1}F \succeq 0 \quad \text{on } \{c : \mathcal{R}^* Zc = 0\}. \quad (\text{S41})$$

The columns of Z may be dependent. If $J \preceq \bar{J} \prec aI$, substitution of $\bar{J}$ gives a sufficient gate. For any finite matrix V of trial coefficients,

$$J \preceq (U - ZV)^*(U - ZV). \quad (\text{S42})$$

Proof. The Q block of (S40) is $aQ - QUU^*Q \succ 0$. Its exact elimination adds

$$PUU^*P + PUU^*Q(aQ - QUU^*Q)^{-1}QUU^*P = aPU(aI - U^*QU)^{-1}U^*P$$

to the adverse finite block. Every $x \in P\mathcal{H}$ has the form Zc ; evaluation gives (S41). Since $\text{ran } \mathcal{R} \subseteq \text{ran } Z$, the entire Q space satisfies the exact rows, so this elimination holds on the stated kernel. Inverse operator monotonicity proves the upper-covariance substitution. Finally $Q(U - ZV) = QU$, and the orthogonal P, Q split of the trial residual proves (S42). $\square$

For injective Z , $J = U^*U - FG^{-1}F^*$. Formula (S42) supplies a complete positive upper without subtracting nearly equal numerical matrices or assuming a projected row Gram floor. It keeps the full residual covariance rather than replacing it by unrelated diagonal carrier bounds.

S9.1 Paid common-source errors

Theorem S12.1 supplies the joint source enclosure, its exact-row shorting and the full congruence payment. The numerical application in Section S13 retains every phase/row/carrier block.

A positive padded covariance $\Gamma \succeq \delta \text{diag}(D_p, D_r)$ has shorted phase block at least δD_p : minimize its quadratic form over the row coefficients. This supplies a lower bound for the padded elimination without requiring a lower bound on the angle between unpadded phase and row spans. The padding is charged in the signed inequality. An original full-row Gram floor applies to its stated receiving map; a complementary projection requires its own analysis or this paid residual construction.

S9.2 Return to the original form

For a finite arithmetic head and $x = \sqrt{\alpha + s^2/5}h$, the exact identity is

$$q_{H_M}[h] + \alpha \|h\|_2^2 = \|x\|_2^2 - \int_{\mathbb{R}} (1/5 - m_M(\xi)) |\widehat{t_\alpha x}(\xi)|^2 \frac{d\xi}{2\pi}. \quad (\text{S43})$$

The original polar row becomes $t_\alpha \sinh(u/2)$ in this variable. To separate the favorable unbounded part, fix a finite Λ and write $d_M = 1/5 - m_M$ as

$$d_M = d_{\text{ret}} + b - g, \quad d_{\text{ret}} = 1_{|\xi| \leq \Lambda} d_M, \quad b = 1_{|\xi| > \Lambda} (d_M)_+, \quad g = 1_{|\xi| > \Lambda} (-d_M)_+ \geq 0.$$

The original graph domain pays each integral. Dropping the favorable $-g$ gives a form upper bound; only d_{ret} and b receive bounded-operator approximations. Suppose these supply the complete upper domination

$$M_{t_\alpha}(1/5 - m_M)(D)M_{t_\alpha} \preceq ZCZ^* + UU^* + \eta I, \quad \eta < 1,$$

as forms on the transformed original core. For example, a finite signed source approximation, a positive defect approximation and complete norm errors supply this domination after retaining the favorable exterior by its sign. Set $a = 1 - \eta$ and apply Theorem S9.1 on the exact transformed rows. A successful gate gives

$$q_{H_M}[h] \geq -\alpha \|h\|_2^2, \quad q_H[h] \geq -(\alpha + \delta_M) \|h\|_2^2,$$

where (7) supplies a full raw arithmetic payment. Finite genuine-null deflation is returned as in Theorem 2.2. All identities begin on the exact polar core and extend through the original closure; bounded retained perturbations preserve its domain.

Thus, for $\alpha_j = 2^{-j}$ and any cofinal $M_j \rightarrow \infty$, complete signed gates returning (S43) yield the vanishing allowances required by Theorem 2.2. Theorem S10.1 supplies finite receiving spaces,

complete positive complements and certified continuous coordinates for every such index. The exact residual gate specifies the signed comparison to be certified in those coordinates.

The coordinated schedule of Theorem S10.3 supplies the explicit original allowance $(1+2^{-37})2^{-j}$. The direct return adds the raw norm error to α_j ; any stronger inverse-normalized error requirement used to establish a chosen finite gate is paid within that gate.

S10 A complete conditioned construction

The receiving space will approximate every frequency source at once. Put

$$\alpha_j = 2^{-j}, \quad t_j = \frac{s}{\sqrt{\alpha_j + s^2/5}}, \quad j \geq 16.$$

Let ν_j be a positive finite measure supported in $[0, \Omega_j]$, with mass A_j , and define the positive operator on $\mathcal{H} = L^2_{\text{odd}}(\mathbb{R})$ by

$$O_j = \frac{1}{\pi} \int |t_j \sin(\xi \cdot)\rangle \langle t_j \sin(\xi \cdot)| d\nu_j(\xi). \quad (\text{S44})$$

For an even nonnegative multiplier b , take $d\nu = b(\xi) d\xi$ on the positive half-line. The identity $\cos(x-y) - \cos(x+y) = 2 \sin x \sin y$ gives exactly (S44) in the Fourier convention of Section 2. Positive atoms are also allowed.

Theorem S10.1 (Conditioned complete complements). *Choose a computable integer $B_j \geq \max(1, A_j)$ and put*

$$\begin{aligned} L_j &= j + \log(1 + B_j), \\ T_j &= j \log 2 + \log(1 + B_j) + 4 \log L_j + 20, \quad R_j = \tfrac{1}{2} \log(T_j/\pi), \\ E_j &= 12\Omega_j R_j + 2 \log(1 + B_j) + 4 \log L_j + 40. \end{aligned}$$

Choose an even integer N_j with $E_j \leq N_j < E_j + 4$. Let V_j be t_j times the odd polynomials of degree less than N_j , restricted to $[-R_j, R_j]$, and let Q_j project onto its entire odd orthogonal complement. Then

$$\|Q_j O_j Q_j\| < \frac{1}{2 \log L_j}, \quad \dim V_j \leq 6\Omega_j R_j + \log(1 + B_j) + 2 \log L_j + 22. \quad (\text{S45})$$

For computable supplied parameters, a terminating certified construction gives a dyadic basis transformation whose actual continuous receiving Gram lies between $3I/5$ and $5I/3$. Its condition number is at most $25/9 < 3$.

Proof. The elementary inequalities

$$L/2 < T < 3L, \quad 0 < R < \tfrac{1}{2} \log L$$

hold for $L \geq 16$. For the upper bound use $j \log 2 + \log(1 + B) \leq L$ and $4 \log L + 20 < 2L$; the latter ratio decreases from its strict endpoint value at 16. For $0 \leq \xi \leq \Omega$, real Taylor approximation supplies an odd polynomial p_ξ of degree less than N with

$$|\sin(\xi u) - p_\xi(u)| \leq \frac{(\Omega R)^N}{N!} < 4^{-N} \quad (|u| \leq R),$$

since $N \geq 12\Omega R$ and $e < 3$. Thus the squared interior error after multiplication by t , where $t^2 < 5$, is at most $10R16^{-N}$. The complete theta upper bound (6), with $x = e^{2|u|}$, gives the full exterior estimate

$$\begin{aligned} \int_{|u|>R} t^2 du &\leq 169 2^j \int_{T/\pi}^{\infty} x e^{-\pi x} dx \\ &= \frac{169 e^{-20}}{(1+B)L^4} \frac{T+1}{\pi^2}. \end{aligned} \quad (\text{S46})$$

Orthogonal projection is at least as accurate as this explicit approximation. Integrating the squared error against the whole positive source gives

$$\|QOQ\| \leq \text{Tr}(QOQ) \leq \frac{A}{\pi} \left(10R16^{-N} + \frac{169 e^{-20}(T+1)}{\pi^2(1+B)L^4} \right). \quad (\text{S47})$$

This trace includes every covariance combination and both infinite spatial tails. Because $N \geq 2\log(1+B) + 4\log L + 40$ and $\log 16 > 2$, $16^{-N} < e^{-80}(1+B)^{-4}L^{-8}$. Multiplying the first term in (S47) by $\log L$ bounds it by $(5/3)e^{-80}(\log L)^2 L^{-8} < 1/4$. The second term times $\log L$ is at most $(676/27)e^{-20}L^{-2} < 1/4$, using $A/(1+B) \leq 1$, $T+1 < 4L$ and $\pi > 3$. This proves (S45). Upward rounding an outward enclosure of E of width less than one to the next even integer gives the required N without a transcendental equality decision.

For conditioning, the first theta atom and the decrease of $r^2 e^{-r}$ for $r > 2$ give, throughout $[-R, R]$,

$$\frac{s^2}{\alpha_j} \geq c_j = \frac{2T^2 e^{-20}}{(1+B)L^4}, \quad \frac{e^{-20}}{2(1+B)L^2} < c_j < 1.$$

Consequently

$$t_j^2 \geq \frac{c_j}{1+c_j/5} > \frac{e^{-20}}{4(1+B)L^2} > \ell_j := \frac{2^{-32}}{(1+B)L^2}. \quad (\text{S48})$$

The final comparison follows from $30\log 2 > 20$. In an ordinary $L^2([-R, R])$ -orthonormal odd Legendre basis the true receiving Gram satisfies $\ell_j I \preceq G_j \preceq 5I$.

Choose a positive rational $\ell'_j \leq \ell_j$ by outward evaluation. Compute a symmetric rational moment enclosure $\tilde{G}_j$ with $\|G_j - \tilde{G}_j\| \leq \ell'_j/4$; entrywise error $\ell'_j/(4\dim V_j)$ suffices. On this compact interval the complete theta series has effective uniform derivative tails and its denominator is positive. Certified partitions and increasing precision therefore terminate. The moments here are physical receiving moments, determined by s, t_j, R_j ; computability of ν_j is needed only when evaluating the source operator or its mass rather than using its supplied bounds.

Since $\tilde{G}_j \succeq 3\ell'_j I/4$, increasingly accurate inverse-Cholesky proposals give a dyadic invertible W_j passing the finite check

$$\frac{3}{4}I \preceq W_j^* \tilde{G}_j W_j \preceq \frac{5}{4}I.$$

The error bound and $G_j \succeq \ell'_j I$ imply $(3/4)G_j \preceq \tilde{G}_j \preceq (5/4)G_j$. Hence

$$\frac{3}{5}I \preceq W_j^* G_j W_j \preceq \frac{5}{3}I.$$

Only the basis changes; the space, its entire complement and the trace bound are unchanged. $\square$

The condition estimate concerns the polynomial phase Gram on V_j . Adjoined exact rows enter the joint signed gate with their actual mixed Gram, as specified in Section S9.

For fixed A, Ω , a fixed B yields $\dim V_j = O(\log j)$ and a complementary norm tending to zero. The next section proves necessity of this order for a nonzero fixed density. Polynomial approximation, finite-band concentration and whitening are classical mechanisms [2, 9, 8]; the explicit theta-dependent wall, whole-tail payment, funded Gram floor and original signed interface specify their use here.

S10.1 Moving arithmetic heads

The construction also applies when the complete arithmetic head grows. Put

$$m_M(\xi) = \Re \Psi(1/4 + i\xi/2) - \log \pi - 2 \sum_{2 \leq n \leq M} \frac{\Lambda(n)}{\sqrt{n}} \cos(\xi \log n), \quad b_M = (1/5 - m_M)_+.$$

Every prime power is included. The following elementary bounds, also used in the signed-frequency companion [16], give concrete inputs.

Lemma S10.2. *For $M \geq 2$, the positive-half support and mass of b_M are bounded by*

$$\Omega_M = e^{7+8(\log 2)\sqrt{M}}, \quad A_M \leq \Omega_M[7 + 8(\log 2)\sqrt{M}].$$

The coordinated choice below turns these inputs into explicit original-domain tail and receiving-space budgets.

Proof. The Chebyshev and Abel argument in Section 3 gives $S_M \leq 4(\log 2)\sqrt{M} - 2\log 2$ by (10). The complete digamma comparison (11) and $m_0 > -6$ give

$$m_M(\xi) > -6 + \log(2|\xi|) - 2S_M.$$

Thus outside Ω_M , $m_M > 1 + \log 2 > 1/5$. Everywhere the defect is bounded by $7 + 2S_M < 7 + 8(\log 2)\sqrt{M}$. Integrating this bound over its positive-half support proves the asserted mass. $\square$

Theorem S10.3 (Coordinated arithmetic and receiving budgets). *For $j \geq 16$, put $\kappa_j = 2^{-j}$ and choose an integer M_j with*

$$E_j \leq M_j < E_j + 2, \quad E_j = \frac{j \log 2 + 3 \log(j+2) + 40}{\pi}. \quad (\text{S49})$$

Then the complete original arithmetic remainder satisfies

$$\delta_{M_j} < 2^{-37} \kappa_j. \quad (\text{S50})$$

For the moving positive source b_{M_j} , Theorem S10.1 gives condition number at most 25/9 and

$$\log \dim V_j \leq 8 \log 2 \sqrt{\frac{\log 2}{\pi}} \sqrt{j} + O(\log j). \quad (\text{S51})$$

The leading coefficient is 2.60467320417428.... If complete signed head gates return $q_{H_{M_j}}[h] \geq -\kappa_j \|h\|_2^2$ on the original recipient, then

$$q_H[h] \geq -(1 + 2^{-37})2^{-j} \|h\|_2^2. \quad (\text{S52})$$

Consequently such a cofinal family satisfies Theorem 2.2.

Proof. An outward enclosure of E_j of width less than one, followed by upward integer rounding, supplies the stated integer. For $j \geq 16$, $\log(j+2) \leq j/2$, $\log 2 < 1$ and $\pi > 3$ imply

$$M_j + 1 < E_j + 3 < \frac{5j+98}{6} \leq 3(j+2).$$

Insert this into (8) and use $M_j \geq E_j$:

$$\delta_{M_j} < \frac{4059e^{-40}}{j+2} 2^{-j} \leq \frac{4059}{18} \left(\frac{3}{8}\right)^{40} 2^{-j} < 2^{-37} 2^{-j}.$$

Here $e > 8/3$; the last comparison is rational. Moreover $M_j = (\log 2/\pi)j + O(\log j)$. Choose B_j by upward enclosure of the mass bound in Lemma S10.2. Then $\log(1+B_j) = 8(\log 2)\sqrt{M_j} + O(\log M_j)$, $L_j = j + O(\sqrt{j})$, and $R_j = O(\log j)$. Substitution into (S45) proves (S51). Finally the bounded raw remainder costs at most $\delta_{M_j}\|h\|_2^2$, proving (S52). Finite genuine-null deflations return by Theorem 2.2. $\square$

The constructed rank in (S51) prices the positive receiving space at the final original-norm scale. With $\epsilon_j = (1 + 2^{-37})2^{-j}$, the same estimate reads

$$\log \dim V_j \leq \frac{8 \log 2}{\sqrt{\pi}} \sqrt{\log(1/\epsilon_j)} + O(\log \log(1/\epsilon_j)).$$

The arithmetic and positive-complement estimates hold at every index. The signed hypothesis in Theorem S10.3 is the finite gate of Section S9, including its complete approximation and residual-covariance payments.

S11 The necessary fixed-source dimension

This lower bound counts the whole retained odd span, including any adjoined receiving directions.

Theorem S11.1 (Sharp dimension order). *Let $b \geq 0$ be a nonzero bounded density on positive frequencies supported in a fixed $[a, \Omega]$ with $a > 0$, extended evenly as a Fourier multiplier. Fix one integer $B \geq \max(1, A)$ and use the corresponding R_j of Theorem S10.1. Let $O_j = M_{t_j}b(D)M_{t_j}$ on the odd space. Set*

$$A = \int_0^\infty b, \quad B_2 = \int_0^\infty b^2, \quad B_\infty = \|b\|_\infty.$$

For any finite spaces W_j with $\|Q_{W_j}O_jQ_{W_j}\| \leq \tau_j \rightarrow 0$,

$$\liminf_{j \rightarrow \infty} \frac{\dim W_j}{R_j - 1} \geq \frac{B_2}{\pi B_\infty^2} > 0.$$

Together with Theorem S10.1, the optimal dimension order for obtaining a complementary norm tending to zero is $\Theta(\log j)$; the constructed rate is the one in (S45).

Proof. Take $L = R_j - 1$. On $|u| \leq L$, the first-atom lower bound gives

$$s^2/\alpha_j \geq 2\pi^2 \exp(j \log 2 - T_j/e^2) \rightarrow \infty.$$

Thus $t_j^2 \geq 4$ there for all sufficiently large j . Compare the full source covariance with the source covariance obtained by replacing t_j by $1_{[-L, L]}$. The former is at least four times the latter in frequency-source space. By equality of the nonzero spectra of U^*U and UU^* , and min-max,

$$\lambda_n(O_j) \geq 4\lambda_n(T_L), \quad T_L = 1_{[-L, L]}b(D)1_{[-L, L]} \quad (\text{S53})$$

on odd functions. This comparison is made in the common source space.

Put $C_L(\omega) = 2 \sin(L\omega)/\omega$, continuously defined at zero. The complete sine-carrier formulas give

$$\begin{aligned}\mathrm{Tr} T_L &= \frac{1}{2\pi} \int_0^\infty b(\xi)[2L - C_L(2\xi)] d\xi, \\ \mathrm{Tr} T_L^2 &= \frac{1}{4\pi^2} \int_0^\infty \int_0^\infty b(\xi)b(\eta)[C_L(\xi - \eta) - C_L(\xi + \eta)]^2 d\xi d\eta.\end{aligned}$$

Consequently $\mathrm{Tr} T_L/L \rightarrow A/\pi$. The measures $C_L(\omega)^2 d\omega/(4\pi L)$ are a positive approximate identity of mass one. Applied to the continuous autocorrelation of the zero-extended b , this gives the difference-term limit B_2/π in $\mathrm{Tr} T_L^2/L$. Since $\xi + \eta \geq 2a$, its square term is $O(1)$ and the mixed term is $O(\sqrt{L})$ by Cauchy–Schwarz. Therefore

$$\mathrm{Tr} T_L^2/L \rightarrow B_2/\pi, \quad \|T_L\| \leq B_\infty.$$

If $N_j = \dim W_j$, min–max and (S53) give $\lambda_{N_j+1}(T_L) \leq \tau_j/4$. Hence

$$\mathrm{Tr} T_L^2 \leq N_j B_\infty^2 + (\tau_j/4) \mathrm{Tr} T_L.$$

Divide by L and take the lower limit. Finally $R_j \sim \frac{1}{2} \log j$ for fixed B . $\square$

The first-two-trace argument belongs to classical time–frequency concentration. Its present use matches the explicit complete-theta upper construction, including arbitrary competing odd receiving spans. Compression by the fixed polar row changes a dimension count by at most one; additional growing constraint dimensions are counted with their actual span.

S12 Complete joint sources and stable row elimination

The phase construction controls a positive complement. To evaluate its signed recipient, collect the actual phase, exact-row and carrier profiles in one source map. Put $q = s^2$, $w_\alpha = q/(\alpha + q/5) = t_\alpha^2$. On a finite interval I , write

$$Yc = \sqrt{w_\alpha} \sum_a c_a f_a, \quad \Gamma = Y^*Y.$$

Supported phase profiles are polynomials on their support cells, the polar profile is $\sinh(u/2)$, and a derivative-row profile is

$$f_k = (\psi^{(k+2)} - \psi^{(k)})/4/q \quad (k \text{ odd}).$$

Literal carriers have profile $\sqrt{\nu_a/\pi} \sin(\xi_a u)$. An exact column transformation is applied to the entire derivative polynomial before evaluation. The original preimages determine genuine-null status; the transformed rows are receiving coordinates.

Theorem S12.1 (Common-weight full covariance). *Let the finitely many measurable profiles belong to $L^2(I, w_\alpha)$, and let a real positive function p satisfy $|p - w_\alpha| \leq \eta w_\alpha$ on I , where $0 < \eta < 1$. Define $\Gamma_p = \int_I f(u)^* f(u) p(u) du$. Then*

$$(1 - \eta)\Gamma \preceq \Gamma_p \preceq (1 + \eta)\Gamma. \tag{S54}$$

For a partition into retained and row coefficients, define the shorted form $S[x] = \inf_z (x, z)^ \Gamma(x, z)$, and similarly S_p . Then*

$$(1 - \eta)S \preceq S_p \preceq (1 + \eta)S.$$

These assertions allow dependent columns and redundant rows.

Let $\hat{Y}_p$ approximate $Y_p = \sqrt{p}f$ and let $D \succ 0$ be a specified coefficient metric. If

$$\|(Y_p - \hat{Y}_p)D^{-1/2}\| \leq e, \quad \|\hat{Y}_p D^{-1/2}\| \leq M,$$

then, with $\delta = 2Me + e^2$,

$$-\delta D \preceq \Gamma_p - \hat{Y}_p^* \hat{Y}_p \preceq \delta D, \quad \Gamma \preceq \frac{\hat{Y}_p^* \hat{Y}_p + \delta D}{1 - \eta}. \quad (\text{S55})$$

Shorting this entire upper gives an upper for S .

Proof. Integrate the pointwise weight inequalities against $|\sum c_a f_a|^2$. Taking the infimum over the same row coefficients proves the shorted inequalities. For the profile change put $E = Y_p - \hat{Y}_p$ and expand $Y_p^* Y_p$. The two cross terms have total D -normalized norm at most $2Me$, and $E^* E$ at most e^2 . Combining with the left inequality of (S54) proves (S55). Quadratic-form order is preserved by the defining infimum. $\square$

This is a source-specific use of classical positive integration and variational Schur order [18, 8]. In particular, the relative-weight step preserves the exact kernel. A profile approximation is priced separately by (S55). No lower angle between the phase and row spans enters these bounds. For any exact coefficient map W , an additive enclosure $-\delta D \preceq \Gamma - \hat{\Gamma} \preceq \delta D$ becomes the enclosure $\delta W^* DW$ after congruence. This is the payment used for enrichment or row changes.

S12.1 Positive polynomial integration

Here is a fully specified realization of the weight step. Suppose w_α is analytic on each closed disk of radius d centered in $[-R, R]$, with $|w_\alpha| \leq H$, and $w_\alpha \geq w_0 > 0$ on the real interval. These are checkable source hypotheses. Split the interval into cells of half-width $h \leq d/8$, also splitting at every supported-profile endpoint. On a cell centered at c , let $L \geq m \geq 1$ and $\zeta_\ell = e^{2\pi i \ell / L}$. Use

$$a_k = \left(\frac{2h}{d}\right)^k \frac{1}{L} \sum_{\ell=0}^{L-1} w_\alpha(c + (d/2)\zeta_\ell) \zeta_\ell^{-k}, \quad p(c + hx) = \sum_{k < m} a_k x^k.$$

Exact conjugate-paired sampling makes p real. Cauchy's coefficient bound and root-of-unity aliasing give

$$\|w_\alpha - p\|_\infty \leq \frac{H(h/d)^m}{1 - h/d} + \frac{H 2^{-L}}{(1 - 2^{-L})(1 - h/d)}. \quad (\text{S56})$$

Indeed the discrete coefficient at index $k < L$ contains only Taylor indices $k + vL$; summing their bounds proves the second term. Literal sample and real dyadic coefficient errors add their full uniform payment. Choosing their sum with (S56) below ηw_0 supplies the positive approximant in Theorem S12.1.

For polynomial profile approximants of degree below N , a positive Gauss–Legendre rule of order at least $N + m$ on each cell integrates $p \hat{f}^* \hat{f}$ exactly: every entry has degree below $2N + m$. Certified node, weight and evaluation errors are included when the rule is represented by literal numbers. Thus its matrix is a single positive common-factor sum. The rule uses

$$O((R/d + 1)(N + m))$$

points on a uniform grid, plus the support-boundary subdivisions. This argument establishes a constructive rule under its stated strip and profile hypotheses.

S12.2 Complete growing-row tails

A full-line covariance also needs physical tails. Write a premixed row profile on $u \geq 0$ as

$$f_a(u) = c_a \sinh(u/2) + Q_a(u)/q(u), \quad Q_a(u) = e^{u/2} \sum_{n \geq 1} P_a(\pi n^2 e^{2u}) e^{-\pi n^2 e^{2u}}.$$

Use odd reflection on the other arm. Let $D \geq 2$ bound all polynomial degrees, K_a bound each polynomial's absolute coefficient sum, and $F_0 = 1 + \sum_a (|c_a|^2 + K_a^2)$.

Proposition S12.2 (Premixed complete row tails). *For $\alpha = 2^{-j}$ and $T = \pi e^{2R} \geq \max(\pi, 4D)$,*

$$\sum_a \int_{|u| > R} w_a |f_a|^2 du \leq 676 2^j F_0 T^{2D} e^{-T}. \quad (\text{S57})$$

For $0 < \varepsilon < 1$, a sufficient wall for this trace to be below ε^2 is

$$T \geq 4D, \quad T \geq 4\{j \log 2 + \log(676 F_0) + 2D \log(8D) + 2 \log(1/\varepsilon) + 10\}.$$

Proof. Put $r = \pi e^{2u}$. For $r \geq 4D$, comparison of successive terms gives $|\sum_n P_a(r n^2) e^{-r n^2}| \leq 2K_a r^D e^{-r}$. For example $n^{2D} e^{-r(n^2-1)}$ is decreasing with a geometrically bounded ratio, using $\log n \leq (n^2 - 1)/2$. The first positive theta atom gives $q \geq 2r^2 e^{-r}$, while (6) gives $q \leq 169 e^{4u-r}$ and $w_a \leq 2^j q$. Apply $|x+y|^2 \leq 2|x|^2 + 2|y|^2$, $\sinh^2(u/2) \leq e^u$, $du = dr/(2r)$ and both arms. Enlarging powers to r^{2D} and constants bounds the trace by $338 2^j F_0 \int_T^\infty r^{2D} e^{-r} dr$. Since $T \geq 4D$, integration by parts or a geometric series gives an upper $2T^{2D} e^{-T}$. Finally $2D \log T \leq T/4 + 2D \log(8D)$ proves the sufficient wall. $\square$

In particular $D = O(j/\log(j+2))$, $\log F_0 = O(j)$ and $\log(1/\varepsilon) = O(j)$ permit $T = O(j)$ and $R = O(\log j)$. If a common strip has $d^{-1} = O(j)$, logarithmic profile bounds $O(j)$, and $\log(w_0^{-1}) = O(j)$, the construction above reaches exponential accuracy with $O(j)$ scaled coefficient precision. For the raw identity metric this also requires logarithmic column count $O(j)$; a different metric additionally pays its inverse norm. High-frequency sine profiles use degree $O(|\xi|h+j)$ on each cell. These statements specify source precision and node counts under the displayed hypotheses.

Complete mixed tails follow from the full source factor: if the row and carrier tail traces are E_r, E_c , their joint tail trace is $E_r + E_c$ and the cross block has norm at most $\sqrt{E_r E_c}$. Supported phase tails vanish outside their support. The full covariance therefore remains a single error-controlled input to Theorem S9.1.

S13 A complete 4473-column source certificate

We give a whole-line application of the common-source construction at $\alpha = 3/10^{12}$. The supplied data specify exact dyadic row coordinates and literal frequency carriers. The polynomial-weight construction of Section S12 is one general realization; this application uses a directly enclosed positive Gaussian rule for the actual theta sources.

Theorem S13.1 (Full mixed physical Gram). *Let Y have the 4080 supported phase columns, 41 complete receiving rows and 352 literal carriers specified below and in the scientific data. Let $\hat{\Gamma}$ be the exact 2^{-100} -dyadic block-arrow matrix in `data/joint_centers.npz`. Then*

$$\|Y^* Y - \hat{\Gamma}\| < 1.384607318 \cdot 10^{-20} < 2^{-65}.$$

The norm is the raw Euclidean coefficient operator norm. All phase/row, phase/carrier and row/carrier terms, the entire global block, quadrature errors and both infinite physical tails are included. In particular, putting $\delta = 2^{-65}$,

$$Y^* Y \preceq \hat{\Gamma} + \delta I, \quad \hat{\Gamma} + 2\delta I \succeq \delta I.$$

S13.1 Exact columns and a portable construction

The negative half-line is split into 100 cells of half-width $r = 1/80$, centered at $c_i = (-199 + 2i)/80$, $0 \leq i < 100$. Odd reflection supplies the positive half-line. On the last 68 negative cells ($32 \leq i < 100$), take the 60 local columns

$$t_\alpha(u) \sqrt{\frac{2k+1}{2r}} P_k((u - c_i)/r), \quad 0 \leq k < 60,$$

where P_k is the Legendre polynomial; outside the paired cell supports they vanish. Their unweighted odd Gram is $2I_{4080}$. Disjoint cell interiors give exact zero off-cell blocks.

The 41 global rows are an exact dyadic triangular mixture of the polar profile and the profiles with odd derivative indices $1, 3, \dots, 79$. The file `row_coordinates.json` gives the literal coefficients and polynomial normalizations. To reconstruct them, start with

$$P_0(x) = 4x^2 - 6x, \quad P_{\ell+1}(x) = (1/2 - 2x)P_\ell(x) + 2xP'_\ell(x).$$

Then $\psi^{(\ell)}(u) = e^{u/2} \sum_n P_\ell(\pi n^2 e^{2u}) e^{-\pi n^2 e^{2u}}$ on $u \geq 0$. Normalize $P_{k+2} - P_k/4$ by its stated largest absolute coefficient and premix the exact polynomials before evaluation. The file `literal_rule.json` gives 352 rational frequencies $2400 < \xi_a < 6000$ and positive dyadic weights ν_a with $\sum_a \nu_a < 2$. Their columns are exactly $t_\alpha \sqrt{\nu_a/\pi} \sin(\xi_a u)$.

The script `generate_rows.py` evaluates all 80,000 negative-arm Gaussian nodes at 512-bit ball precision using 16 theta atoms and a complete polynomial-dependent remainder. It produces double-double source arrays and certified point errors. The global assembly uses those source arrays, computes carrier phases by directed ball arithmetic, and pays all point and fixed-point errors jointly. The exact row coordinates are mathematical inputs; no numerical QR is part of the certificate.

S13.2 Whole quadrature and tail payments

The following explicit bounds specify the analytic contract implemented by `row_budget.py` and `joint_gram.py`. They also permit reproduction without a sampled Gram hypothesis. On each Gaussian cell use the Bernstein ellipse $\rho = 7/5$. Its imaginary height is

$$h = \frac{r}{2}(\rho - \rho^{-1}) < 1/200.$$

The actual theta denominator is bounded in this strip by first-atom comparison on $|\Re u| \leq 7/5$ and by regularization outside. For $v = \pi e^{2z}$ the first scaled atom satisfies $|P_0(v)| \geq 2|v|^2$. The other atoms have relative sum at most $3 \sum_{n \geq 2} n^4 e^{-(n^2-1)\Re v} < 1/10$ for $\Re v > 2.99$, giving the lower denominator modulus $9X_-^2/5$ below. Inner-strip phase comparison gives $\Re \psi > 0$; the outer envelope and $\Re \cosh(z/2) > .99$ give receiving denominator real part above $\alpha/2$. The complete strip envelope for ψ is below 100, which also funds the displayed bound $4 \cdot 10^{19}$ for t_α^2 . The finite checks are

$$13/400 + \pi e^{14/5}/100 < 1, \quad 120e^{63/10 - \pi e^{14/5} \cos(1/100)} < \alpha/4.$$

Use even reflection $z \mapsto -z$ for a negative-real portion of the first positive-arm ellipse; the weight is even and the receiving rows are odd. For a positive-arm cell let $x_- = \max(0, c - r(\rho + \rho^{-1})/2)$ and $x_+ = c + r(\rho + \rho^{-1})/2$ be its extreme real coordinates on that ellipse, $X_\pm = \pi e^{2x_\pm}$, and $c_h = \cos(2h)$. A valid majorant is

$$t_2 = \min\{4 \cdot 10^{19}, (240/\alpha)e^{9x_+/2 - X_- c_h}\}, \quad d_0^{-1} = 5/(9X_-^2).$$

For a premixed derivative polynomial P_a , let A_a be its absolute-coefficient polynomial. Its complete atom numerator is bounded by

$$B_a = \sum_{n=1}^{16} A_a(n^2 X_+) e^{-(n^2-1)X-c_h} + 2A_a(289X_+) e^{-288X-c_h}.$$

The script verifies that the remaining atom ratio is below $1/2$. The corresponding row bound is $\sqrt{t_2} e^{x_+/2} (|S_{0a}| + d_0^{-1} B_a)$. The local Legendre factor is bounded by $70\rho^{59}$ and the carrier sine factor by e^{6000h} . Positive Gaussian integration of an analytic scalar product has error at most

$$\frac{16r\rho^{-1600}}{1-\rho^{-1}} \times (\text{its ellipse supremum}).$$

Trace bounds for row blocks and Frobenius bounds for mixed blocks give a complete summed quadrature error below $5.6 \cdot 10^{-185}$.

For the whole physical tail put $Z = \pi e^5$, and define

$$I_0 = e^{-Z}, \quad I_k = kI_{k-1} + Z^k e^{-Z} = \int_Z^\infty x^k e^{-x} dx.$$

If $A_a(x)^2 = \sum b_{a,k} x^k$, the complete row-tail trace is bounded by

$$E_r = \sum_a \left\{ \frac{4}{\alpha} \sum_{k \geq 2} b_{a,k} I_{k-2} + \frac{240}{\alpha} |S_{0a}|^2 I_2 \right\} < 1.384607235 \cdot 10^{-20}.$$

The full exterior atom sum is bounded by twice its first absolute-polynomial term; the explicit ratio tests at Z in the script verify this for every included degree. The carrier tail has bound

$$E_c = \frac{\sum \nu_a}{\pi} \frac{240}{\alpha} (e^5/3 + 1/9) e^{-3e^5}.$$

The row/carrier mixed tail is at most $\sqrt{E_r E_c}$. Supported phase columns vanish outside their own wall. Adding $E_r + E_c + 2\sqrt{E_r E_c}$ is therefore a conservative full-block tail payment.

Proof of Theorem S13.1. Use the common 800-point positive Gauss–Legendre rule on each of the 100 cells. Complete theta source intervals pay the point uncertainty. Directed 256-bit ball products compute the sampled matrix; an alternative supplied route quantizes the common factors to 2^{-120} and uses exact integer products. Directed products retain the double-double uncertainties in their full matrix intervals. In the fixed-point alternative, the common-factor inequality (S55) pays source uncertainty and factor quantization together. For this factor payment let the phase/global errors be e_p, e_g , and let the represented global trace be T_g . The phase factor norm is at most $\sqrt{10}$ before perturbation, since $t_\alpha^2 < 5$. Thus

$$e = (e_p^2 + e_g^2)^{1/2}, \quad M \leq \{(\sqrt{10} + e_p)^2 + T_g\}^{1/2}, \quad E_{\text{pt}} \leq 2Me + e^2.$$

Center the exact products at 2^{-100} . The block-arrow center-rounding error is at most

$$E_{\text{ctr}} \leq \max(E_p, E_g) + E_{pg},$$

where E_p, E_g are maximal directed row sums of the phase/global center radii and E_{pg} is the Frobenius norm of all mixed radii. This uses the block-diagonal phase norm, the global block norm, and the complete rectangular cross-block Frobenius norm; disjoint-support zeros remain exact. For exact fixed-point products the simpler bound $\{\max(60, 393) + \sqrt{4080 \cdot 393}\} 2^{-100}$ suffices. The portable assembly verifies $E_{\text{pt}} + E_{\text{ctr}} < 8.3 \cdot 10^{-28}$. Add the whole quadrature and tail bounds above to obtain the stated radius. The final two inequalities follow from positive semidefiniteness of Y^*Y and its two-sided enclosure. $\square$

All uncertainty in this theorem concerns the stated literal carriers. A continuous frequency measure is supplied to the original signed return with its own operator-approximation debit. The raw covariance error transports through an enrichment W as δW^*W . The scientific supplement contains the exact columns, complete source arrays, centers and portable generators, together with a short verification command and controls that reject underfunded error caps.

S14 Complete source bounds with uniform null cost

This section proves the actual-source estimates used in Section S7. The Fourier convention, theta weight, original polar row and minimal realization are those of Section 2. No auxiliary positive reference replaces $P_{M,K}$.

The shared arithmetic, logarithmic and complete-theta estimates used here are proved once in Section 3. In this section P_M again denotes the hard-cutoff positive form.

S14.1 Original polar return and count-independent null return

On the ambient odd space let $\bar{P}_M = M_s p_M(D) M_s$ and $h_0 = sg$. Bandlimited L^2 functions have every derivative in L^2 ; complete theta multiplication and smooth compact cutoffs put h_0 in the minimal form domain. Equation (15) and form representation then put it in the ambient operator domain and give

$$\|\bar{P}_M h_0\| \leq 3[33(R_M + \log(2 + F)) + 210]\|g\|. \quad (\text{S58})$$

For example the compact approximants converge in L^2 and their weighted physical derivatives converge in L^2 ; $a \leq 17\xi^2$ pays convergence in the specified minimal form norm.

The original odd polar row is $b = s \sinh(u/2)$. Put $f_b = sb = s^2 \sinh(u/2)$. From (13), its exponentially weighted Fourier norm is at most $32\sqrt{2}$. Since $\sup_{x \geq 0} x^2 e^{-x/8} = 256/e^2 < 36$,

$$\|f_b''\| < 1728, \quad \|\bar{A}b\| < 88128.$$

On each interval $1/4 \leq |u| \leq 1/2$, the first complete theta atom gives $s^2 > 18 \cdot 3^{-10}$ and $\sinh^2(u/2) \geq 1/64$. Consequently

$$\|b\|^2 > \frac{9}{64 \cdot 3^{10}} > 2^{-20}, \quad \frac{\|\bar{A}b\|}{\|b\|} < 2^{27}. \quad (\text{S59})$$

Here $r = \pi e^{2|u|} < \pi e < 10$, $r > 3$ and $e < 3$ suffice for the lower bound. The same compact form approximation and representation justify b 's operator membership. From (12), $\bar{P}_M - \bar{A}$ is bounded by B_M . Writing Q_b for the polar projection and $x = Q_b h_0$, therefore,

$$\|P_M x\| \leq \|\bar{P}_M h_0\| + (2^{27} + B_M)\|h_0\|.$$

The constrained action is $P_M x = Q_b \bar{P}_M x$; preservation of the polar condition by the ambient action is not assumed.

Now let T be an original Hilbert isometry onto any finite-dimensional $K \subseteq \ker H$. Bounded perturbation in the same minimal realization gives $\text{Dom } H = \text{Dom } P_M$ and

$$P_M T = N_M T - W_M T, \quad \|P_M T\| \leq B_M + \delta_M.$$

For $e = Q_K x$, the operator on the further constrained form domain acts as $P_{M,K} e = Q_K P_M e$, whence

$$\|P_{M,K} e\| \leq \|P_M x\| + (B_M + \delta_M)\|x\|.$$

Combining the last displays, $\|x\| \leq \|h_0\| \leq 3\|g\|$ and (S58) proves the promised count-independent bound

$$\|P_{M,K}Q_KQ_b(sg)\| \leq J(M,F)\|g\|, \quad J(M,F) = 3[51R_M + 33\log(2+F) + 210 + 2^{27} + \delta_M]. \quad (\text{S60})$$

This argument uses actual operator-nullity. An arbitrary additional row requires its own graph cost. Independence of $\dim K$ is an analytic norm estimate and does not assert that numerical projection is free.

For arbitrary negative profiles and a common complex coefficient vector,

$$\|g_z\|^2 = \int_0^\infty n_M^2 |\phi_z|^2 \frac{d\xi}{\pi} \leq R_M z^* D z.$$

Their support is bounded by Ω_M , and $J(M, \Omega_M) < J_M$. Equation (S60) proves the complete joint moment in Theorem S7.1. Applying the same inequalities to profile differences and using closedness of $P_{M,K}$ proves the full observation-space graph completion used in Theorem S7.2.

The same graph estimate also pays finite inverse-action trials. If E_0, W_0 are actual negative and positive source families, the latter supported below F_0 , and D_n, D_p dominate their true frequency metrics, put $V = \kappa^{-1}Q_M(E_0 - W_0 F_c)$ and $F_b = \max(\Omega_M, F_0)$. Before multiplication by s , the two profiles have disjoint Fourier sign supports. Thus

$$(PV)^*(PV) \preceq \kappa^{-2}J(M, F_b)^2[R_M D_n + (R_M + \log(2 + F_0))F_c^* D_p F_c].$$

This supports complete source-action residual accounting. Physical mixed covariance after multiplication by s remains present; disjoint raw frequency sign supports do not make the receiving cross block vanish.

S15 Protected critical representers and their spectral floor

We supply the original-domain and uniform Gram arguments used in Theorem S8.2. The complete polarized formula is

$$q_H(h_1, h_2) = \sum_z m_z \mathcal{M}(sh_2)(z) \overline{\mathcal{M}(sh_1)(-\bar{z})}, \quad \mathcal{M}f(z) = \int f(u) e^{zu} du.$$

All centered nontrivial zeros occur with multiplicity.

Lemma S15.1 (Critical evaluation vectors in the operator range). *For every positive critical ordinate γ , $v_\gamma = Q_b(s \sin(\gamma u))$ belongs to $\text{ran } H$.*

Proof. Put $\sigma = i\gamma$ of exact multiplicity m and $P_0 = d^2/du^2 - 1/4$. For a function whose first r transform jets vanish at σ , define

$$R_\sigma^r f(u) = \int_u^\infty \frac{(t-u)^{r-1}}{(r-1)!} e^{\sigma(t-u)} f(t) dt.$$

Its canceled lower-tail expression is $(-1)^r \int_{-\infty}^u (u-t)^{r-1} e^{\sigma(t-u)} f(t) dt / (r-1)!$. The vanishing jets prove equality, and direct integration gives $\mathcal{M}(R_\sigma^r f)(z) = \mathcal{M}f(z) / (z-\sigma)^r$. With $\eta = R_\sigma^m \psi$, the detector $h_\sigma = P_0 \eta / s$ therefore has transform $(z^2 - 1/4)\xi(1/2 + z) / (z-\sigma)^m$. It evaluates only at σ , where its value A is nonzero.

Here is its original minimal-domain return. Each fixed-order theta derivative is bounded on either arm by a polynomial in $r = \pi e^{2|u|}$ times $e^{|u|/2-r}$; the first atom gives the matching positive

lower for s^2 . In the upper primitive use $t = u + v$ and $e^{2v} \geq 1 + 2v$; its complete tail is bounded by a polynomial times $e^{u/2-r}e^{-rv}v^{m-1}$ beyond a fixed onset. Use the canceled lower primitive on the other arm. Differentiation gives the corresponding fixed-order estimates. For compact even cutoffs, $P_0(\chi R\eta)$ has both pole moments zero exactly, converges after division by s in L^2 , and its derivative converges in ordinary L^2 . The bound $a \leq 17\xi^2$ gives convergence in V . Weighted integration by parts gives arbitrary vertical inverse powers, so the $O(T \log T)$ zero count passes the full zero formula to the limit, as in Appendix A.

Oddify by $h_o(u) = h_\sigma(u) - h_\sigma(-u)$. Its only evaluations are A and $-A$ at $\pm i\gamma$. The polarized formula gives, in the conjugate-linear-first convention, $q_H(h_o, h) = 2mi\bar{A}\langle v_\gamma, h \rangle$. This is a bounded Hilbert functional; form representation puts $h_o \in \text{Dom } H$ and $Hh_o = -2miAv_\gamma$. In particular $v_\gamma \perp \ker H$. $\square$

Classical positive critical-line proportion [5] and the local total zero bound $N(T+1) - N(T) = O(\log(T+3))$ give a sequence of distinct critical ordinates with any fixed large separation G and $\gamma_j \leq Dj$. To see this, greedily retain a point and discard the following interval of length G . Each discarded cluster has total multiplicity $O_G(\log T)$ below height T , whereas the critical-line count is at least $cT \log T$. Thus the number retained below T is at least $c_G T$, even when multiplicities are charged to clusters. Discard any fixed initial interval and enlarge D if necessary.

For an even positive Schwartz weight F , its sine Gram is

$$\frac{1}{2}[\hat{F}(\gamma_i - \gamma_j) - \hat{F}(\gamma_i + \gamma_j)].$$

Fourier decay bounds the off-diagonal difference row sums by $C_N \sum_{l \neq 0} (1 + G|l|)^{-N}$; the sum-frequency row sums tend to zero with the first retained height. Schur's test therefore gives uniform upper and lower Riesz bounds, simultaneously for $F = s^2$ and s^4 . The polar coefficients $c_j = \langle b, s \sin(\gamma_j u) \rangle / \|b\|^2$ have an arbitrarily small *whole* ℓ^2 norm after increasing that height. Hence for $h = \sum_{i \leq j} a_i v_{\gamma_i}$ and $f = sh$,

$$l_h \|a\|^2 \leq \|h\|^2 \leq u_h \|a\|^2, \quad \|f\|^2 \geq l_f \|a\|^2.$$

Differentiating $f = \sum a_i s^2 \sin(\gamma_i u) - sb \sum a_i c_i$ and applying the upper cosine/sine Gram bounds with weights s^4 and $|(s^2)'|^2$ gives $\|f'\|^2 \leq C\gamma_j^2 \|a\|^2$ uniformly in j . Fourier Jensen, as displayed after Theorem S8.2, now gives $\langle \mathcal{C}_0 h, h \rangle \geq c_*/j \|h\|^2$ on this j -dimensional span. Lemma S15.1 puts it in $\mathcal{R}$. Compact minmax proves the required floor. Complete theta decay supplies all Schwartz seminorms used here; no constant depends on the number of subsequently removed genuine nulls.

S16 Origin compression and fixed-window separation

This appendix records an additional constraint on the original hard-cutoff origin layer. Put $U = \log M$, $0 < L < \pi/2$ and fix $B < \infty$, $C \geq 0$. On the original polar-only space let N_M^{org} use the negative window $[0, L/U]$, and let $P_M^{[B]}$ use the actual positive window $[0, B/U]$.

Proposition S16.1 (A bounded scaled positive window). *There are compact smooth original polar tests h_M and constants $c, c' > 0$, with an integer k fixed before M , such that eventually*

$$(N_M^{\text{org}} - CP_M^{[B]})(h_M) \geq c\sqrt{M}/U, \quad \|h_M\|^2 \leq c'U^{4k+2}.$$

Thus this fixed positive-window budget has an adverse Rayleigh depth at least a constant times $\sqrt{M}/U^{4k+3}$.

Proof. PNT and Abel summation give uniformly on every fixed scaled interval $m_M(t/U)/\sqrt{M} \rightarrow -4\cos t$, with positive/negative parts converging to $4(-\cos t)_+$ and $4(\cos t)_+$. Choose $b > \max(B, L, \pi/2)$ and $p_k(t) = t(1 - t^2/b^2)^k$. The positive integral on $[0, B]$ is exponentially small in k , since its support starts at $\pi/2$. The negative integral on $[0, L]$ is at least a positive multiple of k^{-3} by restricting to $[0, 1/(2k)]$. Fix k so the negative integral exceeds C times the positive one by a strict amount.

On a fixed compact interval the odd functions $u, u^3, \dots, u^{2k+1}, \sinh(u/2)$ are independent. A positive even bump and their Gram inverse supply compact smooth odd moment duals. Writing $p_k(t) = \sum_{j \leq k} a_j t^{2j+1}$, choose f_M with exact polar moment zero and $\int f_M u^{2j+1} = (-1)^j (2j+1)! U^{2j+1} a_j$. Then $h_M = f_M/s$ is in the original compact polar core, $\|h_M\| = O(U^{2k+1})$, and its sine transform at t/U is $p_k(t) + O(U^{-2})$ uniformly on the two retained intervals. Substitute in the actual observations and use their scaled limits. The fixed strict polynomial gap gives the asserted inequalities. $\square$

There is also a positive finite-frequency compression for any original projection Q . With $Th(\xi) = \langle Q(s \sin(\xi u)), h \rangle$ in the true origin measure, its monomial projection Π_d gives $0 \preceq T^* \Pi_d T \preceq T^* T$. This is a true source projection and keeps its full monomial Gram. A sine Taylor approximation and the complete theta moments give the bound used in the portable controls:

$$\|T^* T - T^* \Pi_d T\| \leq 64 \beta_M \frac{[\Omega \log(4m)/2]^{2m}}{(m!)^2}, \quad m = 2d + 1, \quad \beta_M = \int_0^\Omega n_M d\xi/\pi.$$

For completeness put $r_m = \log(4m)/2$. For $x = |u| \leq r_m$, $x^{2m} \leq r_m^{2m}$. For $x \geq r_m \geq \log 2$, monotonicity of $x e^{-2x}$ gives

$$2m \log(x/r_m) \leq 2m e^{-2r_m} e^{2x} = \frac{1}{2} e^{2x} < \frac{\pi}{2} e^{2x}.$$

Thus $x^{2m} \leq r_m^{2m} e^{r/2}$ everywhere, and both real arms give

$$\|su^m\|^2 \leq 16 r_m^{2m} \int_\pi^\infty r e^{-r/2} dr = 16(2\pi + 4) e^{-\pi/2} r_m^{2m} < 64 r_m^{2m}.$$

The real Taylor remainder is $|\xi u|^m/m!$; the squared observation-map error integrates to the displayed bound. The exact projection has no larger error. At $\Omega = L/\log M$, for any fixed $c > (p+1/2)/8$, taking $d = \lceil c \log M / \log \log M \rceil$ pays M^{-p} eventually, by Stirling. This is a compression estimate in the true observation metric, complementary to the complete ridged comparison of Theorem S4.1.

S17 Source and error contracts for the finite-reference application

This section supplies the source construction and analytic error payments used in Section S6.4, in the original norm and minimal form domain of Section 2. It adapts the author's signed-frequency construction [16]; that citation records the origin of the construction, while the necessary arguments are included here. The hard-cutoff symbols in this section are

$$m_M = m_0 + a - 2 \sum_{2 \leq n \leq M} \Lambda(n) n^{-1/2} \cos(\xi \log n), \quad p_M = (m_M)_+, \quad n_M = (-m_M)_+.$$

In the numerical head $M = 13$. Let $Y = L_{\text{odd}}^2(\mathbb{R})$, $b = s \sinh(u/2)$ and $X = \mathcal{K} = b^\perp \cap Y$. The weight, Fourier measure and physical norm are exactly those of (3)–(4). The original realization

and its bounded paired hard-cutoff return were established in Lemma 3.1; the genuine operator nulls

$$z_{2j+1} = (\psi^{(2j+3)} - \psi^{(2j+1)}/4)/s$$

are the vectors proved legal and null in Lemma G.1. Their Mellin normalization and the polar cancellation are proved in Appendix A. Thus these domain and null facts require no additional predecessor hypothesis. Only the finite scientific inputs, including literal dyadic coefficients and row/source moments, are stored as data; their analytic construction and error bounds are given below.

S17.1 Finite receiving with positive-frequency references

For a nonnegative weight w supported in a bounded frequency set put

$$(T_w h)(\xi) = \sqrt{w(\xi)} \int_{\mathbb{R}} s(u) h(u) \sin(\xi u) du \quad \text{in } L^2(\mathbb{R}_+, d\xi/\pi).$$

For the numerical construction this map acts on the ambient odd Hilbert space $Y = L^2_{\text{odd}}(\mathbb{R})$; its inequalities can then be compressed to X . In particular $N_M = T_{n_M}^* T_{n_M}$ on Y . On frequency intervals take $z_k = \mathbf{1}_I \sqrt{w} T_k(t_I)$, where $t_I : I \rightarrow [-1, 1]$ is affine. For their column map Z , define the raw metric $D = Z^* Z$ and $E = T_w^* Z$.

Lemma S17.1 (Frequency range projection). *For positive-weight columns with metric D_+ , ridge $\rho_+ > 0$ and $W = T_{p_M}^* Z_+$,*

$$P_{\text{fin}} = W(D_+ + \rho_+ I)^{-1} W^* \leq P_M.$$

Such constructions add to at most P_M when their frequency supports have disjoint interiors.

Proof. Singular values give $0 \leq Z(Z^* Z + \rho I)^{-1} Z^* \leq I$. Conjugate by the observation map. For disjoint sets their exact observation energies add, and each finite projection is below its own energy. $\square$

Absorb both ridges into $D_n, D_p > 0$. Set

$$N_\star = E D_n^{-1} E^*, \quad P_{\text{fin}} = W D_p^{-1} W^*, \quad G_{nn} = E^* E, \quad C = E^* W, \quad G_{pp} = W^* W.$$

These are full ambient mixed covariances. An ambient operator inequality restricts exactly to X . The head uses the complete exact-row compression proved below.

Theorem S17.2 (Full mixed variational certificate). *Let $\kappa > 0$, $L = G_{pp} + \kappa D_p$, and fix a matrix F of size $(\dim D_p) \times (\dim D_n)$. The condition*

$$\kappa D_n - G_{nn} + C F + F^* C^* - F^* L F \geq 0 \tag{S61}$$

implies $N_\star \leq P_{\text{fin}} + \kappa I$. If an approximate full Gram satisfies

$$-\delta \text{diag}(D_n, D_p) \leq G - \tilde{G} \leq \delta \text{diag}(D_n, D_p), \tag{S62}$$

it suffices to check

$$\begin{aligned} \kappa D_n - \tilde{G}_{nn} + \tilde{C} F + F^* \tilde{C}^* - F^* (\tilde{G}_{pp} + \kappa D_p) F \\ - \delta (D_n + F^* D_p F) \geq 0. \end{aligned} \tag{S63}$$

Proof. The positive inverse identity gives

$$\kappa E^*(\kappa I + W D_p^{-1} W^*)^{-1} E = G_{nn} - C L^{-1} C^*.$$

Completing the square,

$$C L^{-1} C^* - (C F + F^* C^* - F^* L F) = (L^{-1} C^* - F)^* L (L^{-1} C^* - F) \geq 0.$$

Thus (S61) implies $E^*(\kappa I + P_{\text{fin}})^{-1} E \leq D_n$. Singular values of $(\kappa I + P_{\text{fin}})^{-1/2} E D_n^{-1/2}$ give the desired operator inequality. The Gram-dependent part is $-(I, -F^*) G (I, -F^*)^*$. Apply (S62) to $(v, -Fv)$, giving precisely $\delta v^*(D_n + F^* D_p F)v$. This proves (S63). $\square$

A dyadic F is an exact witness. A numerical inverse can propose it; the proof uses its literal entries and the enclosed sign. The approximate Gram need not be positive. All mixed covariance remains in the sign test. If an additional exact-row compression is used in another construction, its full joint correction must also be retained.

An alternative is a direct certified inverse calculation. A frozen approximate inverse of L is validated by its complete Neumann residual, which encloses L^{-1} ; a directed Schur sign then proves $\kappa D_n - G_{nn} + C L^{-1} C^* \geq 0$. This is the same inverse criterion in the proof above, without exporting an explicit variational F . The supplement specifies which route each receiving group uses.

S17.1.1 Complete exact-row covariance

Let $\mathcal{Z} : \mathbb{C}^r \rightarrow Y$ be an injective row map with $\Gamma = \mathcal{Z}^* \mathcal{Z} > 0$, and set

$$Q = I - \mathcal{Z} \Gamma^{-1} \mathcal{Z}^*, \quad \mathcal{C}_n = \mathcal{Z}^* E, \quad \mathcal{C}_p = \mathcal{Z}^* W, \quad \mathcal{C} = [\mathcal{C}_n, \mathcal{C}_p].$$

The complete physical source Gram on $\text{ran } Q$ is exactly

$$G_Q = G - \mathcal{C}^* \Gamma^{-1} \mathcal{C}. \tag{S64}$$

Projection changes the physical columns; the true frequency metrics D_n, D_p retain their definitions.

Theorem S17.3 (Conservative row-compressed receiving). *Suppose $\bar{\Gamma} \geq \Gamma > 0$, and the ambient model obeys (S62). For a fixed F , put*

$$T_F = \begin{pmatrix} I \\ -F \end{pmatrix}, \quad R_F = \mathcal{C}_n - \mathcal{C}_p F.$$

The condition

$$\kappa(D_n - F^* D_p F) - T_F^* \tilde{G} T_F + R_F^* \bar{\Gamma}^{-1} R_F - \delta(D_n + F^* D_p F) \geq 0 \tag{S65}$$

implies

$$Q N_* Q \leq Q P_{\text{fin}} Q + \kappa Q \quad \text{on } \text{ran } Q.$$

Proof. Inverse order gives $\bar{\Gamma}^{-1} \leq \Gamma^{-1}$ and hence

$$G_Q \leq G - \mathcal{C}^* \bar{\Gamma}^{-1} \mathcal{C}.$$

Thus the left side of (S65) is a lower bound for $\kappa(D_n - F^* D_p F) - T_F^* G_Q T_F$. The ambient model error is paid on the whole vector $T_F v$ by (S62). Apply Theorem S17.2 to $Q E, Q W$ in $\text{ran } Q$. $\square$

All entries of $\mathcal{C}_n, \mathcal{C}_p$ and the proof of $\bar{\Gamma} \geq \Gamma$ belong to this certificate. Enclosing arithmetic evaluates their complete expression (S65); the row source is not replaced by selected entries or by the identity Gram. Direct-kernel locality does not remove its dense correction.

S17.2 Degree-independent locality and actual source funding

Let $q = s^2$ and $\hat{q}(\omega) = \int_{\mathbb{R}} q(u) e^{-i\omega u} du$. The ambient odd feature kernel is

$$\int_{\mathbb{R}} q(u) \sin(\xi u) \sin(\zeta u) du = \frac{1}{2} \{ \hat{q}(\xi - \zeta) - \hat{q}(\xi + \zeta) \}. \quad (\text{S66})$$

Double frequency integration carries $1/(2\pi^2)$. Exact row compression is applied to its full covariance by (S64).

Lemma S17.4 (Complete theta Fourier locality). *For every real ω ,*

$$|\hat{q}(\omega)| \leq \min\{32e^{-|\omega|/2}, 16384e^{-3|\omega|/4}\}.$$

If two frequency blocks start at $a_i, a_j \geq 0$, have separation d_{ij} and masses β_i, β_j , their normalized feature Gram block satisfies

$$\|G_{ij}\| \leq 16\sqrt{\beta_i\beta_j} \left(e^{-d_{ij}/2} + e^{-(a_i+a_j)/2} \right).$$

Normalization uses the true raw metrics plus nonnegative ridges; the estimates are independent of degree.

Proof. On $|\Im u| \leq 1/2$ put $r_n = \pi n^2 e^{2|\Re u|}$. Since $\cos 1 > 1/2$ and $|\cosh(u/2)| \geq (31/32) \cosh(\Re u/2)$, the complete theta series gives

$$|q(u)| \leq \frac{64}{31} \sum_{n \geq 1} (4r_n^2 + 6r_n) e^{-r_n/2}.$$

Its horizontal integral is at most

$$\frac{64}{31} \sum_{n \geq 1} (8\pi n^2 + 28) e^{-\pi n^2/2} < 32.$$

Substitute $r = \pi n^2 e^{2x}$ on both arms; integrate $(4r + 6)e^{-r/2}$ and bound the remaining scalar sum by its first terms and full geometric remainder. The complete double-exponential envelope makes the vertical contour ends vanish. Translation by $-\text{sign}(\omega)i/2$ proves the first bound.

For $|\Im u| \leq 3/4$, use $\cos(3/2) > 1/16$ and $\cos(3/8) > 9/10$. The corresponding full-series bound is

$$|q(u)| \leq \frac{20}{9} \sum_{n \geq 1} (4r_n^2 + 6r_n) e^{-r_n/16}.$$

Its integral over the two real arms is at most

$$\frac{20}{9} \sum_{n \geq 1} (64\pi n^2 + 1120) e^{-\pi n^2/16} \leq \frac{20}{9} (256S_2 + 1120S'_0),$$

where $S'_0 = \sum_{n \geq 1} e^{-3n^2/16} < 2$ and $S_2 = \sum_{n \geq 1} n^2 e^{-3n^2/16} < 15$. Indeed successive ratios give $S'_0 \leq e^{-3/16}/(1 - e^{-9/16})$, and

$$S_2 \leq \frac{32}{3e} \frac{e^{-3/32}}{1 - e^{-9/32}} < 15.$$

The integral is therefore less than 16384. The same complete contour argument at height $3/4$ proves the second bound.

A normalized source coefficient vector represents a weighted polynomial of L^2 norm at most one, since its metric contains the true raw Gram. Its weighted L^1 norm is at most $\sqrt{\beta_i}$. Apply this twice to (S66); the half-strip constant becomes $32/2 = 16$. $\square$

For a union of blocks with total mass B , a uniform retained-kernel error ϵ gives normalized full-matrix error at most $B\epsilon$. The rank-one comparison ($\sqrt{\beta_i\beta_j}$) proves this for the entire coefficient vector. Complete omitted interactions are paid by the same argument and their actual minimum argument. The finite-rank correction in (S64) remains dense.

S17.2.1 The complete real kernel polynomial

The source data supply the fixed exact dyadic polynomial

$$Q(\omega) = \sum_{k=0}^{512} a_k T_k(\omega/210), \quad \sup_{|\omega| \leq 210} |\hat{q}(\omega) - Q(\omega)| < 10^{-60}. \quad (\text{S67})$$

Odd coefficients vanish exactly. Here is the complete analytic construction. The preceding strip proof gives

$$|q(u)| \leq \frac{64}{31} (256/e^2 + 24/e) \frac{e^{-3/4}}{1 - e^{-9/4}} < 128 \quad (|\Im u| \leq 1/2).$$

Use ten physical panels of width $1/4$ on $[0, 5/2]$, 128 Gauss–Legendre nodes per panel and even reflection. For real $|\omega| \leq 210$ their symmetric Gaussian error is at most

$$E_G = 2 \cdot 10 \cdot \frac{8(1/8) 128 e^{630/32} 2^{-256}}{1 - 1/2}.$$

The omitted theta atoms after eight cost $E_\theta = 5 \cdot 16\pi^2 9^4 e^{-81\pi}$, and the full real tail is bounded by

$$E_\infty = \frac{480 e^{4U - \pi e^{2U}}}{2\pi e^{2U} - 4}, \quad U = 5/2.$$

Every enclosing Fourier node receives these complete errors. On the parameter-2 frequency ellipse, $|\Im \omega| < 158$ and

$$|\hat{q}(\omega)| \leq 480 \int_0^\infty e^{162u - 3e^{2u}} du \leq \frac{240 \cdot 80!}{3^{81}} =: M_Q.$$

Chebyshev coefficient bounds and complete Lobatto aliasing cost $4M_Q 2^{-512}$. Add the sum of all DCT coefficient radii when replacing them by exact dyadic midpoints. Odd coefficient balls contain zero before symmetry is imposed. The total is below $5.119 \cdot 10^{-63}$, establishing (S67). This separates analytic integration from coefficient approximation; compare [3].

The executed head uses this Q_{512} on $[-210, 210]$. Local evaluation first expands the *fixed exact polynomial* at a nearby integer center and keeps degree 256, then shifts to the actual fractional center and keeps degree 192. The two omitted polynomial parts are paid separately. Let $M_{24} < 1.26 \cdot 10^{44}$ bound Q on every radius-24 circle centered on $[-210, 210]$. The Bernstein ellipse with semimajor $1 + 24/210$ contains these disks, giving this bound directly from the literal coefficients. The fractional center shift has modulus at most $1/2$, and the actual convolution halfwidth is at most 4. Thus the first real remainder is at most

$$E_{\text{real}} = M_{24} \frac{(4.5/24)^{257}}{1 - 4.5/24}.$$

Cauchy's coefficient bound gives modulus at most $M_{24}/(1 - 16.5/24)$ for the intermediate degree-256 polynomial on the radius-16 disk about the actual center. The second remainder is at most

$$\frac{M_{24}}{1 - 16.5/24} \frac{(1/4)^{193}}{1 - 1/4}.$$

Their sum is below $3.411 \cdot 10^{-72} < 10^{-70}$. This is a bound for two polynomial truncations, independent of the real-kernel error in (S67).

The head retains every real kernel argument through 210. An omitted term has minimum argument at least 202. Consequently the normalized source debit

$$2B \left(16384e^{-3 \cdot 202/4} + 10^{-60} + 10^{-70} \right) \quad (\text{S68})$$

pays its complete kernel, far-image and local-model error. The true raw metrics and their enclosing entry errors are retained; the ridges dominate the model-metric discrepancies. The factor 2 safely covers the common-metric transport. It is paid on $(v, -Fv)$, without a further inverse-ridge factor.

S17.2.2 True symbol integration and source classes

The holomorphic symbol used for source quadrature is

$$m_M(z) = \frac{\Psi_0(1/4 + iz/2) + \Psi_0(1/4 - iz/2)}{2} - \log \pi - 2 \sum_{n \leq M} c_n \cos(z \log n).$$

Complete root brackets have opposite endpoint signs and a strict derivative sign. Every intervening real part is certified by a whole symbol box or monotonicity and endpoints; the entire negative support and positive exterior are included. If a root stripe has width η and $|m'_M| < 100$, its symbol magnitude is at most 100η and its integrated weight in $d\xi/\pi$ is at most $100\eta^2/\pi$. Thus the omitted stripe payment is quadratic in its width. All such stripes remain paid in the continuous source.

Each fixed-sign integration panel has length at most $1/4$. Its parameter-2 ellipse has real halfwidth $5h/4$ and imaginary halfwidth $3h/4$. The actual rectangles check $\Re(1/4 \pm iz/2) > 1/5$ and their distance from 1. The convergent digamma series gives $|\Psi_0(w)| \leq 1 + 6|w - 1|$ on this half-plane. The actual prime coefficients then fund the symbol bounds below. Mapped raw Chebyshev arguments have modulus less than 2, so $|T_k| \leq 5^k$.

For the true complex Fourier kernel,

$$|\hat{q}(z)| \leq \int_{\mathbb{R}} q(u) e^{|u|} du \leq 12, \quad |\Im z| \leq 1.$$

The complete envelope $q(u) \leq 16\pi^2 e^{4|u| - \pi e^{2|u|}}$ and the substitution $r = \pi e^{2|u|}$ give this gamma-integral bound. It applies with one frequency variable complex and the other real. A profile envelope M_{an} and Gaussian order n therefore give errors at most $8LM_{\text{an}}2^{-2n}$ for a raw metric and $16L_1L_2M_{\text{an}}2^{-2n}$ for a joint Gram entry. These errors concern the *true* analytic integrands. Only their real quadrature values are then replaced by fixed polynomials. Factorial-scaled moments evaluate that finite tensor rule exactly up to their enclosing arithmetic, retaining the second-variable sign and the two frequency measures.

Table 1 distinguishes the selected constructions. An observation count is separate from its number of source columns. The dense positive bins have width at most 2 and 16 features; each negative block has 13 features. Their maximal profile index is 15, so the analytic metric and one-variable Gram envelopes fit 10^{34} . The adopted total integration widths are below 270. Raw metric and Gram pads are 10^{-85} and 10^{-80} . Both $13 \cdot 10^{-85} < 10^{-26}$ and $16 \cdot 10^{-85} < 10^{-30}$ fund the corresponding negative and positive ridge comparisons. The remote groups use the same profile counts with their larger actual digamma rectangles. The head has 547 negative and 1680 positive columns, maximal profile index 15, analytic envelope 10^{30} and 10^{-100} raw entry pads. Its ridges are 10^{-25} and 10^{-28} . The other sources keep their own small-interval polynomials, reflected-term bounds, ridges and complete contracts. The typed index `data/m13_certificate.json` binds each group to these literal inputs.

Table 1: Source integration contracts for the selected constructions.

Source class	Observations	Positive bins	Gauss order	Symbol cap
Exact-row head	46	105	256	2000
Three dense groups	26	128	256	10000
Sixteen remote groups	35	variable	256	20000
Four repriced sources	4	5 each	256	own contract
Inherited isolated sources	138	5 each	own contract	own contract

S17.2.3 The original twelve-row head source

Write

$$\Xi(\xi) = \xi_{\text{Riemann}}(1/2 + i\xi) = \int_{\mathbb{R}} \psi(u) e^{-i\xi u} du.$$

Complete-theta integration by parts gives

$$\widehat{sz_{2j+1}}(\xi) = -i(-1)^j \xi^{2j+1} (\xi^2 + 1/4) \Xi(\xi).$$

For exact dyadic premixing coefficients A_{jk} , form

$$\tilde{z}_k = \sum_{j=0}^{10} A_{jk} z_{2j+1}, \quad P_k(\xi) = \sum_{j=0}^{10} A_{jk} (-1)^j \xi^{2j+1} (\xi^2 + 1/4)$$

before source evaluation. The actual mixed column is

$$\langle E_i, \tilde{z}_k \rangle = \int w(\xi) T_i(t_I(\xi)) P_k(\xi) \Xi(\xi) \frac{d\xi}{\pi}.$$

Exact combinations remain genuine nulls; their actual original Gram $A^* \Gamma_{11} A$ is retained. Section S17.3 gives the complete physical quadrature and reciprocal-theta bounds. The supplied complete enclosure differs from I in row-sum norm by less than $1.108 \cdot 10^{-60}$. This is a verified normalization, not a replacement of that Gram by I .

The polar row completes $\mathcal{Z} = [\tilde{z}_1, \dots, \tilde{z}_{11}, b]$. Its null crosses vanish exactly, and the complete Mellin identity gives

$$\|b\|_2^2 = \frac{1}{2} - \hat{q}(0) > 0.$$

Thus Γ is the direct sum of the full eleven-row Gram and this positive scalar. On X , projection off these twelve rows equals $Q_{K_{11}}$. The row b implements a Hilbert constraint; the operator-null transport uses K_{11} alone.

The remaining polar mixed source is

$$\langle E_i, b \rangle = \int w(\xi) T_i(t_I(\xi)) B(\xi) \frac{d\xi}{\pi}, \quad B(\xi) = \int_{\mathbb{R}} \psi(u) \tanh(u/2) \sin(\xi u) du.$$

It has a separately funded odd degree-1536 polynomial on $[-500, 500]$ with real uniform error below 10^{-85} . Its complete construction uses ten physical panels on $[0, 5/2]$, order 192, eight explicit theta atoms and both full omitted tails. The physical integrand bound is $256e^{500 \cdot 3/32}$; its Gaussian payment is

$$2 \cdot 10 \cdot 8(1/8) \frac{256e^{500 \cdot 3/32} 2^{-384}}{1 - 1/2}.$$

The omitted atom and real-tail payments are respectively

$$10 \cdot 16\pi^2 9^4 e^{-81\pi}, \quad \frac{480e^{9U/2-\pi e^{2U}}}{2\pi e^{2U} - 9/2}, \quad U = 5/2.$$

Their sum is below 10^{-90} at every frequency node. On the parameter-2 frequency ellipse a complete bound is $240 \cdot 189!/3^{189}$. Lobatto aliasing costs at most four times this bound times 2^{-1536} ; the sum of all coefficient radii is also paid. The resulting total is below $5.790 \cdot 10^{-89}$. Its local degree-256 polynomial remainder is below $3.784 \cdot 10^{-121}$. This independent polar source funds the actual row, including its sine normalization.

For the true null and polar mixed integrands the complete contour bound $|\Xi(x+iy)| \leq 768e^{-|x|/2}$ and $|B(x+iy)| \leq 768$, $|y| \leq 2$, supplies source majorants. Here is a direct scalar admission for these constants. On either $\text{Im } u = \pm 1/2$, reflection and $\cos 1 > 1/2$ give

$$|\psi(u)| \leq e^{|\text{Re } u|/2} \sum_{n \geq 1} (4r_n^2 + 6r_n) e^{-r_n/2}, \quad r_n = \pi n^2 e^{2|\text{Re } u|}.$$

Since $e^{5x/2} = (r_n/(\pi n^2))^{5/4} \leq r_n^2/(\pi n^2)^{5/4}$ for $x \geq 0$, the two-arm weighted integral is bounded by

$$\pi^{-5/4} \sum_{n \geq 1} n^{-5/2} \int_0^\infty (4r^3 + 6r^2) e^{-r/2} dr < \frac{480}{3} \frac{5}{3} < 768.$$

The double-exponential tails allow translation by $-i \text{sign}(x)/2$ in the Fourier integral, giving the stated Ξ bound. On the real physical line, $|\tanh(u/2)| \leq 1$ and $|\sin((x+iy)u)| \leq e^{2|u|}$; the same larger positive envelope gives the stated polar bound.

On each actual frequency ellipse, $|z| < 500$, $|T_i| < 5^{15}$ and $|m_{13}| < 10000$. The exact coefficient sum $B_k = \sum_\ell |[P_k]_\ell| 500^\ell$ bounds each premixed polynomial. For a total frequency length at most 1000, a safe Gaussian payment is

$$16 \cdot 1000 (10000 5^{16} 768 B_k / \pi) 2^{-512}.$$

On the real line, positivity of ψ gives $|\Xi(\xi)| \leq \Xi(0) < 1$. To justify the strict scalar bound in this normalization, the Mellin identity of Appendix A gives

$$\Xi(0) = \xi(1/2) = \frac{\Gamma(1/4)\eta(1/2)}{8\pi^{1/4}(\sqrt{2}-1)}.$$

The alternating-series upper bound is $\eta(1/2) \leq 1 - 1/\sqrt{2} + 1/\sqrt{3} < 9/10$. Splitting the gamma integral at one gives $\Gamma(1/4) < 4 + e^{-1} < 35/8$. Finally $\pi^{1/4} > 13/10$ and $\sqrt{2} - 1 > 2/5$ yield $(35/8)(9/10) < 8(13/10)(2/5)$, proving $\Xi(0) < 1$. For at most 1000 root stripes of width at most η , their complete mixed-column payment is therefore at most $100 \cdot 1000 \eta^2 B_k / \pi$. The actual stripe endpoints allow the smaller corresponding polynomial coefficient sums. The Gaussian and root-stripe errors are evaluated in each column, with all arithmetic balls retained. Complete row integration costs are below $8.281 \cdot 10^{-104}$ for quadrature and $2.522 \cdot 10^{-75}$ for stripes. Together with the full original Gram, the separate polar model and the complete ambient covariance, these are the joint inputs to (S65). Exact row premixing does not import locality through the projection or through an inverse.

S17.3 Complete physical source for the original null Gram

The covariance in Section S17.2.3 uses the original norm, including its reciprocal-theta factor. Here we give the bounds underlying its physical quadrature. Set

$$F_j = (D^2 - 1/4)D^{2j+1}\psi, \quad G_{jk}^0 = 2 \int_0^\infty \frac{F_j(u)F_k(u) \cosh(u/2)}{\psi(u)} du, \quad 0 \leq j, k \leq 10.$$

Define the exact polynomial operator $\mathcal{D}Q = \frac{1}{2}Q + 2rQ' - 2rQ$ and

$$Q_j = (\mathcal{D}^2 - 1/4)\mathcal{D}^{2j+1}(4r^2 - 6r), \quad d_j = \deg Q_j = 2j + 5, \quad S_j = \sum_\ell |[Q_j]_\ell|.$$

Thus $F_j(z) = e^{z/2} \sum_{n \geq 1} Q_j(\pi n^2 e^{2z}) e^{-\pi n^2 e^{2z}}$. All these polynomials have exact rational coefficients.

Consider $z = x + iy$, $x \geq 0$, $|y| \leq .01$, and put $r = \pi e^{2x}$ and $r_c = \pi e^{2z}$. Then $\Re r_c > .99r$ and $|r_c| = r \geq \pi$. The first theta atom satisfies $|4r_c^2 - 6r_c| \geq 2r^2$. Relative to this atom, the sum of the remaining absolute values is at most

$$3 \sum_{n \geq 2} n^4 e^{-.99r(n^2-1)} \leq 64e^{-3 \cdot .99r} < .1.$$

Indeed the successive-term ratio is at most $(3/2)^4 e^{-5 \cdot .99\pi} < 1/4$. Consequently the complete denominator is nonzero and

$$|\psi(z)| \geq \frac{9}{5} r^2 e^{x/2-r}. \quad (\text{S69})$$

Splitting $.99 = .24 + .75$ and maximizing $t^{d_j} e^{-.24t}$ gives

$$C_j = \frac{S_j(d_j/.24)^{d_j} e^{-d_j}}{1 - e^{-3\pi/4}}, \quad |F_j(z)| \leq C_j e^{x/2-3r/4}.$$

Since $|\cosh(z/2)| \leq e^{x/2}$, it follows that

$$\left| \frac{F_j(z)F_k(z) \cosh(z/2)}{\psi(z)} \right| \leq \frac{5C_j C_k}{9\sqrt{\pi}} r^{-3/2} e^{-r/2} \leq M_{jk} := \frac{5C_j C_k}{9\pi^2} e^{-\pi/2}. \quad (\text{S70})$$

Each F_j is odd and ψ is even, so the Gram integrand is even. This also funds the part of the first quadrature ellipse with negative real part by reflection to $|\Re z|$.

The finite interval $[0, U]$, $U = 3$, is split into 120 equal cells with halfwidth $h = 3/240$. For the Bernstein parameter $\rho = 7/5$, $h(\rho - \rho^{-1})/2 < .01$. Order 512 Gaussian quadrature, including both physical arms, therefore has complete error at most

$$8UM_{jk} \frac{\rho^{-1024}}{1 - \rho^{-1}}.$$

At every real node eight theta atoms are evaluated. The remaining denominator and jet tails are respectively bounded by

$$8e^{u/2} r^2 9^4 e^{-81r}, \quad 2S_j e^{u/2} r^{d_j} 9^{2d_j} e^{-81r}.$$

The successive-term bounds $(10/9)^{2d_j} e^{-19\pi} < 1/2$ fund all omitted atoms. These positive-radius balls are included before the reciprocal of ψ is taken; positivity of the complete real denominator

is checked. Finally $r(U) = \pi e^6 > 1200$. Integrating the first bound in (S70), with $du = dr/(2r)$, funds the two-arm physical tail by

$$\frac{10C_j C_k}{9\sqrt{\pi}} 1200^{-5/2} e^{-600}.$$

Both errors are added to every raw Gram entry. The exact dyadic map A then gives the complete enclosure $A^* G^0 A$, with all interval radii retained. The row-sum allowance stated in Section S17.2.3 uses this enclosure. The portable physical generator evaluates these formulas anew; it does not substitute an identity matrix for the conditioned Gram.

S17.4 Joint error used by the complete inverse-action application

For the declared head input, let B_{mass} be the sum of the actual zeroth raw-metric entries over all negative and positive source blocks, and let the enclosing integration balls be retained throughout. The complete ambient kernel debit is

$$\delta_{\text{src}} = 2B_{\text{mass}} \left(16384e^{-3 \cdot 202/4} + 10^{-60} + 10^{-70} \right) < 4.129 \cdot 10^{-58}.$$

The full raw metrics plus their deterministic ridges dominate the true frequency metrics: each block of size d has total entry uncertainty less than half its ridge divided by d . This comparison and the displayed scalar bound are checked by directed arithmetic on the literal input. The original twelve-row Gram and every row/source cross are enclosing balls, not midpoint replacements. They are inserted into (S64) before the residual covariance is formed, so their quadrature, stripe and reciprocal-Gram uncertainties remain in that full expression. Applying the ambient two-sided source error to the entire map $[I; -F]$ gives its debit $\delta_{\text{src}}(D_n + F^* D_p F)$; the squared residual construction in Section S6.4 pays the resulting full mixed error. No locality or identity-Gram hypothesis is applied after row compression. The code `regenerate_inverse_residual.py` reconstructs this complete tuple; the saved-array reader separately checks the retained residual matrices. The proofs and data have distinct roles: scientific literal coefficients specify the finite object, while the preceding arguments supply its complete continuous, kernel and tail error contract.

S18 Arithmetic bounds and complete signed-deficit coverage

This section combines a complete compact arithmetic bound with a direction-sensitive test of its benefit to a signed certificate. Weighted Schur bounds, constrained inverses and quadratic completion are classical tools [18, 7]. We give the relevant derivations in full. All comparisons of receiving blocks below use one fixed original physical metric, constraint space and receiving core. They do not identify a scalar arithmetic saving with a signed matrix margin.

S18.1 A sharp leading arithmetic bound

Put $\ell = \log K > 0$ and, on $L^2([0, \ell])$ with zero extension, let

$$C_K = \sum_{2 \leq n < K} \frac{\Lambda(n)}{\sqrt{n}} (T_{\log n} + T_{\log n}^*), \quad (T_t f)(x) = f(x - t).$$

Every active prime power and both shift orientations are retained. This is the complete compact arithmetic block when the arithmetic cutoff is at least K . Define the truncated sums

$$\Theta_K(y) = \sum_{2 \leq n < K, n \leq y} \Lambda(n), \quad A_K(y) = \sum_{2 \leq n < K, n \leq y} \frac{\Lambda(n)}{n}, \quad \phi(x) = e^{-x/2} + e^{-(\ell-x)/2}.$$

For $1 < z < K$ put

$$r_K(z) = \frac{\sqrt{K}[\Theta_K(z) + A_K(K/z)] + z[\Theta_K(K/z) + A_K(z)]}{\sqrt{K} + z}, \quad B_{\text{ES}}(K) = \operatorname{ess\,sup}_{1 < z < K} r_K(z).$$

Proposition S18.1 (Complete arithmetic price). *The two-sided bound $\|C_K\| \leq B_{\text{ES}}(K)$ holds. For integer K , the upper is determined by finitely many one-sided values at the rational breakpoints $1, K, n, K/n$. Moreover, using the classical unconditional PNT,*

$$\|C_K\| = (1 + o(1))\sqrt{K} \quad (K \rightarrow \infty).$$

The same leading asymptotic holds on the centered odd space after any finite-dimensional odd row constraints, with no restriction on their dimension as a function of K .

Proof. Exact summation of the allowed shift overlaps gives

$$(C_K\phi)(x) = e^{-x/2}[\Theta_K(e^x) + A_K(e^{\ell-x})] + e^{-(\ell-x)/2}[\Theta_K(e^{\ell-x}) + A_K(e^x)].$$

Thus $(C_K\phi)/\phi = r_K(e^x)$. For each edge use $2|f(x)f(y)| \leq |f(x)|^2\phi(y)/\phi(x) + |f(y)|^2\phi(x)/\phi(y)$. Integration over the full overlaps and summation over both orientations give $|\langle f, C_K f \rangle| \leq \int |f(x)|^2 r_K(e^x) dx$. The operator is self-adjoint; this proves its norm bound without assuming it positive semidefinite. On every open breakpoint cell the four partial sums are constant, so $r_K(z) = (\sqrt{K}a + zb)/(\sqrt{K} + z)$ has derivative of constant sign. Its supremum is at a one-sided endpoint. Duplicate breakpoints must be merged; a value using the wrong side's sums is not an upper. Interval-enclosed noninteger cutoffs require certified ordering and membership decisions.

For the asymptotic, $\Theta_K(y) = y + o(y)$ on growing arguments, $\Theta_K(y) = O(y)$, and $A_K(K) = O(\log K)$. These follow from the unconditional PNT [10] and partial summation. Excluding $n = K$ changes the endpoint by at most $\log K$. Replacing the two Θ_K terms by their arguments gives exactly $\sqrt{K}$. On $K^{1/4} \leq z \leq K^{3/4}$ the relative PNT error tends uniformly to zero. For $z \leq K^{1/4}$ the small-argument error contributes $O(K^{1/4})$, and the other argument is at least $K^{3/4}$, giving $o(\sqrt{K})$. Reflection $z \mapsto K/z$ handles the other end. The two A_K terms form a convex average bounded by $O(\log K)$. Consequently $r_K/\sqrt{K} \rightarrow 1$ uniformly. The Rayleigh quotient at ϕ is its ϕ^2 -weighted average, proving the matching lower. The assertion about odd constraints follows from Lemma S18.2 below, applied after centering. $\square$

For completeness, an exact finite lower uses

$$\|\phi\|^2 = 2(1 - K^{-1}) + 2\ell/\sqrt{K},$$

$$\langle \phi, T_{\log n} \phi \rangle = 2(n^{-1/2} - n^{1/2}/K) + \log(K/n)(\sqrt{n/K} + 1/\sqrt{Kn}).$$

These formulas follow by expanding the four exponential products on $[\log n, \ell]$. The Rayleigh numerator is twice their sum with weights $\Lambda(n)/\sqrt{n}$. No frequency quadrature is involved. Scalar minima of independently valid complete norm uppers are permitted; entrywise minima of Hermitian matrix uppers are not.

Lemma S18.2 (Finite-head invariance of the pure arithmetic norm). *Let C be bounded self-adjoint on the line or a centered interval, commute with reflection, and be an operator-norm limit of finite sums $M_a T_t$ with bounded multipliers, including interval restrictions. For any finite-dimensional row space in the odd Hilbert space, with orthogonal complement projection Q ,*

$$\|QCQ|_{\operatorname{ran} Q}\| = \|C\|.$$

Proof. For $U_\omega f = e^{i\omega x} f$, $U_\omega^* M_a T_t U_\omega = e^{-i\omega t} M_a T_t$. Simultaneous recurrence on each finite torus of shift phases and a diagonal choice through the norm approximants give $\omega_j \rightarrow \infty$ with $\|U_{\omega_j}^* C U_{\omega_j} - C\| \rightarrow 0$ for both signs. To see unbounded recurrence directly, apply compactness to the integer-time orbit of the finite phase vector: arbitrarily close orbit points have differences arbitrarily close to the identity. Either these return times are unbounded, or a bounded subsequence yields an exact return whose multiples are unbounded.

Choose a unit, parity-pure f with $|\langle f, Cf \rangle|$ arbitrarily close to $\|C\|$, possible because the two parity spaces reduce C . Use $v_j = f \cos(\omega_j x)$ if f is odd, and $v_j = f \sin(\omega_j x)$ if f is even. Then v_j is odd, $\|v_j\|^2 \rightarrow 1/2$, and $v_j \rightharpoonup 0$. The modulation identity and Riemann–Lebesgue applied to $\bar{f} C f \in L^1$ give $\langle v_j, C v_j \rangle \rightarrow \langle f, Cf \rangle/2$. Finite rank implies $\|(I - Q)v_j\| \rightarrow 0$; boundedness of C preserves the Rayleigh limit after projection. This proves the lower; compression gives the reverse inequality. $\square$

The lemma applies to C_K and, with the complete norm-convergent prime tail proved in the main paper, to the bounded theta-sandwiched prime block. It does not apply to the unbounded full Weil operator or automatically to an inverse-weighted return. The recurrent frequencies may have large archimedean energy. For legal form tests one may first choose smooth compact Rayleigh approximants and use operator-domain rows; this adds no uniform energy or frequency bound.

S18.2 The same-core receiving improvement

Work in the original physical Hilbert norm on a fixed support window. Let Q project off all fixed constraints and the finite receiving core. All restrictions use the original minimal closed form. Core columns lie in the associated operator domain, and their finite projection preserves the form domain; mixed columns are actual operator actions. Suppose the complete finite polar correction has its range in that excluded space, and the actual tail is

$$T = Q[gI + M_s a(D) M_s + M_s V_{\text{ar}} M_s] Q|_{\text{ran } Q}, \quad V_{\text{ar}} = m_0 I - C_K, \quad g > 0.$$

Assume a paid archimedean comparison $Q M_s a(D) M_s Q \succeq \gamma Q M_{s^2} Q$ and two valid scalar prices $B_0 \geq B_1 \geq \|V_{\text{ar}}\|$, with $G = B_0 - B_1 \geq 0$. For $\beta_i = \gamma - B_i > 0$ define

$$D_{\beta_i} = Q(gI + \beta_i M_{s^2}) Q, \quad Z_{\beta_i} = D_{\beta_i}^{-1}.$$

Thus $T \succeq D_{\beta_1} \succeq D_{\beta_0}$ and $T^{-1} \preceq Z_{\beta_1} \preceq Z_{\beta_0}$. These inverses act on $\text{ran } Q$, extended by zero elsewhere. If W is an orthonormal map spanning the entire excluded finite space, and $d_\beta = g + \beta s^2$, their explicit formula is

$$Z_\beta = d_\beta^{-1} - d_\beta^{-1} W (W^* d_\beta^{-1} W)^{-1} W^* d_\beta^{-1}.$$

Indeed the right side annihilates W , has range in its orthogonal complement, and multiplication by $Q d_\beta Q$ gives Q there. It is not the generally different compression $Q d_\beta^{-1} Q$.

Put $A_s = Q M_{s^2} Q$. The bounded resolvent identity gives

$$Z_{\beta_0} - Z_{\beta_1} = G Z_{\beta_0} A_s^{1/2} (I + G A_s^{1/2} Z_{\beta_0} A_s^{1/2})^{-1} A_s^{1/2} Z_{\beta_0}. \quad (\text{S71})$$

Since $D_{\beta_0} \succeq \beta_0 A_s$, $A_s^{1/2} Z_{\beta_0} A_s^{1/2} \preceq \beta_0^{-1} I$. Hence

$$\frac{G \beta_0}{\beta_0 + G} Z_{\beta_0} A_s Z_{\beta_0} \preceq Z_{\beta_0} - Z_{\beta_1} \preceq G Z_{\beta_0} A_s Z_{\beta_0}, \quad Z_{\beta_0} - Z_{\beta_1} \preceq \frac{G}{\beta_0 + G} Z_{\beta_0}. \quad (\text{S72})$$

The last assertion is scalar functional calculus for $Gt/(1+Gt)$ on $0 \leq t \leq \beta_0^{-1}$ after conjugating by $Z_{\beta_0}^{1/2}$. These bounds include $G = 0$ and use no cutoff-uniform minimum of s .

For the positive reference $T_0 = Q[M_s a(D)M_s + B_1 M_s^2 + gI]Q$, the bounded difference satisfies $0 \preceq T_0 - T \preceq 2B_1 A_s$. Therefore

$$cT_0 \preceq T \preceq T_0, \quad c = \frac{\beta_1}{\beta_1 + 2B_1}.$$

For an already paid core with $\gamma = B_0 + 1$, prices $B_0 = (2 + o(1))\sqrt{K}$ and $B_1 = (1 + o(1))\sqrt{K}$ give $c \rightarrow 1/3$. An exact choice for the former price is $|m_0|$ plus twice the sum of $\Lambda(n)/\sqrt{n}$ for $n \leq \sqrt{K}$, plus its sum for $\sqrt{K} < n < K$. Each killed translation has norm at most one. For $t > \ell/2$, the two end intervals joined by $T_t + T_t^*$ are disjoint; this operator exchanges them isometrically and is zero on the remaining interval, so its norm is one. The unconditional PNT and partial summation give $\sum_{n < x} \Lambda(n)/\sqrt{n} = (2 + o(1))\sqrt{x}$, while the $n \leq \sqrt{K}$ contribution is $O(K^{1/4})$ by the elementary Chebyshev bound. Hence B_0 has the stated leading term. The latter price follows from Proposition S18.1, with $|m_0|$ added to both and the scalar minimum with B_0 taken if needed at finite K . This restores a uniform relative inverse factor on that same paid core. It does not assert that such a core is small.

For clarity, writing $R_0 = T_0^{-1}$ and $C = R_0^{1/2}(T_0 - T)R_0^{1/2}$ gives $0 \preceq C \preceq (1 - c)I$ and $T^{-1} = R_0^{1/2}(I - C)^{-1}R_0^{1/2}$ by coercive form factorization. If $c = 1$, $T = T_0$ and the inverse is exact. For $0 < c < 1$, the Neumann inverse remainder after degree n is bounded by $(1 - c)^{n+1}R_0/c$. For $n \geq 1$, Chebyshev residual polynomials on $[c, 1]$ give the factor $\delta_n = 1/T_n((1+c)/(1-c)) \leq 2[(1 - \sqrt{c})/(1 + \sqrt{c})]^n$: for $p_n(\lambda) = (1 - r_n(\lambda))/\lambda$ and $r_n(\lambda) = T_n((1+c-2\lambda)/(1-c))/T_n((1+c)/(1-c))$, one has $(1 - \delta_n)\lambda^{-1} \leq p_n(\lambda) \leq (1 + \delta_n)\lambda^{-1}$. Full mixed-column congruence retains these inequalities. Approximate reference actions still require their own error enclosures.

S18.3 An optimized complete return and its signed test

Let H_c be the finite signed core and J its complete mixed map into the tail. Every J, E, Y column below takes values in $\text{ran } Q$. Put $X = J^*T^{-1}J$ and $K_{\text{true}} = H_c - X$. For a finite map Y with columns in $\text{Dom } T$, define

$$r = J - TY, \quad L(Y) = J^*Y + Y^*J - Y^*TY, \quad U_Z(Y) = L(Y) + r^*Zr \quad (Z \succeq T^{-1}).$$

Expansion proves the complete two-sided enclosure

$$H_c - L(Y) - r^*Zr \preceq K_{\text{true}} \preceq H_c - L(Y), \quad X = L(Y) + r^*T^{-1}r. \quad (\text{S73})$$

Thus a positive lower certifies this signed test; a negative direction of the upper certifies its failure. A negative lower alone is inconclusive. All action, interval and omitted-tail errors must be included on their proper sides. No $\text{Dom } T^2$ premise is used.

For independent trial columns $E \subset \text{Dom } T$, set

$$V = J^*ZJ, \quad A_E = (TE)^*Z(TE) - E^*TE, \quad F_E = (TE)^*ZJ - E^*J.$$

Then $U_Z(EC) = V - F_E^*C - C^*F_E + C^*A_EC$. Taking $Z = (1 + \varepsilon)Z_\beta$, $\varepsilon > 0$, gives $A_E \succeq \varepsilon E^*TE \succeq \varepsilon gE^*E \succ 0$. Square completion therefore gives the unique optimizer

$$C_{\text{opt}} = A_E^{-1}F_E, \quad U_{\text{opt}} = V - F_E^*A_E^{-1}F_E \succeq X.$$

Enlarging the trial span cannot increase this matrix upper. For the two prices on the same core, write $Z_i = (1 + \varepsilon)Z_{\beta_i}$ and $r_0 = J - TY_0$ at the old optimizer. Evaluation of the new quadratic at Y_0 proves

$$M_{\text{gain}} := U_{\text{opt},0} - U_{\text{opt},1} \succeq r_0^*(Z_0 - Z_1)r_0 \succeq \frac{(1 + \varepsilon)G\beta_0}{\beta_0 + G}(sZ_{\beta_0}r_0)^*(sZ_{\beta_0}r_0) =: G_*.$$

This computable gain lower preserves the full residual and every cross term. Its factor saturates; growing scalar savings need not give growing gains in every coefficient direction.

Proposition S18.3 (Exact deficit coverage). *Let $D = U_{\text{opt},0} - H_c$. The improved certificate passes exactly when $M_{\text{gain}} - D \succeq 0$. For any positive lower $G_* \preceq M_{\text{gain}}$, $G_* - D \succeq 0$ is sufficient. Decompose the coefficient space into $\text{ran } G_* \oplus \ker G_*$ and put $A = -D_{NN}$. Then this sufficient test is equivalent to*

$$A \succeq 0, \quad D_{RN} \ker A = 0, \quad (G_*)_R \succeq D_{RR} + D_{RN}A^\dagger D_{NR}.$$

When $G_* = 0$ the test is simply $D \preceq 0$; when its kernel is empty only the full positive-definite comparison remains.

Proof. The exact identity is $H_c - U_{\text{opt},1} = M_{\text{gain}} - D$. The displayed block conditions follow by completing the square for $G_* - D$ with bottom block A . Positivity forces the off-diagonal to annihilate $\ker A$; after that compatibility the remaining block is $(G_*)_R - D_{RR} - D_{RN}A^\dagger D_{NR}$. $\square$

A numerical near-kernel cannot be treated as an exact kernel. For a strictly positive G_* the test is $\lambda_{\max}(G_*^{-1/2}DG_*^{-1/2}) \leq 1$, with the full matrix-error budget paid. It is not necessary for coverage when the lower gain is strictly smaller than M_{gain} . For example $G_* = \text{diag}(1, 0)$ misses $D = \text{diag}(0, 1/10)$ despite a larger gain trace. For $D = \begin{bmatrix} 0 & 1/10 \\ 1/10 & 0 \end{bmatrix}$ the kernel diagonal is zero but the mixed compatibility fails. Both are exact finite interface examples, not Weil counterexamples.

Finally, repricing must conserve the actual positive surplus. If an arithmetic pair has old prices a, d and new price $\varphi = a + d - G$, then

$$\varphi I - (C_n + C_m) = (aI - C_n) + (dI - C_m) - GI.$$

Increasing the baseline by G while retaining both old surplus features counts that gain twice. For instance the true tail $T = 1$ may be written as $1/4 + 3/4$ or $3/4 + 1/4$; combining the new floor with the old surplus falsely gives $3/2$. At $H_c = 4/5, J = 1$ this changes the true Schur value $-1/5$ to the false positive $2/15$. Rebuild the actual blocks after any core change. For a shifted relative-covariance test, shift the core, mixed map and tail together before applying the identities above. These are complete acceptance and failure tests; their success at growing cutoffs still requires the actual joint matrices and all original-norm return payments.

S19 Actual-prime discrepancy and physical packet energy

This section gives an analytic arithmetic input for the finite-verification framework. Mellin smoothing controls the actual prime-power discrepancy; uniform packet conditioning expresses that control in the original unweighted logarithmic energy. The mother function and separation are fixed throughout, while the number and highest frequency of the packets may grow. All Fourier integrals use the main paper's measure $d\xi/(2\pi)$.

S19.1 A fixed-window Mellin estimate

Assume every nontrivial zero of ζ has real part at most b , where $1/2 \leq b < 1$. Fix $0 < \eta < 1 - b$ and put $d = b - 1/2 + \eta$. For $W \in C_c^\infty(\mathbb{R})$ write $\widetilde{W}(z) = \int W(u)e^{zu} du$.

Lemma S19.1 (Smoothed actual-prime discrepancy). *For each fixed W there is a constant $C_{W,b,\eta}$ such that, for every $T \geq 0$ and $\omega \in \mathbb{R}$,*

$$\left| \sum_{n \geq 2} \Lambda(n) n^{-1/2-i\omega} W(\log n - T) - e^{(1/2-i\omega)T} \widetilde{W}(1/2 - i\omega) \right| \leq C_{W,b,\eta} e^{dT} \log(2 + |\omega|).$$

The constant is bounded by a b, η -dependent multiple of

$$\mathcal{A}_d(W) = \int_{\mathbb{R}} |\widetilde{W}(d + iv)| [1 + \log(2 + |v|)] dv.$$

The main term is the full integral over $x > 0$. It also equals the integral over $x \geq 1$ when $T + \text{supp } W \subset [0, \infty)$.

Proof. We first pay the logarithmic derivative on the line $\sigma_0 = b + \eta$. Euler summation gives, for $\Re s > 0$, $s \neq 1$,

$$\zeta(s) = \frac{s}{s-1} - s \int_1^\infty \{x\} x^{-s-1} dx.$$

The remainder converges absolutely and gives polynomial growth $|\zeta(s)| \leq C(1 + |\Im s|)$ on a fixed strip away from its pole. For $|t| \geq 4$, the disk centered at $2 + it$ with radius $R = 2 - b - \eta/4$ is zero-free and avoids the pole. Its analytic logarithm $h = \log \zeta$ has uniformly bounded center value, since the Euler logarithm there is absolutely bounded by $\sum_{n \geq 2} (n^2 - 1)^{-1} = 3/4$. The growth estimate bounds $\Re h$ above by $C + \log(2 + |t|)$ on the disk.

Here is the needed real-part estimate. If f vanishes at the disk center and $\Re f \leq B$, $B > 0$, then $f/(2B - f)$ maps the disk to the closed unit disk. Schwarz's lemma therefore gives $|f| \leq 2Br/(R - r)$ on its radius- r subdisk. Apply this to $h - h(2 + it)$ with $r = 2 - b - \eta/2$, and then apply Cauchy's derivative estimate on a radius- $\eta/4$ circle about $\sigma_0 + it$. It follows that

$$|\zeta'/\zeta(\sigma_0 + it)| \leq C_{b,\eta} \log(2 + |t|).$$

The same argument is uniform between real parts σ_0 and $3/2$ at large heights. At bounded height the fixed line is zero-free and avoids 1, so continuity supplies its bound there.

Bilateral Laplace inversion on $\Re z = 1$ and absolute convergence of the Dirichlet series give the exact expression

$$\frac{1}{2\pi i} \int_{\Re z=1} -\frac{\zeta'}{\zeta}(1/2 + i\omega + z) e^{zT} \widetilde{W}(z) dz.$$

Shift to $\Re z = d$. The sole crossed pole is at $z = 1/2 - i\omega$, with residue $e^{(1/2-i\omega)T} \widetilde{W}(1/2 - i\omega)$. At each fixed T, ω , horizontal segments at heights larger than $|\omega| + 4$ vanish: repeated integration by parts gives rapid vertical decay of $\widetilde{W}$, while the logarithmic derivative has the bound just proved. The remaining integral is

$$\frac{1}{2\pi} \int_{\mathbb{R}} -\frac{\zeta'}{\zeta}(b + \eta + i(\omega + v)) e^{(d+iv)T} \widetilde{W}(d + iv) dv.$$

Using $\log(2 + |\omega + v|) \leq \log(2 + |\omega|) + \log(2 + |v|)$ pays its entire vertical tail and proves the estimate. Finally, substitution $x = e^{T+u}$ identifies the exact main integral and its stated support case. $\square$

S19.2 Complete mixed packet matrices

Fix a nonzero real even $\phi \in C_c^\infty(-r, r)$. For $\omega \geq 1$ put

$$g_\omega = -\omega^{-2}(\partial_u^2 - 1/4)[\phi(u) \cos(\omega u)], \quad \ell_\omega = \log(2 + \omega).$$

These functions are real, even and have both exponential moments zero. For a finite frequency list $\omega_i \geq 1$ with separation at least $h > 0$, set

$$H_{ij}(s) = \int_{\mathbb{R}} g_{\omega_i}(x + s) \overline{g_{\omega_j}(x)} dx, \quad L_\ell = \text{diag}(\ell_{\omega_i}).$$

The complete actual-prime discrepancy matrix is

$$\mathfrak{E}_{ij}(T) = \sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} H_{ij}(\log n - T) - \int_0^\infty x^{-1/2} H_{ij}(\log x - T) dx.$$

Proposition S19.2 (Uniform mixed-matrix control). *Under the strip hypothesis above,*

$$\|L_\ell^{-1/2} \mathfrak{E}(T) L_\ell^{-1/2}\| \leq C_{\phi, h, b, \eta} e^{dT}, \quad T \geq 0.$$

The constant is independent of list length and highest frequency.

Proof. Expand exactly

$$g_\omega = A_\omega \cos(\omega u) + B_\omega \sin(\omega u), \quad A_\omega = \phi - (\phi'' - \phi/4)/\omega^2, \quad B_\omega = 2\phi'/\omega.$$

Thus $g_\omega = \sum_{\sigma=\pm 1} a_{\omega, \sigma} e^{i\sigma\omega u}$, where $a_{\omega, \sigma} = (A_\omega - i\sigma B_\omega)/2$ has fixed support and uniform smooth seminorms. Each H_{ij} is the sum of four terms $e^{i\sigma\omega_i s} W_{ij, \sigma, \tau}(s)$, with

$$W_{ij, \sigma, \tau}(s) = \int a_{\omega_i, \sigma}(x + s) \overline{a_{\omega_j, \tau}(x)} e^{i(\sigma\omega_i - \tau\omega_j)x} dx.$$

Every envelope has support in $[-2r, 2r]$. Differentiate in s and integrate by parts in x . For all fixed $m, k \geq 0$,

$$\|\partial_s^m W_{ij, \sigma, \tau}\|_\infty \leq C_{\phi, m, k} (1 + |\sigma\omega_i - \tau\omega_j|)^{-k}.$$

The internal phase is independent of s , and compact smooth support removes all boundary terms. The direct bound handles phase magnitude at most one.

The seminorm $\mathcal{A}_d(W)$ in Lemma S19.1 is bounded by a constant times the L^1 norms of $e^{ds}W$ and its second derivative, using the direct and twice-integrated transform bounds. Hence the same off-resonance decay pays the changing-window constants. Apply the lemma to all four carriers, retaining their unit-modulus centering phases. This gives

$$|\mathfrak{E}_{ij}(T)| \leq C_k e^{dT} \ell_{\omega_i} [(1 + |\omega_i - \omega_j|)^{-k} + (1 + \omega_i + \omega_j)^{-k}].$$

Write $\Delta = |\omega_i - \omega_j|$. Since $\ell_{\omega_i} \leq \ell_{\omega_j} + \log(1 + \Delta)$ and $\ell_{\omega_j} \geq \log 3$, taking $k = 3$ bounds each normalized entry by $C e^{dT} (1 + \Delta)^{-2}$. For an h -separated list both its row sums and column sums are bounded by a multiple of $\sum_{m \geq 0} (1 + mh)^{-2}$. The two-sided Schur test proves the induced bound for the entire matrix. The real-even generators also imply $H_{ij} = H_{ji}$ by reflection, though the norm proof does not require it. $\square$

S19.3 The physical logarithmic-energy metric

Put

$$\mathcal{E}(g) = \|g\|_2^2 + \int_{\mathbb{R}} a(\xi) |\widehat{g}(\xi)|^2 \frac{d\xi}{2\pi}.$$

This is the unweighted profile energy. In particular its norm is distinguished from the original theta-weighted norm of h when $f = sh$.

Proposition S19.3 (Uniform physical-energy conditioning). *There are fixed $\Omega_0 \geq 1$, $h_0 > 0$ and $0 < c_\phi \leq C_\phi < \infty$ such that, for every finite list $\omega_i \geq \Omega_0$ separated by at least h_0 , and every complex coefficient vector c ,*

$$c_\phi \sum_i \ell_{\omega_i} |c_i|^2 \leq \mathcal{E} \left(\sum_i c_i g_{\omega_i} \right) \leq C_\phi \sum_i \ell_{\omega_i} |c_i|^2.$$

These constants are independent of list length and highest frequency.

Proof. The main paper's equation (11) gives $\frac{1}{2} \log(1 + 4\xi^2) \leq a(\xi) \leq \frac{1}{2} \log(1 + 4\xi^2) + 4$. Thus $a(\xi) = \log|\xi| + O(1)$ at infinity, with coefficient one. The carrier amplitudes above converge in every Schwartz Fourier seminorm to $\phi/2$ as $\omega \rightarrow \infty$.

Let G be the complete energy Gram. A carrier pairing is

$$\int_{\mathbb{R}} [1 + a(\xi)] \widehat{a}_i(\xi - \sigma\omega_i) \overline{\widehat{a}_j(\xi - \tau\omega_j)} \frac{d\xi}{2\pi}.$$

The symbol satisfies $1 + a(\xi) \leq C\ell_{\omega_i} [1 + \log(2 + |\xi - \sigma\omega_i|)]$. Split this integral where one of the two distances from its carrier is at least half their separation. Extra Schwartz powers on that factor and integration of the remaining weighted decay give, for every k ,

$$\frac{|G_{ij}|}{\sqrt{\ell_{\omega_i} \ell_{\omega_j}}} \leq C_k [(1 + |\omega_i - \omega_j|)^{-k} + (1 + \omega_i + \omega_j)^{-k}].$$

The logarithmic ratio used in the preceding proposition absorbs the normalization with one extra Schwartz power.

On the diagonal, substitute $\xi = \sigma\omega + v$. Dominated convergence with weight $1 + \log(2 + |v|)$ gives $\|\phi\|_2^2/4$ from each of the two equal-carrier terms after division by ℓ_ω . Opposite-carrier terms tend to zero by the separation estimate. Therefore $G_{ii}/\ell_{\omega_i} \rightarrow m_\phi = \|\phi\|_2^2/2 > 0$. Choose Ω_0 so every normalized diagonal G_{ii}/ℓ_{ω_i} lies in $[m_\phi/2, 2m_\phi]$. Then choose the fixed h_0 so the off-diagonal row and column sums, bounded by a constant times $\sum_{k \geq 1} (1 + kh_0)^{-2}$, are at most $m_\phi/4$. Schur's bound gives the asserted two-sided inequality, with $c_\phi = m_\phi/4$ and $C_\phi = 9m_\phi/4$. $\square$

Corollary S19.4 (Actual-prime control in physical energy). *Under the strip hypothesis, for the fixed mother, minimum frequency and separation in Proposition S19.3, let $g_c = \sum_i c_i g_{\omega_i}$ and $g_z = \sum_i z_i g_{\omega_i}$. Then*

$$|c^* \mathfrak{E}(T) z| \leq C e^{dT} \mathcal{E}(g_c)^{1/2} \mathcal{E}(g_z)^{1/2}.$$

There is a fixed T_0 , depending only on the mother support, such that for every $T \geq T_0$ this bound can also be paid by the energies of the odd polar-free profiles $f_{c,T} = \tau_{T/2} g_c - \tau_{-T/2} g_c$ and $f_{z,T}$.

Proof. Combine the two preceding propositions. For the odd-arm assertion take $T > 2r$. Both moments of g_c vanish by integration by parts, so its translated antisymmetrization is odd and exactly polar-free. Its disjoint arms give $\|f_{c,T}\|^2 = 2\|g_c\|^2$. The off-diagonal archimedean kernel is $-J$, where $J(t) = e^{-|t|/2}/(1 - e^{-2|t|})$. If $T - 2r \geq 1$, its cross pairing is at most $\epsilon_T \|g_c\|^2$, with

$\epsilon_T = (8r/3)e^{-(T-2r)/2}$, by Schur's test and $J(t) \leq (4/3)e^{-|t|/2}$. Choose one T_0 so $\epsilon_T \leq 1/2$ for $T \geq T_0$. Then

$$\mathcal{E}(g_c) \leq \mathcal{E}(f_{c,T}) \leq 3\mathcal{E}(g_c).$$

On the actual odd-half supported realization, $D_{\text{full}} = D_L + R_L$ and $0 \preceq R_L \preceq \pi I/2$; the corresponding norm-plus-DN energy is therefore equivalent with factor at most $1 + \pi/2$. Finally the self interaction uses only $|t| \leq 2r$, a fixed finite prime-power shift sum and a bounded continuous main kernel, so its discrepancy costs at most $C_r \|g_c\|^2$. This pays the self fee separately from the complete long cross matrix. $\square$

The supporting estimate retains the explicit coefficient e^{dT} . It controls all mixed entries on the specified, uniformly conditioned packet families. Its constants fix the mother and frequency separation, providing an arithmetic input in the same physical energy coordinates used for these odd test profiles.

S20 Verification of the complete supported example

This section specifies the computer-assisted inputs and the complete verification for the supported example in main-paper Section D. The original realization and closed domain are those proved in the main paper. All functions below are in the odd-half unitary profile space. The primary receiving trial is zero.

S20.1 Exact model and positive step reference

Fix

$$M = 2^{20}, \quad \kappa = M^{-1}, \quad L = \frac{1}{2} \log(4 \log M / \pi) + \frac{1}{10}, \quad J = 74, \quad N = 329.$$

The complete active prime powers are

$$2, 4, 8, 16, 3, 9, 5, 7, 11, 13, 17, 19,$$

each with weight $\Lambda(n)/\sqrt{n}$. Here $21 < e^{2L} < 22 < M$; larger translations have zero supported overlap. Both orientations of every remaining translation are retained. Let $r = \pi e^{2x}$, $r_L = \pi e^{2L}$, and

$$P = 6459 = \lceil 100(r_L - \pi) \rceil, \quad x_i = \frac{1}{2} \log \left(1 + \frac{(r_L - \pi)i}{\pi P} \right), \quad 0 \leq i \leq P.$$

The scientific data specify exact positive dyadic heights on these radial cells. The full-theta brackets proved in the main paper give, with

$$W_1(x) = \frac{(1 + e^{-x})e^{\pi e^{2x}}}{4\pi e^{2x}(2\pi e^{2x} - 3)},$$

the bounds $(128/129)W_1 \leq W \leq W_1$ and $|(\log W)'| < 21\pi e^{2x}$. Thus, on every subcell $[a, b]$, with $v = 21\pi e^{2b}(b - a)$, the directed checks

$$\frac{1}{2}W_1(a)e^v < w_j < \frac{128}{129}W_1(a)e^{-v}$$

prove $W/2 \leq W_{\text{low}} \leq W$ on the whole cell. They are checked for every cell, including the endpoints.

For completeness, the arithmetic row price is a whole-space form bound. On the full interval of length $\log K = 2L$, use the positive reflection-symmetric Schur weight $\phi(u) = e^{-u/2} + e^{-(2L-u)/2}$. For $z = e^{L+x}$, define the finite sums, with one-sided endpoint values,

$$\Psi(y) = \sum_{\substack{n < K \\ n \leq y}} \Lambda(n), \quad H(y) = \sum_{\substack{n < K \\ n \leq y}} \frac{\Lambda(n)}{n}, \quad K = e^{2L}.$$

Direct summation of $C^{\text{ar}}\phi/\phi$ gives

$$\mathfrak{r}(x) = \frac{\sqrt{K}\{\Psi(z) + H(K/z)\} + z\{\Psi(K/z) + H(z)\}}{\sqrt{K} + z}.$$

Applying $2|fg| \leq |f|^2\phi(y)/\phi(x) + |g|^2\phi(x)/\phi(y)$ to each translated edge proves $|\langle f, C^{\text{ar}}f \rangle| \leq \int \mathfrak{r}|f|^2$. Restriction to the odd subspace preserves this bound. The common partition adds every positive breakpoint $\log n - L$ or $L - \log n$ to the radial nodes and has 6471 cells. On each open cell the four partial sums are constant, so $\mathfrak{r} = (\sqrt{K}A + zB)/(\sqrt{K} + z)$ has its extrema at the two one-sided endpoints. The dyadic upper $r_{\text{up},j}$ is checked against both.

Let $p = \sinh(x/2)$, $p_q = \langle p, e_q \rangle$, and $\mathcal{V} = \text{span}(e_0, \dots, e_{328}) \cap p^\perp$. For Q the complete orthogonal complement of $[p, \mathcal{V}]$, the pole-tail argument gives

$$D_L|_Q \succeq \ell_N I, \quad \ell_N = a(k_N) - \delta_N \{a(k_N) - a(k_0)\}, \quad \delta_N \leq \frac{9L^3}{\pi^4(N + 1/2)^3}.$$

Indeed the low-band part of a receiving vector is parallel to the low-band pole projection; its squared fraction is at most the omitted pole fraction. Splitting the DN diagonal below and above N gives the displayed inequality. The exact pole coefficients are

$$p_q = \frac{\sqrt{2/L}(-1)^q \cosh(L/2)}{2(k_q^2 + 1/4)}.$$

The data fix a dyadic $\alpha \leq \ell_N + m_0$, approximately 4.675643352302501, and on each common cell set

$$d_j = \kappa w_j + \alpha - r_{\text{up},j} > 1/200.$$

The complete positive response, the diagonal suffix and the arithmetic row bound therefore prove

$$T = (D_L + R_{L,74} + m_0 - C^{\text{ar}} + \kappa W_{\text{low}})|_Q \succeq d_Q \succeq I/200.$$

This bound covers all receiving directions. Finite inputs and the bounded fixed-window perturbations use the same operator and minimal closed form domains as in the main realization proof.

S20.2 All constraints and the exact inverse direction

Write C for the raw core map with columns $u_j = e_j - (p_j/p_0)e_0$, and let $F = [p, C]$. Its Gram is $G_C = C^*C = I + vv^*$, $v_j = p_j/p_0$. For $G = F^*d^{-1}F$, the full constrained multiplication inverse is

$$Z = d^{-1} - d^{-1}FG^{-1}F^*d^{-1}.$$

Solving $dy = f$ modulo $\text{ran } F$, with $F^*y = 0$, proves this formula and $T^{-1} \preceq Z$ on $\text{ran } Q$. No orthonormality of the raw list is assumed. The archived upper-triangular matrix R_G has exact

entries in $2^{-256}\mathbb{Z}$ and positive diagonal. Complete directed integration and the absolute-row-sum bound give

$$\|R_G^* G R_G - I\| < 4 \times 10^{-75} = \delta, \quad G^{-1} \succeq R_G R_G^* / (1 + \delta).$$

The complete 329-direction congruence is checked; all directions are retained. For example, the needed elementary moments on a constant-weight cell follow from

$$e_q e_r = \frac{\cos((q-r)\pi x/L) - \cos((q+r+1)\pi x/L)}{L}, \quad \int p^2 = \frac{\sinh x - x}{2},$$

and

$$\int p e_q = \frac{\sqrt{2/L} \{ \frac{1}{2} \sin(k_q x) \cosh(x/2) - k_q \cos(k_q x) \sinh(x/2) \}}{k_q^2 + 1/4}.$$

The zero-frequency cosine integral uses its exact length limit.

S20.3 Complete output moments and symbolic ridge cancellation

Let $E = \hat{S}_* - d$. The identical ridge cancels symbolically; the logarithmic DN diagonal remains and acts exactly on finite DN inputs. On each fully split cell the primitive action on e_q is

$$E e_q = \sqrt{2/L} \{ (A_q + \rho) \sin(k_q x) + B_q \cos(k_q x) \} + \sum_{j < 74} r_j(x) V_{jq}, \quad \rho = r_{\text{up},j},$$

where, for each active $t_n = \log n$,

$$\begin{aligned} A_q &= a(k_q) + m_0 - \alpha - \sum_{n < K} \frac{\Lambda(n)}{\sqrt{n}} (\chi_+ + \chi_-) \cos(k_q t_n), \\ B_q &= - \sum_{n < K} \frac{\Lambda(n)}{\sqrt{n}} (\chi_+ - \chi_-) \sin(k_q t_n), \\ V_{jq} &= \frac{\sqrt{2/L} (-1)^q c_j \sqrt{1 + e^{-2c_j L}}}{c_j^2 + k_q^2}. \end{aligned}$$

Here $\chi_+ = 1_{x+t_n < L}$, $\chi_- = 1_{x-t_n > -L}$. The odd extension supplies the reflected negative argument exactly. The full spatial output of every column is retained.

Products of these actions are integrated by elementary endpoint moments. In addition to trigonometric product-to-sum formulas, put $n_j = \sqrt{1 + e^{-2c_j L}}$. The required primitives are

$$\begin{aligned} \int e_q r_j &= \frac{2\sqrt{2/L} e^{-c_j L}}{n_j(c_j^2 + k_q^2)} \{ c_j \cosh(c_j x) \sin(k_q x) - k_q \sinh(c_j x) \cos(k_q x) \}, \\ \int \sqrt{2/L} \cos(k_q x) r_j &= \frac{2\sqrt{2/L} e^{-c_j L}}{n_j(c_j^2 + k_q^2)} \{ c_j \cosh(c_j x) \cos(k_q x) + k_q \sinh(c_j x) \sin(k_q x) \}, \\ r_i r_j &= \frac{2e^{-(c_i+c_j)L}}{n_i n_j} \{ \cosh((c_i + c_j)x) - \cosh((c_i - c_j)x) \}. \end{aligned}$$

The difference-frequency zero is integrated as x . Stable normalized exponentials evaluate the same formulas. Each region's action coefficients are applied before summing its moments; endpoint jumps include both outer endpoints.

For the 393 primitives $\mathbf{e} = (e_0, \dots, e_{392})$, the data contain

$$B = \mathbf{e}^* E \mathbf{e}, \quad K = (E \mathbf{e})^* d^{-1} E \mathbf{e}, \quad H_s = \mathbf{e}^* d^{-1} E \mathbf{e}, \quad P_E = p^* d^{-1} E \mathbf{e},$$

and the sine/response, pre-response-action/response and full response/response blocks VS, XR, RR . For example, with E_0 the displayed trigonometric part,

$$K = E_0^* d^{-1} E_0 + XRV + V^* XR^* + V^* RRV.$$

Thus the response Gram is the full spatial Gram, including every cross term. The relation $p = (n_0 e^{L/2}/2)r_0$ supplies the full pole/action row from the same weighted moments.

S20.4 Finite sign and optional receiving enrichment

The zero-trial main-paper matrix uses

$$H = \begin{bmatrix} 0 \\ G_C \end{bmatrix} + \begin{bmatrix} P_E \\ C^* H_s \end{bmatrix} C, \quad A_\star = \frac{H^* R_G R_G^* H}{1 + \delta} - C^* (B + K) C,$$

where C in these products is its 393-row primitive coefficient map. All 329 weighted constraint rows are applied before finite core conditioning. Since R_G is upper triangular, its core block R_{uu} is invertible, and

$$G_C^{-1} = I - \frac{vv^*}{1 + v^*v}, \quad P_C = G_C^{-1} R_{uu}^{-*}$$

is an invertible congruence. Every received directed pivot of $P_C^* A_\star P_C$ is positive.

The optional trial columns are

$$t_j = e_j - \frac{p_j}{\|p_N\|^2} p_N, \quad 329 \leq j \leq 392, \quad p_N = \sum_{q < N} p_q e_q.$$

They are exactly pole-free and core-orthogonal, with Gram $I + v_{\text{tail}} v_{\text{tail}}^* / \|p_N\|^2$. Together with the core they span the full constrained 393-mode prefix. For any exact dyadic coefficient matrix Z_t , the trial is legal. The original complete coefficient lower A_{all} is tested by

$$[I; -Z_t]^* A_{\text{all}} [I; -Z_t] = A_{uu} - A_{ut} Z_t - Z_t^* A_{tu} + Z_t^* A_{tt} Z_t.$$

A proposed inverse only selects the dyadic trial; the original lower is evaluated afterward. The archived trial has denominator 2^{384} .

Table 2 summarizes the finite sign checks for this case. The complete primitive assembly uses

Table 2: Directed sign checks for the M20 supported example.		
Candidate	Core dimension	Positive directed pivots
Zero receiving trial	328	328
64 legal receiving trials	328	328

2048-bit directed arithmetic. The 4096-bit replay reconstructs both candidates from all 153664 entries of A_{all} , reconstructs the analytical core conditioner, and checks all 107584 conditioner entries. Its four-block trial assembly and right-associated congruence use a different multiplication order from the producer. Completed-column LDL supplies all 656 positive pivots. These are sign certificates in their actual coordinates.

S20.5 Reproducibility files and scope of checks

The companion scientific archive `worked_example_data.zip` contains four numerical inputs and `INPUTS.json`, whose file hashes identify every exported numerical field. Decimal interval strings and exact dyadic integers are retained losslessly. The principal package contains the portable code under `code/worked_example/`. After extracting the companion archive into `data/worked_example/`, run the following from the principal package root:

```
python code/worked_example/check_reference.py -data data/worked_example
python code/worked_example/check_weighted_gram.py -data data/worked_example
python code/worked_example/compute_signed.py -data data/worked_example -output
signed_rebuilt.json
python code/worked_example/check_signed.py -data data/worked_example
```

The first checks all reference cells, theta floors, arithmetic endpoints and the complete receiving admission. The second regenerates the whole weighted Gram and tests its exact dyadic conditioner. The third reconstructs the complete signed lower from the archived mixed moments; the fourth independently receives the saved lower and replays both finite signs. `generate_mixed.py` provides full elementary moment regeneration; its input and output options are given in the reader instructions. The code uses Python with `python-flint/Arb`. The mathematical derivations, whole-domain implications and actual precision checks have the scopes stated above. The code and replays use the same arithmetic library. The zero-trial case is the primary evaluated supported certificate.

S21 Verification of the second moving-window certificate

This section supplies the M24 case of main-paper Section D.1. All definitions use the same original norm, exact polar constraint and minimal closed supported domain. The receiving trial is zero. The formulas in Section S20 remain valid with the new case-specific inputs listed here.

S21.1 Exact new case and complete reference

Set $M = 2^{24}$, $\kappa = 2^{-24}$, $L = \frac{1}{2} \log(4 \log M / \pi) + \frac{1}{10}$, $J = 88$, and $N = 660$. Here $25 < e^{2L} < 26$; the complete active prime powers are

$$2, 4, 8, 16, 3, 9, 5, 25, 7, 11, 13, 17, 19, 23.$$

The same exact dyadic floor construction uses 7,814 radial cells and the common partition has 7,828 cells. The additional endpoints are every interior arithmetic split $\log n - L$ and $L - \log n$. The public reference checker recomputes these exact endpoints, verifies the full prime-power list, checks both one-sided arithmetic row prices and checks $W/2 \leq W_{\text{low}} \leq W$ on every cell using the complete theta brackets. The new diagonal admission satisfies

$$\alpha \leq a(k_N) - \frac{9L^3}{\pi^4(N + 1/2)^3} \{a(k_N) - a(k_0)\} + m_0.$$

Every cell has the exactly represented positive reference $d_j = \kappa w_j + \alpha - r_{\text{up},j} > 1/200$. This retains the infinite receiver before any finite sign test.

S21.2 All constraints, mixed actions and numerical sign

With all 660 columns $F = [p, C]$, the full weighted Gram $G = F^*d^{-1}F$ has 435,600 entries. The exactly represented upper dyadic conditioner has denominator 2^{256} and positive diagonal. Its acceptance uses the complete directed congruence, not the midpoint factorization used to propose it:

$$\|R^*GR - I\| < 4.747 \times 10^{-74} < 10^{-65} < 10^{-60}.$$

The final signed inverse bound conservatively uses $\delta = 10^{-60}$. An inconclusive direct interval factorization at an earlier proposal pivot is not used as a negative-Gram inference or as a reason to discard a constraint direction.

The complete continuous mixed integrals use the elementary primitives of Section S20, retaining all five scalar weights, the pole/action row, and all sine, cosine and Green cross terms. For this larger list, integer and half-integer phases at each endpoint are evaluated directly in interval sine and cosine. This replaces a phase recurrence that can inflate enclosure radii; it does not replace the integrals by a finite output projection or numerical quadrature.

The scientific data contain B_E, K_E, H_s, P_E and the full response blocks for 660 primitive inputs. The original ridge cancels symbolically in $E = \hat{S}_{24} - d$. Assemble the lower $A_{24} = H^*RR^*H/(1 + \delta) - B_E - K_E$ as in the main proof. All 659 pivots of $P_C^*A_{24}P_C$ are strictly positive in directed arithmetic. No additional 64 receiving trials are present in this case. The raw core Gram, all constraint rows and the analytic invertible conditioner are retained.

S21.3 Portable scientific inputs and checks

The M24 companion archive contains four scientific JSON inputs and `INPUTS.json`. Decimal interval strings and exact dyadic integers are retained losslessly. Extract this archive into `data/worked_example_m24/` in the principal source package. The separate `code/worked_example_m24/` programs use only this chosen data folder and a reader-chosen output path:

```
python code/worked_example_m24/check_reference.py -data data/worked_example_m24
python code/worked_example_m24/check_weighted_gram.py -data
data/worked_example_m24
python code/worked_example_m24/compute_signed.py -data data/worked_example_m24
-output signed_m24_rebuilt.json
python code/worked_example_m24/check_signed.py -data data/worked_example_m24
```

The first program checks the whole reference and theta floor. The second regenerates the complete weighted Gram from the reference cells and receives the exact dyadic conditioner. The third reconstructs the signed lower from the supplied full mixed moments. The fourth independently receives the saved lower, reconstructs the analytic core conditioner, and replays completed-column LDL with a different multiplication grouping. `generate_mixed.py` regenerates elementary moments for a requested finite input list; its `-inputs`, `-bits` and output options are documented in the package README. A small-list cross-check, full signed replay and complete source regeneration have different scopes. The general theorem remains analytic; these interval checks certify the stated finite instance within their declared source and error contracts.

S22 Arithmetic certificate transfers in original profile coordinates

These results strengthen the paper's arithmetic verification framework through complete-core representation, certified finite-matrix error bounds, and quantitative transfer of supported lower bounds. They establish a complete-core pullback, a polar-completion theorem under the shared zero-free-strip hypothesis, and a sharper local support-extension theorem. Together, they expand the supporting tools available for the paper's arithmetic matrix criteria while preserving the original operators, norms, constraints and complete mixed terms.

Each subsection states its own support and arithmetic assumptions. The complete-core construction concerns the prescribed DN matrix of Theorem L.6; the polar completion assumes a supplied polar-free raw floor; the local extension starts from a positive whole arithmetic-only bound. These give three distinct interfaces in the same unweighted profile realization.

S22.1 Complete-core representation and certified finite pullbacks

The packet estimates above concern selected separated families. We now construct a complete representation of the ordinary DN core used in Theorem L.6. The representation includes low frequencies, both local parity channels and one polar direction. It supplies a second way of evaluating the same finite matrix, with a paid induced error. Throughout this subsection $L \geq 4$; this restriction does not apply to the other transfer results below.

For a full-line profile set

$$\mathcal{E}(v) = \|v\|_2^2 + \int_{\mathbb{R}} a(\xi) |\widehat{v}(\xi)|^2 \frac{d\xi}{2\pi}, \quad \|v\|_{\mathcal{E}} = \mathcal{E}(v)^{1/2}. \quad (\text{S74})$$

The symbol is the one in the main paper. Its exact paired representation is

$$\mathcal{E}(v) = \|v\|_2^2 + \frac{1}{2} \iint_{\mathbb{R}^2} J(x-y) |v(x) - v(y)|^2 dx dy, \quad J(t) = \frac{e^{-|t|/2}}{1 - e^{-2|t|}}.$$

Indeed $\int_{\mathbb{R}} (1 - \cos(\xi t)) J(t) dt = a(\xi)$ by the positive digamma series and Tonelli's theorem. Also $1 + a(\xi) \asymp \log(2 + |\xi|)$ by the symbol bounds in the main paper. These are unweighted profile coordinates, not the original h norm.

Lemma S22.1 (Compact polar localization and complete frame). *There are fixed smooth compact mothers and a fixed lattice $\omega_k = \pi k/5$, $k \in \mathbb{Z}$, with packets*

$$g_k(u) = -\frac{(\partial_u^2 - 1/4)[\theta(u)e^{i\omega_k u}]}{1 + \omega_k^2}, \quad \ell_k = \log(2 + |\omega_k|),$$

having the following properties. Every smooth profile v supported in $(-L, L)$ with both exponential moments zero has an exact expansion

$$v = \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}} d_{jk} g_k(\cdot - j), \quad \sum_{j,k} \ell_k |d_{jk}|^2 \leq C \mathcal{E}(v).$$

Only $|j| < L+2$ occur. Synthesis is bounded from the entire weighted coefficient space to the energy space:

$$\mathcal{E} \left(\sum_{j,k} d_{jk} g_k(\cdot - j) \right) \leq C \sum_{j,k} \ell_k |d_{jk}|^2.$$

Analysis, synthesis and their identity extend to the minimal energy closure of these profiles. Individual atoms have both moments zero but need not be odd. All k , including zero, are retained.

Proof. Let $P = \partial_u^2 - 1/4$. Its inverse has kernel $-e^{-|u|/2}$, because $Pe^{-|u|/2} = -\delta_0$. Thus $h = P^{-1}v$ has the same support as v : to the right of its support it equals $-e^{-u/2} \int e^{x/2} v(x) dx = 0$, and the other moment treats the left. Fourier multiplication gives

$$\sum_{r=0}^2 \mathcal{E}(h^{(r)}) \leq C\mathcal{E}(v).$$

Fix an even smooth partition $\sum_j \eta(u-j) = 1$, with η supported in $(-2, 2)$. One explicit choice is to divide $b(u/2)$ by its positive periodic translate sum, where $b(t) = e^{-1/(1-t^2)}$ for $|t| < 1$ and zero otherwise. Fix an even $\theta \in C_c^\infty(-4, 4)$ equal to one on $[-2, 2]$. Such a cutoff can be specified by the normalized integral of the same bump. All support, derivative and overlap constants below are fixed.

Write $h_j(u) = \eta(u-j)h(u)$ and $v_j = Ph_j$. Each v_j has both moments zero by integration by parts, and $\sum_j v_j = v$ exactly. For any bounded-overlap family of smooth translates b_j , the paired jump formula gives

$$\sum_j \mathcal{E}(b_j z) \leq C\mathcal{E}(z).$$

To see this, split each difference into $b_j(x)(z(x) - z(y)) + (b_j(x) - b_j(y))z(y)$ and use $\sum_j |b_j(x) - b_j(y)|^2 \leq \min(C, C|x-y|^2)$. Its integral against $J(x-y)$ is finite, both near zero and at infinity. Applying this to $\eta_j, \eta'_j, \eta''_j$ and the preceding derivative bound proves both $\sum_j \sum_{r=0}^2 \mathcal{E}(h_j^{(r)}) \leq C\mathcal{E}(v)$ and $\sum_j \mathcal{E}(v_j) \leq C\mathcal{E}(v)$.

Expand $h_j(j+u) = \sum_k c_{jk} e^{i\omega_k u}$ on the fixed period $(-5, 5)$. It vanishes near the period boundary, and $\theta h_j = h_j$. Consequently $d_{jk} = -(1 + \omega_k^2)c_{jk}$ gives the stated expansion. For completeness, periodizing J gives Fourier eigenvalues $a(\omega_k)$. Its zero-image term is controlled by full-line energy; all other images are uniformly bounded on the support $[-2, 2]$ and contribute at most a fixed multiple of the squared L^2 norm. Apply this comparison to h_j, h'_j, h''_j and use periodic Parseval. Since $(1 + \omega^2)^2 \asymp 1 + \omega^2 + \omega^4$, it proves the weighted coefficient bound, including the zero mode.

We also need synthesis, not only an analysis inequality. At one cell put $z = -\sum_k d_k e^{i\omega_k u} / (1 + \omega_k^2)$. Periodic Parseval gives

$$\sum_{r=0}^2 \mathcal{E}_{\text{per}}(z^{(r)}) \leq C \sum_k \ell_k |d_k|^2.$$

For any fixed smooth b supported away from the period endpoints, $\mathcal{E}(bz) \leq C_b \mathcal{E}_{\text{per}}(z)$. Inside the period this follows from the same difference split, because $J(x-y) \leq \sum_m J(x-y+10m)$. For pairs with one point outside it, the fixed gap between $\text{supp } b$ and its endpoints makes the exterior integral uniformly bounded by $C_b \|z\|_2^2$. Apply this to $\theta, \theta', \theta''$ in $P(\theta z)$. The translated outputs have a fixed support overlap. At a pair (x, y) , at most twice that overlap many differences are nonzero; Cauchy-Schwarz before the paired jump integral pays their entire sum. This proves the full synthesis bound for finite lists and then for their weighted closure. The analysis and synthesis maps are bounded, so their identity on smooth tests extends to the stated minimal closure. Neither a lower Riesz bound for arbitrary redundant coefficients nor a high-frequency restriction has been used. $\square$

Lemma S22.2 (A uniformly conditioned smooth polar direction). *Fix a nonzero even nonnegative $\phi \in C_c^\infty(-\rho, \rho)$, $r > \rho > 0$, and $L \geq r + \rho + 1$. In full odd coordinates put*

$$p_L(x) = \sinh(x/2), \quad u_L(x) = \phi(x - (L - r)) - \phi(x + (L - r)).$$

Then u_L is supported in $(-L, L)$, $\|u_L\|_2^2 = 2\|\phi\|_2^2$, and

$$\frac{|\langle p_L, u_L \rangle|}{\|p_L\|_2 \|u_L\|_2} \geq c_{\phi, r} > 0, \quad \mathcal{E}(u_L) + \|a(D)u_L\|_2^2 \leq C_{\phi, r}.$$

Every supported odd profile in the energy domain splits exactly as $v = v_0 + \alpha u_L$, with $v_0 \perp p_L$ and $\mathcal{E}(v_0) + |\alpha|^2 \leq C\mathcal{E}(v)$.

Proof. Set $A_\phi = \int \cosh(t/2)\phi(t) dt > 0$. Evenness gives $\langle p_L, u_L \rangle = 2A_\phi \sinh((L-r)/2)$, while $\|p_L\|_2^2 = \sinh L - L \leq e^L/2$. Since $L-r \geq 1$, the ratio is bounded below by $A_\phi e^{-r/2}(1-e^{-1})/\|\phi\|_2$. The two arms are disjoint. Translation invariance and the triangle inequality give $\mathcal{E}(u_L) \leq 4\mathcal{E}(\phi)$ and $\|a(D)u_L\|_2 \leq 2\|a(D)\phi\|_2$; smoothness makes both finite. Take $\alpha = \langle p_L, v \rangle / \langle p_L, u_L \rangle$. The angle bound controls $|\alpha|$ by $C\|v\|_2$; the energy triangle inequality proves the assertion. The seed is smooth in the genuine supported operator domain, and the polar functional is continuous in L^2 , so the split is valid throughout the closed local domain. This is geometric conditioning only, with no signed form assertion. $\square$

Use $\rho = 1/4$, $r = 1$ from now on. Let $J_L : L^2(0, L) \rightarrow L^2(\mathbb{R})$ be the odd isometry given by $J_L v(x) = v(x)/\sqrt{2}$ on $(0, L)$, its odd reflection on $(-L, 0)$, and zero elsewhere. Apply the frame lemma to the polar-free part of $J_L v$ and keep the single exact seed coordinate. This covers the full ordinary DN core, rather than its polar-free intersection. Indeed that intersection alone cannot approximate the whole core at fixed support: if P_N is the ordinary DN projection and $p = \sinh(x/2)$ on $(0, L)$,

$$\text{dist}\left(\frac{P_N p}{\|P_N p\|}, p^\perp\right) = \frac{\|P_N p\|}{\|p\|} \rightarrow 1.$$

Proposition S22.3 (Exact host and complete energy control). *Fix the host buffer $h_* = 9$. On all full-line profiles supported in $(-L - h_*, L + h_*)$, not merely odd individual atoms, define*

$$\hat{q}_L = D_{\text{full}} + (m_0 + 2)I - C_{L+h_*}^{\text{ar}} - J_L R_L J_L^*.$$

Here J_L^* includes restriction and odd projection, and R_L is the old complete Green response. Then

$$\hat{q}_L[J_L v, J_L w] = \langle v, \mathcal{A}_{L,2} w \rangle \quad (\text{S75})$$

as closed forms, and

$$|\hat{q}_L[f, g]| \leq F_L \mathcal{E}(f)^{1/2} \mathcal{E}(g)^{1/2}, \quad F_L = 12 + 12e^{L+h_*}.$$

The complete frame with its seed converges in this same host form. For the paired arithmetic operator on this host, define $\text{Err}(f, g) = \langle f, (C_{L+h_*}^{\text{ar}} - C_{L+h_*}^{\text{main}})g \rangle$, where $C_{L+h_*}^{\text{main}}$ has kernel $e^{|x-y|/2}$ restricted to the same host. Thus the discrepancy includes both shift orientations and the complete continuous main term. Under the strip premise of Lemma S19.1, it has the bound

$$|\text{Err}(f, g)| \leq C e^{2dL} \mathcal{E}(f)^{1/2} \mathcal{E}(g)^{1/2}, \quad d = b - 1/2 + \eta, \quad (\text{S76})$$

for represented old profiles. The factor e^{2dL} records the support dependence of this arithmetic error bound.

Proof. Nonzero cells satisfy $|j| < L+2$, and their atoms have support within four of their cell centers. Thus all partial syntheses lie in the stated fixed host. Infinite synthesis equals the original profile, although finite partials can occupy its collar. On old profiles every additional host arithmetic shift

has zero overlap. The exact original realization gives $D_{\text{full}}|_{J_L L^2(0,L)} = J_L(D_L + R_L)J_L^*$ as forms. Subtracting the transported old response therefore proves (S75). Replacing it by the host response would change the target.

The full paired arithmetic norm is bounded by $2 \sum_{n < e^{2(L+h_*)}} \Lambda(n)/\sqrt{n} < 12e^{L+h_*}$, using Lemma M.1. Also $|m_0 + 2| < 8$ and $\|J_L R_L J_L^*\| \leq \pi/2$ by Lemma B.1. Cauchy–Schwarz for the nonnegative multiplier D_{full} and $\|f\|_2 \leq \mathcal{E}(f)^{1/2}$ prove the stated F_L bound. It includes all mixed, collar and parity terms. Energy convergence thus implies convergence in the host form.

Here are the additional uniformities needed for the last assertion. Factoring out $e^{i\omega_k u}$ in g_k leaves

$$\frac{(\omega_k^2 + 1/4)\theta - \theta'' - 2i\omega_k \theta'}{1 + \omega_k^2},$$

whose fixed-order smooth seminorms are bounded uniformly for every $k \in \mathbb{Z}$, including zero. In a pairing of cells j, l , integrate by parts in the internal variable as in the proof of Proposition S19.2. For every fixed m this bounds each window seminorm by $C_m(1 + |\omega_k - \omega_{k'}|)^{-m}$. Lemma S19.1 then bounds the corresponding entry by

$$C_m e^{d|j-l|} \ell_k (1 + |\omega_k - \omega_{k'}|)^{-m}.$$

The continuous term here is the actual positive-shift integral over $y \geq 1$. When the translated window meets the origin, replacing its full Mellin main by this truncated main costs only a fixed compact-range term with the same smooth window bound. A window strictly to the negative side has zero actual sum and integral. The finitely wide origin range is bounded directly, retaining its carrier phases. The opposite shift orientation is paid in the same manner. Thus no fictitious primes below one, discarded orientation or unpriced negative window occurs. The logarithmic ratio of ℓ_k and $\ell_{k'}$ is absorbed by one extra inverse separation power. Since the full signed lattice has fixed spacing, the two-sided Schur test yields a weighted coefficient norm bound $Ce^{d|j-l|}$ for each cell pair. The seed adds two translates of one fixed smooth mother at carrier zero; its pairings satisfy the same argument with bounded multiplicity. It need not have zero polar moment. Finally

$$\sup_{|j| < L+2} \sum_{|l| < L+2} e^{d|j-l|} \leq C_d e^{2dL}.$$

The same estimate holds for the two additional seed centers after changing the fixed constant. Schur’s test on cells and the complete bounded analysis map prove (S76) for finite syntheses. Pass to the energy limit in the bounded host arithmetic form. All color, low-frequency, seed and spatial cross terms have been retained. This is not a lower Riesz assertion for the redundant dictionary. $\square$

Proposition S22.4 (Quantitative finite-core pullback). *Let $U_N : \mathbb{C}^N \rightarrow L^2(0, L)$ be the ordinary first- N DN isometry, $k_{\max} = k_{N-1}$, and*

$$M_N^2 = 1 + a(k_{\max}) + \pi/2.$$

Retain every active cell, every local frequency $|\omega_k| \leq \Omega$, and the exact seed. The resulting synthesis map S_Ω from $\mathbb{C}^N$ to the common host satisfies, for $\Omega \geq 2$,

$$\|S_\Omega - J_L U_N\|_{\mathbb{C}^N \rightarrow \mathcal{E}}^2 \leq C_{\text{cov}} \Omega^{-1/4} N(1 + k_{\max})^{1/2}. \quad (\text{S77})$$

Its pullback $H_\Omega = S_\Omega^ \hat{q}_L S_\Omega$ approximates the same core matrix $H_N = U_N^* \mathcal{A}_{L,2} U_N$ with*

$$\|H_\Omega - H_N\| \leq \eta_\Omega := F_L \epsilon (2M_N + \epsilon) \quad (\text{S78})$$

whenever the left map norm in (S77) is at most ϵ . The fixed C_{cov} is independent of L, N, Ω ; a numerical use must certify it or directly certify that map error.

Proof. A normalized zero-extended DN sine has, for $s = 1/4$,

$$\|J_L e_q\|_{H^s(\mathbb{R})}^2 \leq C_s(1 + k_q)^{2s}, \quad L \geq 4.$$

This follows by expressing its two arms as interval indicators times modulations. The squared fractional seminorm of a normalized interval indicator is $C_s L^{-2s}$; modulation costs at most $C_s(1 + k_q)^{2s}$. The triangle and Cauchy–Schwarz inequalities give

$$\|J_L U_N z\|_{H^s}^2 \leq C_s N(1 + k_{\max})^{1/2} \|z\|^2.$$

The polar-seed subtraction preserves this bound because its coefficient is bounded by $C\|z\|$ and its translated smooth H^s norm is fixed.

The analysis proof in Lemma S22.1 works with H^s in place of logarithmic energy: replace $J(t)$ by $|t|^{-1-2s}$ in the difference estimates. Its small-distance error integral converges for $s < 1$, and its exterior/periodic image integrals converge for $s > 0$. The multiplier P^{-1} gains two derivatives in this norm as well. Periodic Parseval therefore gives

$$\sum_{j,k} (1 + \omega_k^2)^s |d_{jk}|^2 \leq C_s \|v_0\|_{H^s}^2.$$

For $|\omega_k| > \Omega \geq 2$, $\ell_k/(1 + \omega_k^2)^{1/4} \leq C\Omega^{-1/4}$. Apply bounded energy synthesis to precisely these discarded coefficients to prove (S77). Finite DN zero extensions belong to the required minimal energy closure: cut off their boundary strips and mollify. For any $0 < t < 1/2$ the strips of a fixed piecewise smooth vector have H^t norm tending to zero, hence energy norm tending to zero. Correcting the one polar moment by the fixed smooth seed preserves this limit. This verifies the actual input domain rather than assuming smooth DN endpoint values.

By the original Green identity and Lemma B.1, $\|J_L U_N\|_{\mathcal{C}^N \rightarrow \mathcal{E}} \leq M_N$. Write $S_\Omega = J_L U_N + E_\Omega$ and expand the full form pullback into two mixed terms and one error square. The host continuity bound gives $2F_L M_N \epsilon + F_L \epsilon^2$, proving (S78) as an induced matrix estimate, without an additional entrywise dimension fee. $\square$

Corollary S22.5 (Certified acceptance on the original core range). *Write the finite synthesis as a dictionary map times its actual coefficient map, $S_\Omega = \mathcal{D}_\Omega A_\Omega$, retaining the seed row. Let G_Ω be the complete host-form dictionary Gram, so $H_\Omega = A_\Omega^* G_\Omega A_\Omega$. If a Hermitian center C_Ω has certified induced error r_Ω to this exact pullback and*

$$C_\Omega - (r_\Omega + \eta_\Omega)I \succeq I/5,$$

then $H_N \succeq I/5$. At every cutoff admitted by Theorem L.6, it follows that $\mathcal{A}_{L,2} \succeq 52I/405$. Under RH, each fixed admitted L, N can be recognized in this representation by refinement.

Proof. The first assertion is (S78) and the directed center error. Apply Theorem L.6 only at its prescribed cutoff and with its actual domain and commutator hypotheses. Under RH that theorem gives $H_N \succ 2I/5$. Taking $\eta_\Omega, r_\Omega \leq 1/40$ makes the displayed lower exceed $(2/5 - 2\eta_\Omega - 2r_\Omega)I \succeq 3I/10$. For example $\epsilon \leq \min\{1, [80F_L(M_N + 1)]^{-1}\}$ suffices for $\eta_\Omega < 1/40$. The two comparison losses of each error are retained. Increasing Ω and the numerical precision supplies conditional fixed-instance recognition, not a successful cofinal sign.

All finite coefficients are effectively enclosable. The specified smooth bumps and their normalized integrals have effective derivative and support bounds. Analysis coefficients use their integrals and the explicit compact P^{-1} kernel. Arithmetic entries involve only the finite active prime-power

list and smooth overlap integrals. For D_{full} use Fourier integrals; integration by parts bounds the omitted Fourier tails by any chosen inverse power, while the symbol grows logarithmically. The old response entries can also be enclosed without a false norm limit: for a bounded old half-profile w , $|\langle r_j, w \rangle| \leq \|w\|_\infty / c_j$, so its paired scalar series tail is bounded by the product of the two suprema times $c_j^{-2} + (2c_j)^{-1}$. Finite directed quadrature and all coefficient, coordinate and Gram errors must then be included in r_Ω . A numerical receipt cannot replace C_{cov} by an unspecified existence constant. $\square$

This criterion tests the pullback on the actual original-core input range. Positivity of G_Ω on arbitrary redundant dictionary coefficients is neither assumed nor required. A sufficient cutoff is $\Omega \geq C_{\text{cov}}^4 \epsilon^{-8} N^4 (1 + k_{\text{max}})^2$. There are $O((L+1)(1+\Omega))$ dictionary atoms and one seed coordinate. The bound is polynomial in N, e^L, L and the inverse fixed margin; on the elementary near-threshold CO schedule it is $\exp(O(e^L)) = \exp(O(\sqrt{\log M}))$. It counts dictionary size, not arithmetic operations or bit complexity, and native H_N evaluation can be smaller. The representation transfers a certified finite sign through the displayed complete error budget; the arithmetic discrepancy retains its explicit support-dependent factor.

S22.2 Conditional completion of the polar direction

The preceding polar-free constructions admit an exact one-dimensional completion on the original odd supported space. The estimate below retains the raw prime operator and expresses the additional debit in the same unweighted profile norm. Two arithmetic hypotheses give alternative bounds for that debit.

Supported operators and a smooth polar seed. Let $\mathcal{H}_L = L_{\text{odd}}^2(-L, L)$, with profiles extended by zero outside $(-L, L)$, and let

$$(J_L g)(x) = 2^{-1/2} \text{sgn}(x) g(|x|)$$

be the unitary map from the odd-half coordinates to $\mathcal{H}_L$. Write $\mathcal{D}_L$ for the operator associated with the supported closed form of the full-line multiplier $a(D)$ on its minimal odd domain $\mathcal{V}_L$. The complete supported arithmetic action is

$$(\mathcal{C}_L f)(x) = \sum_{2 \leq n < e^{2L}} \frac{\Lambda(n)}{\sqrt{n}} [f(x + \log n) + f(x - \log n)], \quad |x| < L.$$

Both channels use the zero extension of the same odd profile. Define

$$\tilde{B}_L = \mathcal{D}_L + m_0 I - \mathcal{C}_L, \quad \mathcal{R}_L = J_L R_L J_L^*, \quad \tilde{\mathcal{A}}_{L,c} = J_L \mathcal{A}_{L,c} J_L^* = \tilde{B}_L - \mathcal{R}_L + cI,$$

where c is fixed. Here $0 \leq \mathcal{R}_L \leq (\pi/2)I$ by Lemma B.1, and $\mathcal{D}_L = J_L(D_L + R_L)J_L^*$ as forms. These raw operators have the common form domain $\mathcal{V}_L$.

Fix $0 < \rho < r$ and a nonzero, nonnegative, real even $\phi \in C_c^\infty(-\rho, \rho)$. For $L \geq r + \rho + 1$ put

$$a_L = L - r, \quad p_L(x) = \sinh(x/2), \quad u_L(x) = \phi(x - a_L) - \phi(x + a_L), \quad m_L = \langle p_L, u_L \rangle.$$

Lemma S22.2 gives $m_L \asymp e^{L/2}$, a uniform moment angle, and the uniform seed-action bound $\|a(D)u_L\|_2 \leq 2\|a(D)\phi\|_2$. In particular u_L belongs to the operator domains of both raw supported operators. For every $f \in \mathcal{V}_L$,

$$\alpha = \frac{\langle p_L, f \rangle}{m_L}, \quad f_0 = f - \alpha u_L \perp p_L, \quad \|f_0\|_2 + |\alpha| \leq C_\phi \|f\|_2. \quad (\text{S79})$$

The split preserves the closed form domain, since the local polar functional is L^2 -continuous and the seed is smooth and compact.

The raw main kernel and scalar completion. Let $\mathcal{C}_L^{\text{main}}$ be the supported integral operator with kernel $e^{|x-y|/2}$, the continuous $y \geq 1$ counterpart of the paired arithmetic shifts. On the full odd space,

$$\mathcal{C}_L^{\text{main}} = -2p_L \otimes p_L - \mathcal{K}_L, \quad 0 \preceq \mathcal{K}_L \preceq 4I, \quad (\text{S80})$$

where $\mathcal{K}_L$ has kernel $e^{-|x-y|/2}$ and $(p_L \otimes p_L)f = p_L \langle p_L, f \rangle$. Indeed $e^{|t|/2} = 2 \cosh(t/2) - e^{-|t|/2}$, and the even $\cosh(x/2) \cosh(y/2)$ channel vanishes on odd inputs. Positivity and the bound for $\mathcal{K}_L$ follow from its full-line Fourier multiplier $(\xi^2 + 1/4)^{-1}$. The signed-pole geometric counterpart is $\tilde{B}_L - 2p_L \otimes p_L$; its restriction to $p_L^\perp$ agrees with the raw form.

Lemma S22.6 (Completion from a supported seed residual). *For $\mathcal{T}_L = \tilde{B}_L$ or $\mathcal{T}_L = \tilde{\mathcal{A}}_{L,c}$, suppose*

$$\|(\mathcal{C}_L - \mathcal{C}_L^{\text{main}})u_L\|_{L^2(-L,L)} \leq \varepsilon(L), \quad \varepsilon(L) \geq 1, \quad \varepsilon(L) = o(e^L).$$

Then, for all sufficiently large L , every $f = f_0 + \alpha u_L$ as in (S79) satisfies

$$q_{\mathcal{T}_L}[f] \geq q_{\mathcal{T}_L}[f_0] - Ce^{-L}\varepsilon(L)^2\|f_0\|_2^2.$$

In particular, if $b_0(L) \geq 0$ and $q_{\mathcal{T}_L}[g] \geq -b_0(L)\|g\|_2^2$ for all $g \in \mathcal{V}_L \cap p_L^\perp$, then

$$q_{\mathcal{T}_L}[f] \geq -C[b_0(L) + e^{-L}\varepsilon(L)^2]\|f\|_2^2, \quad f \in \mathcal{V}_L. \quad (\text{S81})$$

All constants may depend on the fixed seed and shift, but not on L .

Proof. Put $e_L = (\mathcal{C}_L - \mathcal{C}_L^{\text{main}})u_L$. Equation (S80) gives

$$\tilde{B}_L u_L = 2m_L p_L + (\mathcal{D}_L + m_0 I + \mathcal{K}_L)u_L - e_L.$$

The non-rank-one remainder has norm at most $C\varepsilon(L)$. The same assertion holds for $\tilde{\mathcal{A}}_{L,c}$ by the uniform response bound. Therefore

$$\langle u_L, \mathcal{T}_L u_L \rangle = 2m_L^2 + O(\varepsilon(L)) \geq c_0 e^L, \quad |\langle f_0, \mathcal{T}_L u_L \rangle| \leq C\varepsilon(L)\|f_0\|_2.$$

The first inequality holds eventually because $\varepsilon(L) = o(e^L)$; the second uses $f_0 \perp p_L$. Completing the scalar square in $q_{\mathcal{T}_L}[f_0 + \alpha u_L]$ gives the first conclusion. The nonnegative base debit and the uniform conditioning in (S79) give (S81). The operator-domain seed justifies the mixed pairing throughout $\mathcal{V}_L$. $\square$

Proposition S22.7 (Completion under a zero-strip hypothesis). *Assume every nontrivial zeta zero has real part at most b , where $1/2 \leq b < 1$. Fix $0 < \eta < 1 - b$ and put $d = b - 1/2 + \eta$. If a nonnegative $b_0(L)$ gives the polar-free lower bound in Lemma S22.6, then*

$$q_{\mathcal{T}_L}[f] \geq -C[b_0(L) + e^{(4b-3+4\eta)L}]\|f\|_2^2 \quad (f \in \mathcal{V}_L) \quad (\text{S82})$$

for $\mathcal{T}_L = \tilde{B}_L$ and $\tilde{\mathcal{A}}_{L,c}$, respectively, using each operator's own polar-free lower bound.

Proof. For real T , define the exact truncated-main discrepancy

$$F_\phi(T) = \sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} \phi(\log n - T) - \int_1^\infty y^{-1/2} \phi(\log y - T) dy.$$

For $T \geq \rho$ the main integral is the full Mellin residue, so Lemma S19.1 at $\omega = 0$ gives $|F_\phi(T)| \leq Ce^{dT}$. For $0 \leq T \leq \rho$, the difference between the full residue and the truncated main integral is bounded

on a fixed compact interval. For $-\rho \leq T < 0$, both terms use only $0 \leq \log y \leq 2\rho$, a fixed finite arithmetic list and a bounded main integral. For $T < -\rho$ both terms vanish. Enlarging one constant therefore gives, for every real T ,

$$|F_\phi(T)| \leq C e^{d \max(T, 0)} \mathbf{1}_{\{T \geq -\rho\}}.$$

Evenness of ϕ gives the exact four-window identity

$$e_L(x) = F_\phi(a_L - x) - F_\phi(-a_L - x) + F_\phi(x - a_L) - F_\phi(x + a_L), \quad |x| < L.$$

This retains both translation directions and all active prime powers; the compact seed makes every shift beyond the supported cutoff vanish. In particular

$$|e_L(x)| \leq C [e^{d|x-a_L|} + e^{d|x+a_L|}].$$

The complete spatial norm is paid by

$$\int_{-L}^L e^{2d|x-a_L|} dx = \frac{e^{2d(L+a_L)} + e^{2d(L-a_L)} - 2}{2d} \leq C_{d,r} e^{4dL},$$

and the same bound holds for $x + a_L$. Hence $\|e_L\|_2 \leq C e^{2dL}$, with no factor depending on the receiving profile or on the interval length outside this exponential. Since $0 < 2d < 1$, this is an admissible $\varepsilon(L)$ in Lemma S22.6. Its squared cost divided by the raw seed reserve e^L gives (S82). $\square$

Proposition S22.8 (Completion under a prime-primitive bound). *Let $\psi_C(y) = \sum_{n \leq y} \Lambda(n)$ be the actual Chebyshev function. Suppose it obeys*

$$|\psi_C(y) - (y - 1)| \leq C_\sigma y^\sigma \quad (y \geq 1), \quad \frac{1}{2} < \sigma < 1, \quad \psi_C(1) = 0.$$

For either raw supported operator, a nonnegative polar-free debit $b_0(L)$ then gives

$$q_{\mathcal{T}_L}[f] \geq -C[b_0(L) + e^{(4\sigma-3)L}] \|f\|_2^2, \quad f \in \mathcal{V}_L, \quad (\text{S83})$$

for all sufficiently large L .

Proof. Set $E(y) = \psi_C(y) - (y - 1)$ and, for each fixed $|x| < L$,

$$G_x(y) = y^{-1/2} [u_L(x + \log y) + u_L(x - \log y)].$$

Stieltjes integration by parts gives exactly

$$e_L(x) = \int_{[1, \infty)} G_x(y) dE(y) = - \int_1^\infty E(y) G'_x(y) dy.$$

Here $E(1) = 0$ removes the lower boundary, and compact support removes the upper boundary. Only the fixed smooth seed is differentiated: $|G'_x(y)| \leq C_\phi y^{-3/2}$ on at most four fixed-log-width bands, whose nonnegative centers lie near $|x - a_L|$ and $|x + a_L|$. Bands meeting $\log y = 0$ contribute a bounded constant. With $\gamma = \sigma - 1/2 > 0$, integration on these bands gives

$$|e_L(x)| \leq C [e^{\gamma|x-a_L|} + e^{\gamma|x+a_L|}].$$

The spatial integral used above, with d replaced by γ , yields $\|e_L\|_2 \leq C e^{(2\sigma-1)L}$. Since $\sigma < 1$, the shared scalar completion applies and proves (S83). $\square$

The primitive hypothesis with $\sigma < 3/4$ gives a vanishing additional debit, and at $\sigma = 3/4$ a bounded additional debit. Under the zero-strip hypothesis, $b < 3/4$ permits $\eta < 3/4 - b$ and again a vanishing debit; at $b = 3/4$, each fixed $\eta > 0$ gives the explicit factor $e^{4\eta L}$. The constants may depend on the chosen positive η , so this endpoint statement retains that factor. For every allowed $\sigma < 1$ or $b + \eta < 1$, the additional debit is $o(e^L)$. These completions transport a separately supplied polar-free lower bound to the full raw supported operator, retaining its original unweighted norm and the exact polar main term. The unitary identity $\|J_L g\|_2 = \|g\|_2$ transports the same debit to $\mathcal{A}_{L,c}$ in its odd-half norm. In the theta-congruent coordinates $f = sh$, the norm factor is exactly $\|sh\|_2^2$; the bound $\|s\|_\infty < 3$ of the main paper further gives $\|sh\|_2^2 \leq 9\|h\|_2^2$ whenever those original supported coordinates are used.

S22.3 Actual-prime thin collars and local support extension

We work in the ordinary odd-half profile space. Here $0 < \ell < U$, $J = (\ell, U)$ and $\delta = U - \ell$. The support hypotheses in this subsection are independent of those in the full-core construction. The physical old-interval/collar decomposition also occurs in the support-variation analysis of [4, Section 4]. The following estimate uses the integer spacing of the actual prime channels for the arithmetic-only recipient of this paper.

Lemma S22.9 (Complete thin-collar arithmetic payment). *If $\delta e^{2U} \leq 1/4$, the complete paired arithmetic action satisfies*

$$\|1_{(0,\ell)} C_U^{\text{ar}} 1_J\|^2 \leq 2 \sum_{n < e^{2U}} \frac{\Lambda(n)^2}{n} \leq 24U^2 + 12U, \quad \|1_J C_U^{\text{ar}} 1_J\| \leq 2Ue^{-\ell}.$$

Proof. Put $t_n = \log n$ and $w_n = \Lambda(n)/\sqrt{n}$. On a collar input u , the old-output block is exactly

$$\sum_{n < e^{2U}} w_n [u(x + t_n) 1_J(x + t_n) - u(t_n - x) 1_J(t_n - x)], \quad 0 < x < \ell.$$

The ordinary $x - t_n$ arm cannot reach J . For fixed x , the two displayed channels require integers in $(e^{\ell-x}, e^{U-x})$ and $(e^{\ell+x}, e^{U+x})$, respectively. Each interval has length less than one, since its length is at most $\delta e^{2U} \leq 1/4$. Thus at most two scalar summands occur. Pointwise Cauchy–Schwarz pays their entire cross term by twice the sum of squares. For each n , the two changes of variables give the disjoint source pieces $J \cap (t_n, t_n + \ell)$ and $J \cap (t_n - \ell, t_n)$. Their squared input integrals sum to at most $\|u\|^2$. Summation proves the first inequality with every prime power retained.

The proof of Lemma M.1 gives $\Psi(X) < 3X$ for $X > 1$. Partial summation therefore gives

$$\sum_{n \leq X} \frac{\Lambda(n)}{n} \leq 3(1 + \log X), \quad \sum_{n \leq X} \frac{\Lambda(n)^2}{n} \leq 3 \log X (1 + \log X).$$

Take $X = e^{2U}$. For the collar diagonal, $\delta < \log 2$ excludes ordinary shifts between collar points. A reflected channel requires $n \in (e^{2\ell}, e^{2U})$, an interval of length at most $2\delta e^{2U} \leq 1/2$. There is at most one such integer. Its coefficient is at most $2Ue^{-\ell}$, and its compressed reflection is a self-adjoint contraction. This proves the remaining bound, including activation endpoints, whose overlaps have measure zero. $\square$

Proposition S22.10 (Paid local support extension). *Let $c \geq 0$ be fixed and suppose the entire arithmetic-only operator $\mathcal{A}_{\ell,c} \succeq \mu I$, $\mu > 0$. Set $\bar{U} = \ell + 1$ and*

$$B_{\text{mix}} = 2 + 3\pi/2 + \sqrt{24\bar{U}^2 + 12\bar{U}}, \quad w_* = 2\bar{U}e^{-\ell},$$

$$\tau = \max(1, 2B_{\text{mix}}^2/\mu),$$

$$T_* = \tau + |m_0| + w_* + 1.$$

Choose any positive δ with

$$\delta \leq \min\{\ell/16, 1, e^{-2\bar{U}}/4, (\pi/9)e^{-2T_*}\}.$$

For every $\ell \leq L \leq \ell + \delta$, on the complete original odd-half domain,

$$\mathcal{A}_{L,c} \succeq \frac{\min(\mu/2, \tau)}{(1 + B_{\text{mix}}/\tau)^2} I \succ 0.$$

For $\ell \geq 1$ and $0 < \mu \leq 1$, a safe dyadic choice may be made with $\log(1/\delta) = O((\ell+1)^2/\mu + \ell)$, with fixed symbol constants.

Proof. The full raw form is compatible under physical zero extension. The true Green features in (45) give, on the old interval,

$$R_\ell - R_L|_{(0,\ell)} = \sum_{j \geq 0} 4 \left(\frac{1}{e^{2c_j \ell} + 1} - \frac{1}{e^{2c_j L} + 1} \right) \sinh(c_j \cdot) \otimes \sinh(c_j \cdot) \succeq 0.$$

The order holds first for positive partial sums and then for the bounded strong limits. Extra arithmetic shifts have zero overlap on the old support. Consequently the old compression of $\mathcal{A}_{L,c}$ is at least $\mathcal{A}_{\ell,c} \succeq \mu I$.

The spatial-cut argument at (52) preserves the original logarithmic domain and bounds the complete archimedean cross block by $2 + \pi$. Subtracting R_L , whose norm is at most $\pi/2$, gives a DN cross bound $2 + 3\pi/2$. Lemma S22.9 then bounds the complete old/collar cross block by B_{mix} , uniformly for all the stated L . The scalar shift has no cross block.

To bound the whole collar, choose an integer $n \in [\ell/(8\delta), \ell/(4\delta)]$. Its existence follows from $\delta \leq \ell/16$. On a vector f supported in (ℓ, L) , the first n DN modes satisfy

$$\|P_n f\|^2 \leq \frac{2n(L-\ell)}{L} \|f\|^2 \leq \frac{1}{2} \|f\|^2.$$

The positive symbol is increasing. Its defining digamma sum, compared with the integral of the decreasing function $2k^2/[t(t^2 + k^2)]$ on the mesh $t = 1/2 + 2j$, gives $a(k) \geq \frac{1}{2} \log(1 + 4k^2) \geq \log k$. Also $L \leq 17\ell/16$ and $k_n(L) \geq n\pi/L > \pi/(9\delta) > 1$. Hence the entire collar has

$$D_L[f] \geq \frac{1}{2} \log(\pi/(9\delta)) \|f\|^2.$$

Its arithmetic compression costs at most w_* by the same lemma. The chosen δ and $c \geq 0$ therefore make the whole collar block at least τI .

The old/collar split is valid on the same closed form domain by the spatial-cut proof just cited; the bounded Green and arithmetic terms preserve it. Write its blocks as A_{00}, C, T . Then $T \succeq \tau I$ and $A_{00} - CT^{-1}C^* \succeq (\mu - B_{\text{mix}}^2/\tau)I \succeq \mu I/2$. Complete square completion uses a triangular map whose inverse norm is at most $1 + B_{\text{mix}}/\tau$. This proves the displayed whole-operator bound, retaining all new directions and mixed terms. For the final assertion, $B_{\text{mix}} = O(\ell + 1)$, $w_* = O(1)$ and $\tau = O((\ell + 1)^2/\mu)$; the other logarithmic radius prices are $O(\ell)$ or smaller. A dyadic below the explicit minimum changes its logarithm by at most a fixed additive constant. $\square$

Corollary S22.11 (Local extension of the shift-two example). *Let $\ell = \frac{1}{2} \log(96 \log 2/\pi) + 1/10$ be the support of Theorem L.8. Throughout $\ell \leq L \leq \ell + 2^{-1300000}$, $\mathcal{A}_{L,2} \succeq I/1281$.*

Proof. The retained reference enclosures of the shift-two example give $\|C_\ell^{\text{ar}}\| < 7$, receiving floor greater than 1.182, and complete core Schur lower $I/10$. The whole triangular payment is therefore at most $(1 + 7)^2$, giving $\mathcal{A}_{\ell,2} \succeq I/640$. The exact support expression and $2/3 < \log 2 < 7/10$, $3 < \pi < 4$ give $1 < \ell < 37/20$. Since $\delta < 1/8 < 2 - \ell$, the proposed interval has $L < 2$, and one may sharpen the common budgets to $B_{\text{mix}} < 18$, $w_* \leq 4$ and $|m_0| < 6$. Take $\tau = 420000 > 2 \cdot 18^2 \cdot 640 = 414720$. For $\delta = 2^{-1300000}$, the thinness and $\ell/16$ guards hold, and

$$\frac{1}{2} \log(\pi/(9\delta)) - 10 > 1299967/3 > 420000,$$

using $\log 2 > 2/3$ and $\log(\pi/9) > -2$. Thus the same proof applies with these sharper budgets. Its lower is

$$\frac{1/1280}{(1 + 18/420000)^2} = \frac{3828125}{4900420009} > \frac{1}{1281}.$$

This uses the complete previously certified starting example and exact scalar comparisons. $\square$

S23 The earlier complete supported certificate

The finite criterion can be evaluated on a complete receiving space. The following worked example uses an alternative fixed support and receiving reference in the same original-form normalization. Put

$$M = 2^{20}, \quad \kappa = 2^{-20}, \quad L = \frac{1}{2} \log(4 \log(2^{20})/\pi) + \frac{1}{10}, \quad J = 74, \quad N = 329.$$

Use the exact original supported realization and normalization defined above. Let W_{low} be the dyadic theta floor specified by the accompanying certificate data, with $W/2 \leq W_{\text{low}} \leq W$, and put

$$S_\star = D_L + R_{L,74} + m_0 - C_L^{\text{ar}} + \kappa W_{\text{low}}.$$

Every prime power $n < e^{2L}$ and both killed-shift orientations occur in C_L^{ar} .

Theorem S23.1 (Evaluated complete supported certificate). *On the original minimal closed odd supported profile domain, subject to $\langle p, f \rangle = 0$, the form $S_\star$ is nonnegative. Consequently every original odd polar-free test supported in $(-L, L)$ satisfies*

$$q_H[h] \geq -2^{-20} \|h\|_2^2.$$

Proof. Apply the common complete receiver construction of main-paper Section D with $N = 329$, $J = 74$ and the support specified above. Thus C has 328 polar-free columns, $F = [p, C]$, $G_C = C^*C$ and $Q = \text{proj}_{\ker F^*}$. The reference-cell verification in the supplementary section *Verification of the complete supported example* certifies the complete arithmetic row upper r_{up} , the leakage-paid diagonal lower α and

$$(S_\star)_Q \succeq d_Q, \quad d = \kappa W_{\text{low}} + \alpha - r_{\text{up}} > 1/200.$$

The exact invertible dyadic conditioner R_G retains all 329 constraints and satisfies

$$\|R_G^*(F^*d^{-1}F)R_G - I\| < \delta, \quad \delta = 4 \times 10^{-75}.$$

Form B_E, K_E, H from the complete outputs as in main-paper (44). The directed lower matrix is

$$A_\star = \frac{H^* R_G R_G^* H}{1 + \delta} - B_E - K_E.$$

All 328 directed pivots of $P_C^* A_\star P_C$ are positive, where $P_C = G_C^{-1} R_{uu}^*$ and R_{uu} is the 328×328 core block of R_G . The supplementary verification gives the exact panels, elementary primitives, interval data and replay. The common full receiving comparison therefore proves $S_\star \succeq 0$, and its physical congruence proves the original-norm lower bound. $\square$

The zero receiving trial suffices. A separately verified 64-trial extension is supplementary reproducibility evidence for the same instance. The evaluated instance supplies a concrete realization of the complete finite-certificate mechanism at the stated support and cutoff.

S24 Complete fixed-support ridge continuation

The supported examples can also be studied as parameter families. The following comparison retains the full constrained reference inverse and all mixed columns. Support, constraints, Green model and legal trials remain fixed throughout each comparison.

Write $S_\kappa = S_0 + \kappa W_0$, $d_\kappa = d_0 + \kappa W_0$, where W_0 is a bounded positive multiplier on the fixed support. Let Q be the complete receiving projection and assume $QS_\kappa Q \succeq Qd_\kappa Q$, with d_κ uniformly positive on the parameter interval considered. Extend $Z_\kappa = (Qd_\kappa Q)^{-1}$ by zero off $\text{ran } Q$. For a fixed legal finite input map V , set

$$\Gamma_\kappa = V^* S_\kappa V - (S_\kappa V)^* Z_\kappa (S_\kappa V), \quad X_\kappa = V - Z_\kappa S_\kappa V.$$

These are sufficient reference-inverse lower matrices. The complete Schur comparison retains the corresponding actual receiving operator.

Proposition S24.1 (Complete derivative and downward loss). *On the common domain,*

$$\Gamma'_\kappa = X_\kappa^* W_0 X_\kappa \succeq 0.$$

At an admitted anchor κ_0 , put $D_0 = X_{\kappa_0}^* W_0 X_{\kappa_0}$ and $a = \text{ess sup } W_0/d_{\kappa_0}$. For $0 \leq t \leq h$ and $ha < 1$,

$$\Gamma_{\kappa_0-t} \succeq \Gamma_{\kappa_0} - \frac{t}{1-ta} D_0, \quad \Gamma'_{\kappa_0-t} \preceq (1-ta)^{-2} D_0.$$

Assume $\kappa_0 > 0$ and $0 < h < \kappa_0$. If a positive accepted anchor lower $L_0 \preceq \Gamma_{\kappa_0}$ satisfies $\kappa_0 D_0 \preceq cL_0$, and $b = 1 - ha > 0$, $g = 1 - ch/(\kappa_0 b) > 0$, then throughout that band

$$\kappa \Gamma'_\kappa \preceq \frac{c}{b^2 g} \Gamma_\kappa.$$

Proof. The constrained resolvent identity gives $Z'_\kappa = -Z_\kappa W_0 Z_\kappa$. Differentiating the entire quadratic expression, including both mixed terms, gives the derivative square. Put $A = W_0^{1/2} Z_{\kappa_0} W_0^{1/2}$. The multiplier-minus-finite-rank formula implies $0 \preceq A \preceq (W_0/d_{\kappa_0})I$, so $\|A\| \leq a$. Subtracting the constrained equations for X yields

$$W_0^{1/2} X_{\kappa_0-t} = (I - tA)^{-1} W_0^{1/2} X_{\kappa_0}.$$

Spectral calculus proves the derivative bound; integration gives the loss $t/(1-ta)$. The reference remains positive because $d_{\kappa_0-t} \geq (1-ta)d_{\kappa_0}$. Finally $D_0 \preceq c\Gamma_{\kappa_0}/\kappa_0$ and the loss bound give $\Gamma_\kappa \succeq g\Gamma_{\kappa_0}$. Combining these matrix orders proves the relative estimate without simultaneous diagonalization. $\square$

Application to the complete M24 example. With exactly the supported model and input contracts of the M24 example, the complete derivative-weight integrals and their independently reconstructed 659×659 derivative Gram certify a continuous ridge band. Its upper endpoint is $\kappa_0 = 2^{-24}$, and its exact lower endpoint is $\kappa_0 - h$, where

$$h = \frac{1125889145388605}{75557863725914323419136}, \quad a = \frac{61260186864964717714045042811731968}{3651321882289427623988064773}.$$

The lower endpoint is approximately $4.47036260082 \times 10^{-8}$. Complete directed matrix tests establish $\kappa_0 D_0 \preceq (1 + 2^{-20})L_0$ and the downward-band lower. Substitution in Proposition S24.1 gives

$$\kappa \Gamma'_\kappa \preceq 3\Gamma_\kappa \quad (\kappa_0 - h \leq \kappa \leq \kappa_0).$$

All 434281 derivative-matrix entries and all 659 directions are retained. This application uses new derivative-weight primitives; the original M24 moment archive alone does not reproduce that additional claim. Its scientific inputs, exact congruences and numerical contracts accompany this extension in the separate ridge-continuation dataset and portable verifier. The interval keeps the same support; a changing support changes the operator and requires a separate comparison.

Proposition S24.2 (Uniform capped return comparison). *Fix an admitted capped model with $S_w = S_{\text{base}} + w$, $d_w = d_{\text{base}} + w$, $d_{\text{base}} \geq \beta I$, and $Q(S_{\text{base}} - d_{\text{base}})Q \succeq 0$. For any $0 \leq w_1 \leq w_0 \leq \beta I$, keep the same constraints and legal input V . Put $\Delta = w_0 - w_1$, $A = \Delta^{1/2} Z_{w_0} \Delta^{1/2}$ and $B = \Delta^{1/2} X_{w_0}$. Then*

$$0 \preceq A \preceq \frac{1}{2}I, \quad \Gamma(w_1) = \Gamma(w_0) - B^*(I - A)^{-1}B.$$

For $M_j = B^* A^j B$ and every integer $m \geq 0$,

$$\sum_{j < m} M_j \preceq B^*(I - A)^{-1}B \preceq \sum_{j < m} M_j + 2^{1-m} M_0.$$

This holds also for $w_1 = 0$, including changes across clipping knots.

Proof. Inverse order gives $Z_{w_0} \preceq d_{w_0}^{-1}$ and $\Delta/(d_{\text{base}} + w_0) \leq 1/2$. Minimize the complete quadratic functional $V^* S_w V - 2\Re((S_w V)^* y) + y^* d_w y$ over $y \in \text{ran } Q$. Writing $y = Z_{w_0} S_{w_0} V + z$, completion of the square at w_1 subtracts $B^* B + B^* \Delta^{1/2} Z_{w_1} \Delta^{1/2} B$ from $\Gamma(w_0)$. The constrained resolvent identity makes its middle factor $(I - A)^{-1}$. The positive geometric series and $A^m (I - A)^{-1} \preceq 2^{1-m} I$ prove the moment enclosure. No inverse of the possibly singular Δ is used. $\square$

The contraction is uniform under the stated admissions. Certification also includes the size of the complete observation Gram M_0 and a directed signed lower bound after subtraction of the return charge. Every capped-model test uses its own recomputed inputs, including the moment, full-response and inverse errors.

S25 An alternative growing-floor arithmetic matrix criterion

The fixed-shift test has a complementary growing-floor version whose scalar tail payment uses the complete spatial action Gram. Proposition N.1, with exponent $a = 0$, permits an unweighted negative floor growing subexponentially with support. The construction below composes that implication with the complete arithmetic source Gram. Its matrix contains no Green action or receiving inverse; the complete receiving contribution is paid by an explicit scalar floor.

For each integer $j \geq 4$, set

$$L = j, \quad \beta_j = e^{\sqrt{j}}, \quad P_j = 8je^j.$$

Choose a certified rational τ_j and an outward integer N_j with

$$4P_j^2/\beta_j \leq \tau_j \leq 4P_j^2/\beta_j + 1, \quad X_j = P_j + |m_0| + \tau_j + 1, \quad \frac{je^{X_j}}{\pi} \leq N_j \leq \frac{je^{X_j}}{\pi} + 2. \quad (\text{S84})$$

These choices are effective: approximate the real lower threshold for τ_j to a rational interval of width less than one and use its upper endpoint. An interval of width less than one for je^{X_j}/π supplies an integer in the stated outward range, without an exact-real equality decision. Fix these same chosen values in all subsequent matrices.

Let U_j be the ordinary isometry onto the first N_j DN modes at support j , and $Q_j = I - U_j U_j^*$. Use the complete supported self-adjoint arithmetic operator $C_j = C_j^{\text{ar}}$, with both killed translations and the reflected odd channel for every $n < e^{2j}$. Put

$$\begin{aligned} A_j &= D_j + m_0 - C_j + \beta_j I, \\ A_j^{\text{ar}} &= U_j^* C_j U_j, & G_j^{\text{full}} &= (C_j U_j)^* (C_j U_j) = U_j^* C_j^2 U_j, \\ G_j^{\text{rec}} &= G_j^{\text{full}} - (A_j^{\text{ar}})^2, & H_j &= \text{diag}_{q < N_j} a(k_q) + (m_0 + \beta_j)I - A_j^{\text{ar}}, \\ \Gamma_j &= H_j - \tau_j^{-1} G_j^{\text{rec}}. \end{aligned} \tag{S85}$$

The square in $(A_j^{\text{ar}})^2$ is a matrix product. In particular, G_j^{full} integrates the complete spatial arithmetic outputs, not a finite receiving-output projection.

Theorem S25.1 (Explicit growing-floor matrix criterion). *Each Γ_j is a finite, effectively enclosable matrix. The following are equivalent:*

1. RH ;
2. $\Gamma_j \succ 0$ for every sufficiently large integer j ;
3. for every sufficiently large integer j , Γ_j admits a strictly positive directed lower enclosure with every coefficient and matrix error paid.

Under RH , $\Gamma_j \succeq \beta_j I/2$ for every $j \geq 4$. A two-sided matrix-center radius $e_j \leq \beta_j/8$ then leaves an outward lower at least $\beta_j I/4$. The specified sufficient dimension satisfies

$$\log N_j = O(je^j + j^2 e^{2j - \sqrt{j}} + \log j).$$

For $j = \lceil L_0(M) \rceil$, where $L_0(M) = \frac{1}{2} \log(10 \log M / \pi) + 2$, this gives $N_j = M^{o(1)}$. Here M labels support growth, not a shrinking original-norm allowance.

Proof. The elementary bounds $\Lambda(n) \leq \log n \leq 2j$ for $n < e^{2j}$ and $\sum_{2 \leq n < K} n^{-1/2} \leq 2\sqrt{K}$ give

$$\sum_{2 \leq n < e^{2j}} \frac{\Lambda(n)}{\sqrt{n}} \leq 4je^j, \quad \|C_j\| \leq P_j.$$

The second estimate retains the entire paired arithmetic operator; odd restriction cannot increase its norm. On the whole DN suffix, $k_{N_j} > e^{X_j} > 1$ and $a(k) \geq \log k$ imply

$$T_j = Q_j A_j Q_j|_{\text{ran } Q_j} \succeq [a(k_{N_j}) + m_0 - P_j + \beta_j]I \succeq (\tau_j + 1 + \beta_j)I \succeq \tau_j I.$$

This is the full receiving floor on the same minimal supported domain. The exact mixed map is $C_j^{\text{mix}} = -Q_j C_j U_j$, and therefore

$$(C_j^{\text{mix}})^* C_j^{\text{mix}} = U_j^* C_j Q_j C_j U_j = G_j^{\text{rec}} \succeq 0.$$

Inverse order proves the complete Schur lower

$$H_j - (C_j^{\text{mix}})^* T_j^{-1} C_j^{\text{mix}} \succeq \Gamma_j.$$

Hence a positive directed lower for Γ_j proves $A_j \succeq 0$ on its entire supported domain by Schur square completion.

Every arithmetic input action is a finite collection of killed and reflected sine pieces. A common finite endpoint partition reduces both A_j^{ar} and G_j^{full} to continuous piecewise trigonometric integrals, with all paired cross terms retained. The digamma diagonal also has directed enclosures. This proves finite entry computability. Subtraction in G_j^{rec} must be enclosed in the full matrix order; its exact positivity alone does not price numerical cancellation.

Under RH, the odd pole calculation and Lemma B.1 give $A_j \succeq (\beta_j - \pi/2)I$. Since $\|C_j^{\text{mix}}\| \leq P_j$ and $\tau_j \geq 4P_j^2/\beta_j$,

$$\Gamma_j \succeq (3\beta_j/4 - \pi/2)I \succeq \beta_j I/2.$$

Here $j \geq 4$ gives $\beta_j \geq e^2 > 2\pi$. Subtracting a two-sided center radius loses at most $2e_j$, giving the stated directed margin and effective recognition at each fixed j .

Conversely, suppose the matrices pass for every integer $j \geq J$. The raw supported form $B_j = A_j + R_j - \beta_j I$ then satisfies $B_j \succeq -\beta_j I$. Embed any compact real odd polar-free profile supported in $[-L, L]$ at $j = \lceil L \rceil \geq J$. The complete forms agree on it under zero extension and the half-unitary map preserves its unweighted norm. Thus

$$q_{\text{geo}}[f] \geq -e^{\sqrt{\lceil L \rceil}} \|f\|_2^2.$$

For every $\varepsilon > 0$, $\sqrt{\lceil L \rceil} \leq \sqrt{L+1} \leq \varepsilon(L+1) + 1/(4\varepsilon)$. This supplies one support-growth constant $C_\varepsilon = e^{\varepsilon+1/(4\varepsilon)}$ for all admitted profiles and sufficiently large supports. Proposition N.1 with $a = 0$ proves RH. No arbitrary gapped support subsequence is used in this step.

Finally the upper guards in (S84) give $\log N_j = O(\log j + P_j + P_j^2/\beta_j)$, yielding the stated bound. For $j = \lceil L_0(M) \rceil$, $e^{2j} = O(\log M)$ and $j = O(\log \log M)$; hence

$$\frac{j^2 e^{2j-\sqrt{j}}}{\log M} = O(j^2 e^{-\sqrt{j}}) \longrightarrow 0, \quad \frac{j e^j + \log j}{\log M} \longrightarrow 0.$$

This proves the sufficient dimension assertion for the prescribed near-threshold choices. Entry count, precision and validation costs are separate; this dimension estimate is not a total runtime bound. $\square$

Corollary S25.2 (A prescribed controlled-sparse support sequence). *Fix a strictly increasing unbounded integer sequence j_k such that $j_{k+1} = o(j_k^2)$. RH is equivalent to existence of strictly positive, fully paid lower enclosures of the matrices (S85) at every sufficiently large j_k . In particular the prescribed sequence $j_k = 2^k$ suffices.*

Proof. Under RH the preceding strict margin applies at every support. Conversely take an arbitrary sufficiently large support L and its first containing sequence support. If $j_k < L \leq j_{k+1}$, the accepted form gives the common floor $e^{\sqrt{j_{k+1}}}$ for every legal profile at that L . Moreover

$$\frac{\sqrt{j_{k+1}}}{L} \leq \frac{\sqrt{j_{k+1}}}{j_k} \longrightarrow 0.$$

Equality $L = j_k$ uses that support directly. Thus for every $\varepsilon > 0$ the floor is at most $e^{\varepsilon L}$ for all sufficiently large real L , uniformly over the profiles. The support-growth theorem proves RH. For $j_k = 2^k$, the next containing support is at most $2L$, giving the explicit envelope $e^{\sqrt{2L}} \leq e^{1/(2\varepsilon)} e^{\varepsilon L}$. $\square$

Along a sparse sequence the resource label is its native $M_j = \exp\{(\pi/10)e^{2(j-2)}\}$, for which $L_0(M_j) = j$ and $N_j = M_j^{o(1)}$. Rounding an arbitrary $L_0(M)$ up to its next doubling support can change this comparison and does not establish a uniform $N = M^{o(1)}$ bound in that arbitrary smaller label. The criterion specifies an infinite signed matrix target; finitely many passing tests do not decide its all-needed-support premise.

References

- [1] J. Behrndt, S. Hassi, H. S. V. de Snoo and R. Wietsma, Monotone convergence theorems for semibounded operators and forms with applications, 2009, Theorem 4.2. <https://www.applied.math.tugraz.at/~behrndt/BehrndtHassiSnooWietsma09.pdf>.
- [2] A. Bonami and A. Karoui, Spectral decay of time and frequency limiting operator, 2015. <https://arxiv.org/abs/1509.02646>.
- [3] F. Bornemann, *On the numerical evaluation of Fredholm determinants*, Math. Comp. 79 (2010), 871–915. <https://arxiv.org/abs/0804.2543>.
- [4] bramble_trace, *Boundary traces and support variation for localized Weil operators*, rStation Riemann hypothesis paper 67, version 1, 27 October 2026. https://rstation.ai/problems/riemann_hypothesis/papers/67.
- [5] J. B. Conrey, More than two fifths of the zeros of the Riemann zeta function are on the critical line, *J. reine angew. Math.* **399** (1989), 1–26.
- [6] R. G. Douglas, On majorization, factorization, and range inclusion of operators on Hilbert space, *Proc. Amer. Math. Soc.* **17** (1966), 413–415.
- [7] G. Dusson, I. M. Sigal and B. Stamm, The Feshbach–Schur map and perturbation theory, 2021. <https://arxiv.org/abs/2105.02058>.
- [8] N. J. Higham, *Accuracy and Stability of Numerical Algorithms*, second ed., SIAM, 2002. <https://doi.org/10.1137/1.9780898718027>.
- [9] A. Israel and A. Mayeli, On the eigenvalue distribution of spatio-spectral limiting operators in higher dimensions, 2023. <https://arxiv.org/abs/2301.09616>.
- [10] D. R. Johnston and A. Yang, *Some explicit estimates for the error term in the prime number theorem*, arXiv:2204.01980v2, 2022. <https://arxiv.org/abs/2204.01980v2>.
- [11] NIST Digital Library of Mathematical Functions, Section 5.11, equation 5.11.2. <https://dlmf.nist.gov/5.11.E2>.
- [12] S. Schubr, *Certified complementary bounds for an odd Weil receiver*, Luminary Research Archive, author-posted preprint, 2026. <https://research.luminaryoddities.com/certified-complementary-bounds-odd-weil-receiver/v/1>.
- [13] S. Schubr, *Joint Cutoffs and Negative-Spectrum Decay for Weighted Weil Operators*, Luminary Research Archive, author-posted preprint, edition 2, 2026. <https://research.luminaryoddities.com/joint-cutoffs-negative-spectrum-decay-weighted-weil-operators/v/2>.
- [14] S. Schubr, *Quantitative Null-Space Compression for a Weighted Weil Operator*, Luminary Research Archive, author-posted preprint, 2026. <https://research.luminaryoddities.com/quantitative-null-space-compression-weighted-weil-operator/v/1>.
- [15] S. Schubr, *RH, Approximately: Certified Lower Bounds for Odd Weil Positivity*, Luminary Research Archive, author-posted preprint, 2026. <https://research.luminaryoddities.com/rh-approximately-certified-lower-bounds-odd-weil-positivity/v/1>.

- [16] S. Schubr, *Signed Frequency Certificates and Constraint Costs for Odd Weil Forms*, Luminary Research Archive, author-posted preprint, 2026. <https://research.luminaryoddities.com/signed-frequency-certificates-constraint-costs-odd-weil-forms/v/1>.
- [17] A. N. Tikhonov and V. Y. Arsenin, *Solutions of Ill-Posed Problems*, Winston, Washington, DC, 1977.
- [18] F. Zhang (ed.), *The Schur Complement and Its Applications*, Springer, 2005. <https://doi.org/10.1007/b105056>.