

Statistical inference for empirical heat-kernel Sinkhorn metrics

Scott Schubr
Luminary Oddities LLC

October 22, 2026

Abstract

We estimate fixed smooth pullback entries of the inverse weighted-Laplacian metric of an unknown smooth strictly positive density on a flat torus. The estimator uses symmetric heat-kernel Sinkhorn normalization of the sample and prescribed score functions, without density evaluations. For every fixed dimension we establish a joint, population-centered root- n central limit theorem as the heat bandwidth tends to zero. The proof gives a signed-measure Taylor expansion on one Sobolev space, an explicit influence function including normalization and inverse response, and a consistent empirical covariance estimator using exact solves. A scalar Hessian identity with the moving integration measure improves the sufficient remainder budget. The population tensor has even finite asymptotic jets; fixed multiband extrapolation removes enough bias to retain the same limiting covariance. The influence is the canonical gradient, and the estimator attains the pointwise information bound under smooth local density submodels. Finally, an observed scaling, spectral-gap and inverse-residual certificate produces an actual finite first-hit output with negligible root- n metric error. Its certified-arithmetic realization requires effective convergent input enclosures. Bandwidth conditions are conservative; no practical runtime, optimal smoothing rate or numerically certified covariance algorithm is asserted.

Keywords: Sinkhorn metric; diffusion maps; inverse weighted Laplacian; asymptotic linearity; semi-parametric efficiency; observed stopping.

1 Introduction

The local metric of a Sinkhorn divergence contains an inverse whose nonconstant spectral gap shrinks with regularization. Consequently a fixed-bandwidth delta method, or consistency of a normalized graph operator, does not by itself give a central limit theorem for the limiting inverse metric. A further distinction is important statistically: replacing the outer population integral by a sample average while freezing the normalized operator omits part of the influence of the sample.

We study a specific periodic heat-kernel model. For an unknown smooth positive density f and fixed smooth functions u_a , the target is the finite matrix of inverse weighted-Laplacian energies of the centered scores $u_a - \int u_a f$. The sample supplies both the normalization and the integration measure. We prove a bandwidth-uniform scalar expansion, remove population bias by fixed-band extrapolation, and identify the complete joint limiting distribution. A deterministic certificate makes the numerical metric target explicit, including the change of mass and centering caused by an imperfect scaling iterate.

Relation to earlier work. Sinkhorn divergence geometry and its Hessian metric are established objects [8]. Symmetric normalization of diffusion kernels, spectral convergence and observed stopping were studied by Wormell–Reich [1]; approximate bistochastic normalization and graph-Laplacian convergence are treated by Cheng–Landa [2]. Kessler–Magnet–Feydy [3], in particular their mass-variation formulas, provide direct antecedents for differentiating the normalization. These are predecessors of the present construction, rather than consequences of it.

Fixed-regularization differentiability and distributional limits for entropic transport are developed by Goldfeld–Kato–Rioux–Sadhu [4]. Root- n graph eigenvalue fluctuations, variance estimation and information-geometric interpretations are studied by Li–García Trillos–Li–Suchan [5]. Efficient inference for smooth transport functionals already appears in Manole et al. [6] and, using elliptic functional derivatives, in Kono [7]. Our observable is instead a fixed-score inverse metric with empirical Sinkhorn normalization and a simultaneous vanishing heat bandwidth. Horizontal metric limits on a fixed finite support [9] concern a different perturbation and sampling regime.

The population inverse-metric limit and smooth-input bias were studied in [10, 11]. In particular the reversibility mechanism behind the even tensor expansion belongs to that population analysis. Here we give its finite-jet form at every required order and use it to simplify the empirical bias correction. The new statistical work is the shrinking-bandwidth scalar remainder, complete same-sample influence and covariance, and their combination with certified numerical equivalence. Sobolev perturbation, extrapolation, iid central limits, LAN information bounds and residual-Gram identities are classical tools. We make no claim to introduce those methods.

2 Model and main results

Let $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$ have volume one. Throughout d is fixed, $\mu = f \, dx$ is a probability measure with $f \in C^\infty(\mathbb{T}^d)$ strictly positive, and $u_1, \dots, u_p$ are fixed real smooth functions. Neither p nor these functions grow with the sample size. Write $\nu(\nu) = \int \nu \, d\nu$ and use the periodic heat kernel

$$H_h = e^{h\Delta/2}, \quad H_h \nu(x) = \int k_h(x-y) \nu(y) \, dy, \quad h > 0.$$

For a probability measure ν let $a = a_{h,\nu} > 0$ solve

$$aH_h(av) = 1, \quad P_{h,\nu} \nu = aH_h(av\nu). \quad (1)$$

For smooth μ the small- h solution is constructed below. For a finite positive measure on distinct points it is the usual unique symmetric positive scaling; its extension to all x is specified by (1). The operator preserves constants and ν , is selfadjoint and nonnegative in $L^2(\nu)$, and has a simple constant eigenvalue. Define $\Pi_\nu \nu = \nu(\nu)1$ and

$$T_{h,\nu} = \frac{h}{2} \left\{ (I - P_{h,\nu} + \Pi_\nu)^{-1} - (I + P_{h,\nu})^{-1} - \frac{\Pi_\nu}{2} \right\}. \quad (2)$$

Thus $T1 = 0$, while on the mean-zero space $T = hP(I - P^2)^{-1}$. In particular the last projection in (2) cannot be omitted when the measure varies. The empirical pullback metric is

$$G_{ab}(h, \nu) = \nu[u_a T_{h,\nu} u_b]. \quad (3)$$

For the entropic interpretation use $\varepsilon = 2h$ and the periodic cost $c_h(x, y) = -2h \log[k_h(x-y)/k_h(0)]$. If $\text{OT}_{\varepsilon,c}(\nu, \rho)$ denotes the infimum of $\int c \, d\pi + \varepsilon \text{KL}(\pi \mid \nu \otimes \rho)$ over couplings, its debiased divergence subtracts the two half self-costs. The established diagonal quadratic-coefficient formula is $(\varepsilon/2)P(I - P^2)^{-1}$ on mean-zero scores [8, 10]; hence the convention (2). The analysis below can equivalently take (3) as its definition. Equivalently one may center each input at ν . Its prescribed statistical family is the exponential tilt

$$\nu_\theta = \frac{\exp(\sum_a \theta_a u_a) \nu}{\nu[\exp(\sum_a \theta_a u_a)]}.$$

This definition requires no evaluation of an unknown density ratio. It does not estimate the geometry of an independently prescribed unknown parametric density family.

Put $u_a^0 = u_a - \mu(u_a)$ and

$$D_f = -f^{-1} \text{div}(f \nabla), \quad D_f \phi_a = u_a^0, \quad \mu \phi_a = 0, \quad G_{ab}^0 = \mu(u_a^0 \phi_b) = \mu(\nabla \phi_a \cdot \nabla \phi_b). \quad (4)$$

Smooth elliptic theory gives the indicated solution. Define

$$\psi_{ab} = u_a^0 \phi_b + u_b^0 \phi_a - \nabla \phi_a \cdot \nabla \phi_b - G_{ab}^0, \quad \Sigma_{ab,cd} = \mu(\psi_{ab} \psi_{cd}). \quad (5)$$

The vectorization below uses entries $a \leq b$.

Theorem 2.1 (Population-centered empirical inference). *Let $X_1, \dots, X_n$ be iid with law μ and $\mu_n = n^{-1} \sum_i \delta_{X_i}$. Choose an integer $s > d/2$ and set*

$$q = 1 + s/2, \quad M = 2 + 5s/2. \quad (6)$$

Take an integer K with $2K > M$, a constant $1/(4K) < \beta < 1/(2M)$, and $h_n = n^{-\beta}$. Let $c_0, \dots, c_{K-1}$ be the unique real coefficients satisfying

$$\sum_{\ell=0}^{K-1} c_\ell = 1, \quad \sum_{\ell=0}^{K-1} c_\ell 4^{-\ell j} = 0 \quad (1 \leq j < K). \quad (7)$$

Then

$$\sqrt{n} \operatorname{vec} \left\{ \widehat{G}_n - G^0 \right\} \implies N(0, \Sigma), \quad \widehat{G}_n = \sum_{\ell=0}^{K-1} c_\ell G(h_n/2^\ell, \mu_n). \quad (8)$$

More precisely, the left side is $n^{-1/2} \sum_i \operatorname{vec} \psi(X_i) + o_{\mathbb{P}}(1)$. The density is not an input to $\widehat{G}_n$. The covariance may be singular. There is a consistent sample-computable covariance estimator using exact empirical influence solves, given in Section 6.

Corollary 2.2 (Certified numerical equivalence). *Set $C_K \geq \sum_\ell |c_\ell|$. If the computed metric at each band has a certified operator-norm error at most $n^{-3/4}/C_K$, replacing the exact band metrics in (8) preserves its conclusion. Section 8 constructs a finite first-hit algorithm with this property on every finite distinct sample set. A certified finite-arithmetic realization requires effective convergent enclosures of sites, kernel and score values, and eventually informative spectral-gap enclosures. It asserts neither fast runtime nor fixed-precision success. The metric certificate alone does not certify numerical covariance or studentization.*

Theorem 2.3 (Information bound). *In the model of all smooth strictly positive densities, the vector $\operatorname{vec} \psi$ is the canonical gradient of $\operatorname{vec} G^0$. The estimators in Theorem 2.1 and Corollary 2.2 are regular along each fixed finite smooth exponential density submodel under local $n^{-1/2}$ alternatives. They attain the pointwise semiparametric covariance bound Σ for this notion of regularity. Singular directions are allowed. This is not uniform inference over a density class, a finite-sample variance inequality, or an optimal bandwidth statement.*

The metric is a periodic heat-kernel metric, whose cost changes with h ; fixed Euclidean squared-cost problems and boundary domains are not included. Signed extrapolation can produce an indefinite matrix at finite n . If G^0 is positive definite, consistency implies positive definiteness with probability tending to one.

3 A fixed-space measure perturbation

We give the analytic estimates needed for the simultaneous limit. Constants may depend on the fixed density, scores, dimension and finitely many derivatives, but not on small h or n . Use the Sobolev norm with Fourier weight $(1 + 4\pi^2 |j|^2)^s$. Since $s > d/2$, H^s is an algebra, embeds in C^0 , and its multipliers act on H^{-s} . In particular

$$\|H_h\|_{H^{-s} \rightarrow H^s} \leq C h^{-s}. \quad (9)$$

All measure perturbations have total mass zero. Signed measures and distributions are used locally to define derivatives; probability measures are used for the statistical values.

Lemma 3.1 (Population scale and gap). *At every fixed sufficiently large integer Sobolev order m ,*

$$a_{h,\mu} = f^{-1/2} + O_{H^m}(h).$$

The plus inverse $(I + P_{h,\mu})^{-1}$ is uniformly bounded on H^m . The nonconstant $L^2(\mu)$ gap of $P_{h,\mu}$ is at least ch ; on H^m ,

$$\|(I - P_{h,\mu} + \Pi_\mu)^{-1}\| \leq Ch^{-1-m/2}. \quad (10)$$

Proof. Write $a = f^{-1/2}r$. At $r = 1$ the normalization residual is $f^{-1/2}H_h\sqrt{f} - 1 = O_{H^m}(h)$. The derivative in r is the sum of multiplication by $1 + O_{H^m}(h)$ and $P_0 = f^{-1/2}H_h(\sqrt{f} \cdot)$. Conjugating by multiplication by $\sqrt{f}$ turns $I + P_0$ into $I + H_h$, whose inverse has norm at most one on H^m . Algebra estimates bound the second derivative uniformly on a fixed neighborhood. A contraction/Newton argument therefore gives $r = 1 + O_{H^m}(h)$ and a uniformly bounded local normalization inverse. This construction works at any prescribed finite high order; uniqueness identifies the solutions. Positivity follows from the C^0 bound.

For the exact scale, conjugation by $\sqrt{f}$ gives $Q = bH_hb$, $b = a\sqrt{f} = 1 + O_{H^m}(h)$. Thus $I + Q$ is an $O(h)$ bounded perturbation of $I + H_h$ on H^m , proving the plus assertion. In $L^2(\mu)$, kernel symmetry gives

$$\mu[v(I - P)v] = \frac{1}{2} \iint k_h(x - y)a(x)a(y)f(x)f(y)(v(x) - v(y))^2 \, dx \, dy.$$

The coefficients are uniformly bounded above and below. Fourier Poincaré for $I - H_h$ gives a lower bound ch times the Lebesgue variance. The latter controls the μ variance, since f is bounded above and below and variance is the minimum over constant centers. This proves the gap. Positivity of P follows from that of H_h . Its constant eigenspace is simple by strict positivity of the kernel. Finally, writing $R_- = (I - P + \Pi_\mu)^{-1}$,

$$R_- = I + PR_- - \Pi_\mu, \quad \|R_-\|_{L^2(\mu) \rightarrow L^2(\mu)} \leq C/h, \quad \|P\|_{L^2(\mu) \rightarrow H^m} \leq Ch^{-m/2}$$

gives (10). $\square$

Lemma 3.2 (Local normalization and inverses). *Let $\epsilon = \|\zeta\|_{H^{-s}}$ and $v = \mu + \zeta$. There is a constant $c > 0$ such that on*

$$\epsilon \leq ch^{s+q} \quad (11)$$

the normalization and (2) are smooth maps on fixed Sobolev spaces. Uniformly there,

$$\|a_{h,v}/a_{h,\mu} - 1\|_{H^s} \leq C\epsilon h^{-s}, \quad (12)$$

$$\|DP_{h,v}\| \leq Ch^{-s}, \quad \|D^2P_{h,v}\| \leq Ch^{-2s}, \quad (13)$$

$$\|R_{-,v}\| \leq Ch^{-q}, \quad \|R_{+,v}\| \leq C. \quad (14)$$

Derivative norms take H^{-s} directions and output bounded H^s operators; $R_{-,v} = (I - P_{h,v} + \Pi_v)^{-1}$ and $R_{+,v} = (I + P_{h,v})^{-1}$.

Proof. Put $a = a_{h,\mu}$ and $a_{h,v} = ar$. The equation is

$$r\{Pr + aH_h(ar\zeta)\} = 1.$$

Its derivative at $(r, \zeta) = (1, 0)$ is $I + P$, with the uniform inverse from Lemma 3.1. The residual is $O(\epsilon h^{-s})$. On a fixed H^s neighborhood the second r derivative is $O(1 + \epsilon h^{-s})$, and the perturbation of its first derivative is $O(\|r - 1\|_{H^s} + \epsilon h^{-s})$. A quantitative implicit argument gives (12) and

$$\|Dr\| \leq Ch^{-s}, \quad \|D^2r\| \leq Ch^{-2s}$$

already on $\epsilon \leq ch^s$. Product differentiation then gives (13) and $\|P_{h,v} - P_{h,\mu}\| \leq C\epsilon h^{-s}$. The rank-one projection perturbation has norm at most $C\epsilon$. By (10) the Neumann criterion for the minus inverse is precisely (11) with sufficiently small c ; the plus inverse has a weaker criterion. This proves the claims.

All identities $P1 = 1$, $\nu P = \nu$ and kernel symmetry persist on the signed local extension. On probability measures the local scale is positive by the C^0 estimate. For a finite positive measure it agrees with the unique finite scale: that scale is the critical point of the strictly convex logarithmic scaling energy

$$\frac{1}{2} \sum_{i,j} w_i w_j K_{ij} e^{x_i + x_j} - \sum_i w_i x_i.$$

Strict positivity of the entries, including the diagonal, makes this energy strictly convex and coercive. Its critical equation is the scaling equation. The extension away from the atoms is $a(x) = 1/H_h(a\nu)(x)$ and hence also agrees. Restriction to the atoms intertwines P, Π_ν , both shifted operators and their inverses. Thus the fixed-space functional is the exact finite sample metric, not a quadrature proxy. $\square$

4 Finite smooth expansions and even bias

The next argument also supplies the strong smooth-input estimates used in the scalar Hessian and influence calculations. We include the remainder justification because a formal series alone would not suffice for population centering.

Lemma 4.1 (Finite smooth expansions). *For each prescribed integer order N and each fixed Sobolev order, the population scale, $h(I - P_h)^{-1}u^0$ and $(I + P_h)^{-1}u^0$ have finite asymptotic expansions through order N , with controlled next-order remainders, for every fixed smooth u . Their leading terms are respectively $f^{-1/2}$, $2D_f^{-1}u^0$ and $u^0/2$.*

Proof. On smooth functions the heat Taylor formula is

$$H_h v = \sum_{j=0}^N \frac{h^j}{2^j j!} \Delta^j v + O_{H^m}(h^{N+1});$$

its remainder is bounded by a constant times a sufficiently high finite Sobolev norm of v , by the integral semigroup remainder. Insert $a^{[N]} = \sum_{j=0}^N h^j a_j$ in $aH_h(af) = 1$. At order zero $a_0 = f^{-1/2}$; at each order j the new coefficient is multiplied by $2\sqrt{f}$. This determines smooth a_j recursively. The residual is $O_{H^m}(h^{N+1})$. The uniform local normalization inverse upgrades this to the same order of error for the exact scale. Consequently on fixed smooth inputs

$$P_h = I - \frac{h}{2} D_f + \sum_{j=2}^N h^j L_j + O(h^{N+1}). \quad (15)$$

Each L_j is a finite-order smooth differential operator. Since the exact P_h fixes constants and preserves μ , uniqueness of finite asymptotic coefficients gives $L_j 1 = 0$ and $\mu L_j = 0$.

Let $y_h = h(I - P_h)^{-1}u^0$ with $\mu y_h = 0$. Construct $y^{[N]} = \sum_{j=0}^N h^j y_j$ with $y_0 = 2D_f^{-1}u^0$ and all coefficients mean zero. Matching successive orders of $(I - P_h)y = hu^0$ requires

$$\frac{D_f}{2} y_j = \sum_{\ell=2}^{j+1} L_\ell y_{j+1-\ell}.$$

Every right side has mean zero; smooth elliptic solvability applies. The residual of a sufficiently long truncation is $O_{H^m}(h^{N+2})$. Applying the coarse inverse of Lemma 3.1 bounds its error by $Ch^{N+2-(1+m/2)}$. For any desired expansion order, start with a longer truncation than that order plus this fixed loss. The low coefficients are unchanged, giving an expansion to every fixed order. The plus equation is treated by its uniformly bounded inverse and a residual expansion beginning with $u^0/2$. These arguments use finite truncations only. $\square$

Proposition 4.2 (Even population jets). *For each fixed integer $K \geq 1$,*

$$G_{ab}(h, \mu) = G_{ab}^0 + \sum_{j=1}^{K-1} b_{ab,2j} h^{2j} + O(h^{2K}). \quad (16)$$

This is the all-finite-orders form of the population reversal mechanism used in [11]; it is used here for empirical bias removal. It does not define backward heat evolution.

Proof. By Lemma 4.1, genuine positive- h expansions exist to every fixed order. We identify their unique coefficients using formal series with smooth coefficients. In this paragraph negative h denotes substitution in finite jets, never a bounded heat operator at negative time. The formal differential-operator series $H(h)$ satisfies $H(h)H(-h) = I$ coefficientwise by the binomial identity. The normalization coefficients are unique. From $H(h)(a(h)f) = 1/a(h)$ one obtains formally $H(-h)(1/a(h)) = a(h)f$. Hence $\tilde{a} = 1/(fa(h))$ solves the normalization at $-h$ and has the same leading coefficient; uniqueness gives

$$a(-h) = \frac{1}{fa(h)}.$$

If $A(h)$ is multiplication by $a(h)$ and M_f by f , then

$$P(h) = A(h)H(h)A(h)M_f, \quad P(h)^{-1} = M_f^{-1}A(h)^{-1}H(-h)A(h)^{-1} = P(-h).$$

The order of multiplication is essential here. The fixed μ -mean-zero smooth space is invariant coefficientwise. On it $I - P(h)^2$ has leading term hD_f , so elliptic invertibility gives a unique formal Laurent inverse, also selected by the finite-residual construction. Since all these factors are functions of the same $P(h)$,

$$(-h)P(h)^{-1}(I - P(h)^{-2})^{-1} = hP(h)(I - P(h)^2)^{-1}.$$

The factor h removes the pole. Thus the tensor jet is even; extend it by zero on constants. Every odd coefficient of each fixed smooth bilinear form vanishes. The controlled expansions from Lemma 4.1 now prove (16) for actual positive h . There is no analyticity or convergent infinite-series assertion. $\square$

5 Scalar Hessian and exact influence

For a fixed smooth real u set $\Phi_h(v) = v[uT_{h,v}u]$.

Proposition 5.1 (Uniform scalar Taylor bound). *On the neighborhood (11),*

$$\|D^2\Phi_h(v)\| \leq Ch^{-M}, \quad M = 2 + 5s/2. \quad (17)$$

In particular

$$|\Phi_h(\mu + \zeta) - \Phi_h(\mu) - D\Phi_h(\mu)[\zeta]| \leq Ch^{-M}\|\zeta\|_{H^{-s}}^2. \quad (18)$$

Proof. The improved scalar estimate uses the fixed smooth input before applying a coarse inverse bound. Let B_v be either $I - P_v + \Pi_v$ or $I + P_v$ and $R_v = B_v^{-1}$. Lemma 4.1 gives $\|R_\mu u\|_{H^s} \leq C/h$, including its bounded constant part. The identity

$$R_v u = (I + R_\mu(B_v - B_\mu))^{-1}R_\mu u$$

and the same Neumann neighborhood give $\|z\|_{H^s} \leq C/h$ for $z = R_v u$, uniformly in v . Only an inverse acting on a general perturbed input will need the larger loss h^{-q} .

We account explicitly for the moving pairing. Along $v_t = v + t\zeta$ put $A = \dot{B}_v$ and $F(v) = v[uR_v u]$. Then

$$\dot{z} = -R_v A z, \quad \dot{F} = \zeta(uz) - v(zAz).$$

Although B_ν is symmetric in the ν pairing, its derivative A need not be. Differentiating the symmetry identity for fixed ν, w yields

$$\nu(\nu Aw) - \nu(wAv) = \zeta(wB_\nu\nu - \nu B_\nu w).$$

With $\nu = z, w = \dot{z}, B_\nu z = u$ and $B_\nu \dot{z} = -Az$, this becomes

$$\nu(zA\dot{z}) - \nu(\dot{z}Az) = \zeta(u\dot{z} + zAz).$$

Differentiating $\dot{F}$ and cancelling the indicated terms gives the exact identity

$$\ddot{F} = 2\nu[(Az)R_\nu(Az)] - \nu[z\ddot{B}_\nu z] - 2\zeta(zAz). \quad (19)$$

It is valid on the signed extension: kernel symmetry and the rank-one projection give the algebraic symmetry there as well. All pairings are continuous by the H^s algebra and duality. For a unit H^{-s} direction, Lemma 3.2 bounds the three terms by

$$Ch^{-2-2s-q}, \quad Ch^{-2-2s}, \quad Ch^{-2-s},$$

respectively. Here ν has bounded H^{-s} norm on the ball. Finally the exact uncentered representation is

$$\Phi_h(\nu) = \frac{h}{2} \left\{ F_-(\nu) - F_+(\nu) - \frac{\nu(u)^2}{2} \right\}.$$

The last term has a bounded second derivative times h . Multiplying the preceding bounds by h gives $M = 1 + q + 2s = 2 + 5s/2$. Polarization bounds mixed directions; integrating the second derivative along the segment proves (18). A full operator Hessian estimate would have spent three coarse inverse factors and would give a weaker scalar budget; (19) avoids that loss. $\square$

Proposition 5.2 (Influence at smooth and atomic laws). *At a probability measure ν in the local neighborhood, center $g = u - \nu u$ and set*

$$z_- = (I - P_\nu)^{-1}g \quad (\nu z_- = 0), \quad z_+ = (I + P_\nu)^{-1}g, \quad C_\nu = (I - P_\nu)(I + P_\nu)^{-1}.$$

The centered derivative representative is

$$\Psi_{h,\nu} = \frac{h}{2} \{g(z_- - z_+) + C_\nu(z_- P_\nu z_- + z_+ P_\nu z_+)\}, \quad (20)$$

$$\psi_{h,\nu} = \Psi_{h,\nu} - \Phi_h(\nu), \quad D\Phi_h(\nu)[\zeta] = \zeta(\psi_{h,\nu}), \quad \nu\psi_{h,\nu} = 0. \quad (21)$$

These formulas hold for every mass-zero H^{-s} direction, including directions off the support of an atomic ν . At $\nu = \mu$,

$$\psi_{h,\mu} \longrightarrow 2u^0\phi - |\nabla\phi|^2 - G^0(u) \quad (22)$$

in every prescribed finite Sobolev order as $h \downarrow 0$.

Proof. Define $S_\nu\zeta = a_\nu H_h(a_\nu\zeta)$. Differentiating normalization gives $\alpha = \dot{a}/a = -(I + P_\nu)^{-1}S_\nu\zeta$ and

$$\dot{P}_\nu z = \alpha P_\nu z + P_\nu(\alpha z) + S_\nu(z\zeta).$$

Kernel symmetry directly gives

$$\begin{aligned} \nu[zS_\nu(z\zeta)] &= \zeta[zP_\nu z], \\ \nu[B(I + P_\nu)^{-1}S_\nu\zeta] &= \zeta[P_\nu(I + P_\nu)^{-1}B]. \end{aligned}$$

Therefore

$$\nu[z\dot{P}_\nu z] = \zeta[C_\nu(zP_\nu z)]. \quad (23)$$

These are dual-kernel identities, not a density argument using perturbations $\nu\nu$. Such perturbations are not dense at a fixed finite atomic law. All their kernels and products are smooth and their pairings extend continuously to H^{-s} .

Differentiate the quadratic resolvent forms. Derivatives of the input center multiply a mean-zero resolvent output and vanish; the derivative of the rank-one projection in the minus inverse has zero quadratic pairing for the same reason. The minus form has derivative $\zeta(gz_-) + \nu(z_- \dot{P}z_-)$; the plus form has derivative $\zeta(gz_+) - \nu(z_+ \dot{P}z_+)$. Equation (23) proves (20). Since $\nu C_\nu = 0$, $\nu\Psi_{h,\nu} = \Phi_h(\nu)$, which proves the centering and derivative assertions.

At the population law Lemma 4.1 gives $y_h = hz_- \rightarrow 2\phi$ and $z_+ \rightarrow u^0/2$ in arbitrarily high prescribed Sobolev orders. Equation (15) and the uniform plus inverse give, on smooth inputs,

$$C_h/h \rightarrow D_f/4, \quad \|C_h/h\|_{H^{m+2} \rightarrow H^m} \leq C.$$

The bound follows from $(I - P_h)/h$ and the plus inverse on H^{m+2} ; the limit follows first on fixed smooth functions and then on convergent high-Sobolev inputs. Rewriting the singular contribution to (20) as $\frac{1}{2}(C_h/h)(y_h P_h y_h)$ shows

$$\Psi_{h,\mu} \rightarrow u^0\phi + \frac{1}{2}D_f(\phi^2) = 2u^0\phi - |\nabla\phi|^2.$$

The identity $D_f(\phi^2) = 2\phi D_f\phi - 2|\nabla\phi|^2$ fixes both the sign and coefficient. Its mean is $G^0(u)$ by integration by parts. This proves (22). $\square$

6 Statistical limit and covariance

We first prove the finite-band centered expansion and then address the limit target and covariance.

Proof of Theorem 2.1. For $\zeta_n = \mu_n - \mu$, Fourier duality and independence give

$$\mathbb{E}\|\zeta_n\|_{H^{-s}}^2 \leq \frac{1}{n} \sum_{j \in \mathbb{Z}^d} (1 + 4\pi^2|j|^2)^{-s} \leq C/n. \quad (24)$$

Thus $\epsilon_n = O_{\mathbb{P}}(n^{-1/2})$. The condition $\sqrt{n}h^M \rightarrow \infty$ implies that (11) contains ζ_n with probability tending to one, because $M > s + q$ and $h \rightarrow 0$. Propositions 5.1 and 5.2 give

$$\sqrt{n}\{\Phi_h(\mu_n) - \Phi_h(\mu)\} = n^{-1/2} \sum_{i=1}^n \psi_{h,\mu}(X_i) + O_{\mathbb{P}}(n^{-1/2}h^{-M}).$$

The remainder tends to zero. The centered triangular sum differs from the sum of the limiting influence by a random variable with variance $\|\psi_{h,\mu} - \psi_{0,\mu}\|_{L^2(\mu)}^2 \rightarrow 0$. The iid central limit theorem applies, including variance zero. Polarization, for instance $G_{ab} = \{\Phi(u_a + u_b) - \Phi(u_a - u_b)\}/4$, gives the joint expansion with influence (5).

The same event and argument control every fixed number of scores and bands, since all bandwidths are fixed multiples of h_n . Equation (16) and (7) give $\sum_{\ell} c_{\ell} G(h_n/2^{\ell}, \mu) = G^0 + O(h_n^{2K})$. Its scaled bias vanishes since $\beta > 1/(4K)$; the empirical remainders vanish since $\beta < 1/(2M)$. All bands use the same sample and have the same limiting influence. Their weights sum to one, so their linear terms combine into that influence, not into a sum of independent-band variances. This proves (8) and its asymptotic linear representation. $\square$

For the covariance estimator compute (20) at μ_n using a single bandwidth h_n and exact finite solves, and polarize to obtain $\widehat{\psi}_{ab,h_n}$. Set

$$\widehat{\Sigma}_{ab,cd} = \mu_n[\widehat{\psi}_{ab,h_n}\widehat{\psi}_{cd,h_n}]. \quad (25)$$

Proposition 6.1 (Covariance consistency). *Under Theorem 2.1, $\widehat{\Sigma} \rightarrow \Sigma$ entrywise in probability. For any fixed contrast with positive limiting variance, the corresponding studentized statistic tends to $N(0, 1)$. Inverse-covariance inference requires nonsingularity of the covariance on the selected contrast space.*

Proof. For a scalar score let χ_ν be the unique zero-Lebesgue-mean H^s representative of $D\Phi_h(\nu)$ modulo constants. Duality of mass-zero H^{-s} with H^s modulo constants and (17) give

$$\|\chi_\nu - \chi_\mu\|_{H^s} \leq Ch^{-M} \|\nu - \mu\|_{H^{-s}}.$$

The direct atomic calculation in Proposition 5.2 identifies the derivative at an atomic law as well. Thus $\psi_{h,\nu} = \chi_\nu - \nu\chi_\nu$. The population χ_μ is bounded uniformly in H^s by the high-order smooth-input expansions. Since ν is bounded in H^{-s} on the local ball,

$$\|\psi_{h,\nu} - \psi_{h,\mu}\|_{H^s} \leq Ch^{-M} \|\nu - \mu\|_{H^{-s}}. \quad (26)$$

This uses the mean-value integral on the whole segment, not differentiation of a pointwise asymptotic formula. For $\nu = \mu_n$ the right side is $o_{\mathbb{P}}(1)$. Split the variance error into

$$\mu_n(\psi_{h,\mu_n}^2 - \psi_{h,\mu}^2) + (\mu_n - \mu)(\psi_{h,\mu}^2) + \mu(\psi_{h,\mu}^2 - \psi_{0,\mu}^2).$$

The first vanishes by (26) and the algebra bound, the second is $O_{\mathbb{P}}(\epsilon_n)$, and the third vanishes by smooth influence convergence. Polarization and the same product estimates prove every covariance entry. The finite-band limiting influence is unchanged, so the one-band covariance serves (8). Slutsky's theorem gives the studentized conclusion when the limiting denominator is positive. $\square$

Corollary 6.2 (Exact nonparametric bootstrap). *Under Theorem 2.1, let $X_1^*, \dots, X_n^*$ be conditionally iid from μ_n , and compute the same exact multiband functional $\widehat{G}_n^*$ at $\mu_n^* = n^{-1} \sum_i \delta_{X_i^*}$, with the same n , bandwidths and coefficients. Its conditional law satisfies*

$$d_{\text{BL}}\left(\mathcal{L}\left[\sqrt{n} \text{vec}(\widehat{G}_n^* - \widehat{G}_n) \mid X_1, \dots, X_n\right], N(0, \Sigma)\right) \rightarrow 0$$

in probability, where d_{BL} is the bounded-Lipschitz metric on laws in the fixed entry space. The covariance may be singular. This assertion concerns exact empirical metrics, not an uncertified numerical bootstrap implementation.

Proof. Repeated bootstrap sites are merged and given their positive multinomial weights; zero weights are omitted. This leaves the measure and normalized metric unchanged. A one-site law has the zero mean-zero metric. Thus duplicates cause no definitional problem.

Let $E = H^{-s}$ and $C_s = \|\delta_x\|_E^2 < \infty$, independent of x . Conditional Hilbert-space variance gives

$$\mathbb{E}^* \|\mu_n^* - \mu_n\|_E^2 = \frac{1}{n} \{\mu_n \|\delta_x\|_E^2 - \|\mu_n\|_E^2\} \leq C_s/n.$$

The cross term with $\mu_n - \mu$ has conditional mean zero. Together with (24) this yields $\mathbb{E} \|\mu_n^* - \mu\|_E^2 \leq 2C_s/n$. Both measures consequently lie in the Taylor neighborhood centered at the deterministic true law μ , at all fixed bands, with joint probability tending to one. No uniform Taylor estimate centered at a random atomic law is needed.

Let ψ_h denote the finite-band population influence vector, and put $b_n = \sum_\ell c_\ell \psi_{h_n/2^\ell}$. It has mean zero, converges uniformly to $\text{vec } \psi$, and is uniformly bounded. Apply (18) separately to μ_n and μ_n^* at the same μ and subtract:

$$\sqrt{n} \text{vec}(\widehat{G}_n^* - \widehat{G}_n) = \sqrt{n}(\mu_n^* - \mu_n)b_n + r_n^*, \quad r_n^* = o_{\mathbb{P}}(1)$$

under the joint original/bootstrap law. Indeed, on the good neighborhood the expected remainder is at most $Cn^{-1/2}h_n^{-M} \rightarrow 0$, while its complement has probability tending to zero. The population tensor

and its bias cancel exactly. For each $\epsilon > 0$, $\mathbb{E}\mathbb{P}^*(\|r_n^*\| > \epsilon) = \mathbb{P}(\|r_n^*\| > \epsilon) \rightarrow 0$. Markov's inequality therefore gives conditional negligibility in original-sample probability.

Conditionally the leading term is a sum of n iid centered vectors $b_n(X_i^*) - \mu_n b_n$, divided by $\sqrt{n}$. They are uniformly bounded, and their conditional covariance

$$\mu_n(b_n b_n^T) - (\mu_n b_n)(\mu_n b_n)^T \longrightarrow \Sigma$$

in probability by uniform convergence of b_n and the ordinary laws of large numbers for the fixed limiting influence. Conditional Lindeberg holds. To obtain convergence of laws, take any subsequence and a further subsequence on which these covariances converge almost surely. On almost every such sample path the usual triangular-array CLT applies to the conditional arrays, including singular covariance. The subsequence principle proves bounded-Lipschitz convergence in probability on the original sequence. Finally, uniformly over bounded 1-Lipschitz tests, the effect of r_n^* is at most $\epsilon + 2\mathbb{P}^*(\|r_n^*\| > \epsilon)$. Letting $\epsilon \downarrow 0$ proves the corollary.

The bootstrap mechanism is the classical one for asymptotically linear functionals; fixed-regularization Sinkhorn and transport-functional precedents include [4, 7]. Here the shrinking-bandwidth remainder justifies its use for this inverse-metric statistic. $\square$

7 Canonical gradient and local regularity

Proof of Theorem 2.3. For a smooth centered density score v , let $f_t = f e^{tv} / \mu(e^{tv})$. Work modulo constants in the weak Poisson equation

$$E_f(\phi_a, z) = L_a(f; z), \quad E_f(x, z) = \mu(\nabla x \cdot \nabla z), \quad L_a(f; z) = \mu[(u_a - \mu u_a)z].$$

Uniform ellipticity for small t and the inverse on the quotient give differentiable smooth solutions; this can also be obtained by an implicit inverse between fixed Sobolev quotient spaces. The derivatives are

$$\dot{E}_f(x, z) = \mu[v \nabla x \cdot \nabla z], \quad \dot{L}_a(f; z) = \mu(v u_a^0 z) - \mu(v u_a) \mu z.$$

Using the centered baseline solutions as tests, differentiation of $G_{ab}^0 = E_f(\phi_a, \phi_b)$ and the two weak equations gives

$$\begin{aligned} DG_{ab}^0(f)[v] &= \dot{L}_a(f; \phi_b) + \dot{L}_b(f; \phi_a) - \dot{E}_f(\phi_a, \phi_b) \\ &= \mu[v(u_a^0 \phi_b + u_b^0 \phi_a - \nabla \phi_a \cdot \nabla \phi_b)] = \mu(v \psi_{ab}). \end{aligned} \quad (27)$$

This includes derivatives of the score means; they vanish here because the test solutions have baseline mean zero. Smooth centered functions are dense in $L_0^2(\mu)$, since f is bounded above and below. The smooth centered vector ψ is therefore the canonical gradient.

We state regularity precisely rather than presume a uniform density-class remainder. Fix a finite vector v of smooth centered functions and a fixed local vector t . Under the local exponential law $f_{n,t} = f \exp(t^T v / \sqrt{n}) / \mu \exp(t^T v / \sqrt{n})$ the likelihood satisfies

$$\log L_n = t^T n^{-1/2} \sum_i v(X_i) - \frac{1}{2} t^T \mu(v v^T) t + o_{\mathbb{P}}(1).$$

Boundedness of v also gives $\mathbb{E}_f L_n^2 = O(1)$, by a second-order expansion of its exponential moments. Hence these alternatives are contiguous to the baseline product laws. The baseline asymptotically linear representation in Theorem 2.1 retains its negligible remainder under each such alternative. The joint iid CLT for (ψ, v) and Le Cam's third lemma shift the mean of its linear term to $\mu(\psi v^T) t$, leaving covariance Σ unchanged. Alternatively this follows by expanding

$$\left\{ \frac{\mu \exp[(t^T v + i z^T \psi) / \sqrt{n}]}{\mu \exp[t^T v / \sqrt{n}]} \right\}^n.$$

Equation (27) gives exactly the same shift for $\sqrt{n}\{G^0(f_{n,t}) - G^0(f)\}$. After target centering the limit is $N(0, \Sigma)$, independently of t . This proves regularity along every fixed finite smooth exponential submodel. The certified numerical error also transfers by contiguity.

The information conclusion uses classical LAN convolution theory, not the covariance calculation alone; see [12, Theorems 25.20–25.21], and the transport applications [6, 7]. For a scalar contrast $b^T \text{vec } G^0$, write $\chi = b^T \text{vec } \psi$. The supremum of the parametric bounds $(\mu\chi v)^2/\mu(v^2)$ over smooth centered v is $\mu\chi^2$; if nonzero it is attained by the smooth score $v = \chi$. The convolution bound gives this Gaussian component for regular limiting laws, and the present estimator has exactly that normal limit. For the vector, choose a basis of the finite span of its smooth influence components and apply the multivariate LAN bound, or test every contrast. Thus the covariance bound is Σ , including singular directions, and it is attained. This is the claimed pointwise smooth-submodel efficiency; it entails no uniform coverage or minimax smoothing-rate conclusion. $\square$

8 Observed metric-error stopping

We now give an actual numerical target and a finite first-hit construction. This section is deterministic and uses no asymptotic empirical gap lower bound. All matrix products and inequalities in the certificates must be evaluated exactly or enclosed rigorously; floating eigenvalues alone are not certificates.

Fix distinct sites, positive probability weights w_i , a symmetric positive definite matrix K with strictly positive entries, a bandwidth $h > 0$, and finitely many prescribed score columns. The periodic heat Gram has these properties. Indeed its Fourier quadratic form has strictly positive multipliers; vanishing of that form forces the finite signed atomic measure to have all Fourier coefficients zero, hence to vanish. Duplicate sites can instead be merged.

8.1 Backward interpretation of a scaling iterate

For any positive vector b put

$$r_i = b_i \sum_j K_{ij} b_j w_j, \quad R = \sum_i w_i r_i, \quad v_i = \frac{w_i r_i}{R}, \quad a_i^{\text{tilt}} = \frac{\sqrt{R} b_i}{r_i}, \quad P_{ij} = \frac{K_{ij} b_i b_j w_j}{r_i}. \quad (28)$$

Direct substitution proves $a_i^{\text{tilt}} \sum_j K_{ij} a_j^{\text{tilt}} v_j = 1$. Thus P is exactly normalized at the residual-tilted measure v , with detailed balance $v_i P_{ij} = w_i w_j K_{ij} b_i b_j / R$. Its Euclidean symmetric representative is

$$\tilde{Q}_{ij} = \frac{K_{ij} b_i b_j \sqrt{w_i w_j}}{\sqrt{r_i r_j}}, \quad \tilde{\Pi} = (\sqrt{v})(\sqrt{v})^T. \quad (29)$$

Perron simplicity and positivity imply that its nonconstant eigenvalues lie in $(0, 1)$. The correct score columns at this intermediate law are centered at v , not at w . For example, if $w = (1/2, 1/2)$ but $v = (1/2 + e, 1/2 - e)$, the old-centered vector $(1, -1)$ has new mean $2e$ and cannot be the right side of a mean-zero inverse equation. This issue cannot be removed by merely making a normalization residual small.

8.2 A posteriori scale and tensor bounds

Let a be the exact scale at the original weights w , and let $m = \min_i w_i$. Define observed quantities

$$\eta_r = \|r - 1\|_{L^2(w)}, \quad r_{\min} = \min_i r_i, \quad r_{\max} = \max_i r_i, \quad R_0 = 2\eta_r / r_{\min}, \quad M_0 = R_0 / \sqrt{m}. \quad (30)$$

Lemma 8.1 (Observed logarithmic scale bound). *If*

$$q_0 = 2(r_{\max}/r_{\min})(e^{2M_0} - 1) \leq 1/2, \quad (31)$$

then $\|\log(a/b)\|_{\infty} \leq M_0$.

Proof. In $L^2(w)$ write $B = \text{diag}(b)K_w \text{diag}(b)$, where $(K_w v)_i = \sum_j K_{ij} w_j v_j$, and $F(x) = e^x b K_w (e^x b) - 1$. Positivity gives $B \geq 0$ and $\|B\| \leq r_{\max}$, the latter by the nonnegative row-sum bound and symmetric similarity. $J_0 = DF(0) = \text{diag}(r) + B \geq r_{\min} I$. On the closed $L^2(w)$ ball of radius R_0 we have $\|x\|_\infty \leq M_0$. If $D = \text{diag}(e^x)$, then

$$DF(x) = \text{diag}(r(x)) + DBD, \quad \|DF(x) - J_0\| \leq 2r_{\max}(e^{2M_0} - 1).$$

For the diagonal part use $e^{-2M_0} r_i \leq r_i(x) \leq e^{2M_0} r_i$; for the matrix part expand $DBD - B = (D - I)BD + B(D - I)$. The Newton map $x - J_0^{-1}F(x)$ is q_0 -Lipschitz on the ball and its value at zero has norm at most $R_0/2$. It therefore maps the ball to itself and contracts. Its fixed point is the unique exact positive scale. The zero-residual case is immediate. The ℓ^∞ estimate follows from the $L^2(w)$ radius. No smallest kernel eigenvalue is used. $\square$

Let Q be the exact symmetric matrix at w and $\Pi_w = (\sqrt{w})(\sqrt{w})^T$. With a certified upper bound $p_0 \geq \|\tilde{\Pi} - \Pi_w\|$ define

$$L = M_0 + \frac{1}{2} \max_i |\log r_i|, \quad \epsilon_0 = e^{2L} - 1, \quad e_0 = \epsilon_0 + p_0. \quad (32)$$

For example $p_0 = 2\|\sqrt{v} - \sqrt{w}\|_2$ is valid. Equation (29) gives $\tilde{Q} = DQD$ with $D_{ii} = b_i/(a_i \sqrt{r_i})$. Hence Lemma 8.1 implies $\|\tilde{Q} - Q\| \leq \epsilon_0$. Let $g \in (0, 1]$ be a certified lower bound on the nonconstant gap of $\tilde{Q}$ and require $e_0 < g$. Both shifted matrices $I - Q + \Pi_w$ and $I - \tilde{Q} + \tilde{\Pi}$ are positive definite; Weyl's inequality bounds the former below by $(g - e_0)I$. The resolvent identity and (2) give

$$\|T(\tilde{Q}, \tilde{\Pi}) - T(Q, \Pi_w)\| \leq B_0 := \frac{h}{2} \left\{ \frac{e_0}{g(g - e_0)} + \epsilon_0 + \frac{p_0}{2} \right\}. \quad (33)$$

Indeed the plus inverse norms are at most one, while the minus inverse product costs $1/[g(g - e_0)]$.

Let S be the matrix of the prescribed score values and form the common-Euclidean columns

$$\begin{aligned} Y_w &= \text{diag}(\sqrt{w})[S - 1(w^T S)], & Y_v &= \text{diag}(\sqrt{v})[S - 1(v^T S)], \\ D_Y &= Y_v - Y_w, & y &= \|Y_w\|_F, & d_Y &= \|D_Y\|_F. \end{aligned}$$

The exact tilted tensor has norm at most $h/(2g)$. Expanding the difference of the two metric matrices as

$$Y_w^T(\tilde{T} - T)Y_w + D_Y^T \tilde{T} Y_w + Y_w^T \tilde{T} D_Y + D_Y^T \tilde{T} D_Y$$

therefore proves the observed allowance

$$\|G(h, v) - G(h, w)\|_{\text{op}} \leq E_{\text{scale}} := B_0 y^2 + \frac{h}{2g} (2y d_Y + d_Y^2). \quad (34)$$

Changing only the operator and keeping the old mass factors or the old center would not establish this inequality.

8.3 Inverse residuals and the complete certificate

Work now in $L^2(v)$ with score columns $Y = S - 1(v^T S)$. On its mean-zero subspace let $A_- = I - P$; let $A_+ = I + P$. Choose minus trial columns X_- centered at v and arbitrary plus trial columns X_+ . Set

$$\begin{aligned} Z_\pm &= Y - A_\pm X_\pm, \\ W_\pm &= Y^* X_\pm + X_\pm^* Y - X_\pm^* A_\pm X_\pm, & G^{\text{trial}} &= \frac{h}{2}(W_- - W_+), \end{aligned}$$

where stars include the actual v weights. Expanding the residual proves the classical identity

$$Y^* A_\pm^{-1} Y - W_\pm = Z_\pm^* A_\pm^{-1} Z_\pm.$$

As $A_- \geq gI$ on the complement and $A_+ \geq I$, it follows in full matrix Loewner order that

$$-\frac{h}{2}Z_+^*Z_+ \leq G(h, \nu) - G^{\text{trial}} \leq \frac{h}{2g}Z_-^*Z_-.$$

In particular

$$E_{\text{solve}} := \max \left\{ \frac{h}{2g} \|Z_-\|_{L^2(\nu), F}^2, \frac{h}{2} \|Z_+\|_{L^2(\nu), F}^2 \right\} \geq \|G(h, \nu) - G^{\text{trial}}\|_{\text{op}}. \quad (35)$$

The full residual matrices retain off-diagonal correlations; no entrywise independence is assumed. Combining (34)–(35) gives the complete observed certificate

$$E_{\text{scale}} + E_{\text{solve}} \leq \eta \implies \|G^{\text{trial}} - G(h, w)\|_{\text{op}} \leq \eta. \quad (36)$$

Its target is the metric at the original empirical law.

8.4 Actual first hit and finite termination

Proposition 8.2 (Finite first-hit construction). *For every finite distinct sample set and every $\eta > 0$, there is an actual scaling and inverse iteration that stops with (36). In ideal real arithmetic it terminates finitely. It has a certified finite-arithmetic realization under effective convergent enclosures of inputs and gap certification that eventually detects the positive gap.*

Proof. Start $b = 1$ and perform damped symmetric scaling

$$b_i^{\text{next}} = \sqrt{\frac{b_i}{(K_w b)_i}}. \quad (37)$$

To see actual convergence, let a be the exact scale and $t = b/a$. Then $t^{\text{next}} = \sqrt{t/(P_* t)}$, where P_* is the exact balanced matrix. For a strictly positive matrix its Hilbert projective contraction gives

$$\text{osc}(\log(P_* t)) \leq \theta \text{osc}(\log t), \quad \theta = \tanh(\Delta(K)/4) < 1,$$

where $\Delta(K) = \max_{i,j,k,l} \log(K_{ik}K_{jl}/(K_{il}K_{jk}))$. Consequently $\text{osc}(\log t^{\text{next}}) \leq (1 + \theta) \text{osc}(\log t)/2$. Moreover each new log coordinate lies between $-\text{osc}(\log t)/2$ and $\text{osc}(\log t)/2$, since $P_* t$ lies between the minimum and maximum of t . Thus $\log(b/a) \rightarrow 0$, not merely convergence modulo constants. Classical positive-kernel contraction is also discussed in [1, 3].

At each iterate evaluate (31), a certified gap g for $\tilde{Q}$, and (34). Stop scaling at the first iterate for which

$$q_0 \leq 1/2, \quad e_0 < g, \quad E_{\text{scale}} < \eta/2.$$

This hit is finite: convergence implies $r \rightarrow 1$, $\nu \rightarrow w$ and $M_0, q_0, e_0, p_0, d_Y, B_0, E_{\text{scale}} \rightarrow 0$. The exact original finite matrix has a positive gap, and so do the nearby tilted matrices. An eventually informative gap enclosure can have a lower bound at least half their positive limiting gap. These strict tests are then satisfied.

Freeze that iterate and its tilted law. Start $X_- = X_+ = 0$ and apply

$$X_-^{\text{next}} = P X_- + Y, \quad X_+^{\text{next}} = (I - P) X_+ / 2 + Y / 2.$$

Minus columns remain centered in exact arithmetic. On that space the first iteration contracts with nonconstant eigenvalues in $(0, 1)$; the second has eigenvalues $(1 - \lambda)/2 \in [0, 1/2]$. Their actual residuals tend to zero. Stop at the first inverse iteration with $E_{\text{solve}} < \eta/2$. This hit is also finite and proves (36).

For certified finite arithmetic, arbitrary real observations do not alone specify effective numerical input. Assume convergent rigorous enclosures for sites, scores and kernel values. Spectral enclosures

must eventually detect a positive gap; an arbitrary valid but excessively small lower bound is not enough. Increase precision or dovetail it with iterate count. For an existence-level realization, represent each exact finite iterate as a finite real-arithmetic expression, enclose it at increasing precision, choose positive rational approximations, and freshly certify the displayed tests at those concrete vectors. A sufficiently late exact iterate satisfies all strict tests with a positive margin. Continuity ensures sufficiently close rational approximations do too. Treat the inverse trials in the same way, recentring minus columns and certifying their actual residuals. This avoids assuming convergence of an arbitrarily rounded recurrence. The dovetailing finds a valid finite hit, with no efficient expression-evaluation or bit-complexity assertion. $\square$

Proof of Corollary 2.2. Use Proposition 8.2 at each band with $\eta = n^{-3/4}/C_K$. The combined numerical error is at most $n^{-3/4}$ pathwise on every finite distinct sample set. Multiplying by $\sqrt{n}$ gives $n^{-1/4} \rightarrow 0$; Slutsky's theorem applies. Distinctness holds almost surely under the smooth density. No statistical lower bound on the realized empirical gap was used in this numerical transfer. $\square$

There is no practical runtime conclusion hidden in finite termination. In particular $\Delta(K)$ can grow as $1/h$, making the projective bound extremely poor. Likewise a normalization first hit does not perform an inverse solve. A raw unnormalized inverse is unsafe: for a two-point exact matrix with tangent eigenvalue $\lambda = e^{-ch}$, replacing the exact scale by $b = \lambda^{-1/2}a$ gives residual $r_i = \lambda^{-1} = 1 + O(h)$ but raw tangent eigenvalue one, so its supposed tensor inverse is singular, although $\nabla \log r = 0$ for its constant residual. The residual-tilted row normalization and the separate inverse residual test are substantive parts of the certificate.

9 A circle example and the complete variance

Let $f = 1$ on $\mathbb{T}$, $u(x) = \sqrt{2} \cos(2\pi kx)$ with $k \neq 0$, and $\omega = 4\pi^2 k^2$. Then $\phi = u/\omega$, $G^0 = 1/\omega$ and

$$\psi_0(x) = \frac{3}{\omega} \cos(4\pi kx), \quad \Sigma = \frac{9}{2\omega^2}.$$

The exact finite-band population multiplier is

$$\lambda = e^{-h\omega/2}, \quad t_h = \frac{h\lambda}{1 - \lambda^2} = \frac{h}{2 \sinh(h\omega/2)}.$$

Here the exact centered influence is $A_h \cos(4\pi kx)$ with

$$A_h = t_h \left\{ 1 + \frac{(1 + \lambda^2)^2}{1 + \lambda^4} \right\}. \quad (38)$$

For completeness, the scaling derivative in a mass direction ζ solves $(I + H_h)\alpha = -H_h\zeta$. Since $u^2 = 1 + \cos(4\pi kx)$ and the heat multiplier on the doubled frequency is λ^4 , kernel duality gives

$$\langle u, \dot{P}u \rangle_{L^2} = \lambda \frac{1 - \lambda^4}{1 + \lambda^4} \zeta[\cos(4\pi kx)].$$

The filter $q_h(z) = hz/(1 - z^2)$ has derivative $q'_h(\lambda) = h(1 + \lambda^2)/(1 - \lambda^2)^2$. Adding the outer-measure derivative $t_h\zeta(u^2)$ gives (38), hence $A_h \rightarrow 3/\omega$.

Taking $s = 1$ gives $M = 9/2$. Three bands suffice:

$$(c_0, c_1, c_2) = (1/45, -4/9, 64/45), \quad \sum |c_\ell| = 17/9.$$

They cancel h^2 and h^4 . With $\beta = 2/21$, both $\sqrt{n}h^6$ and $n^{-1/2}h^{-9/2}$ equal $n^{-1/14}$. This is a conservative sufficient balancing example, not an optimal rate or a practical sample-size recommendation.

Freezing the population operator and merely averaging $u\phi$ would instead produce variance $1/(2\omega^2)$. That is an oracle quadrature proxy for a different experiment; it omits normalized-operator and inverse sensitivity. It does not contradict the full-model information bound. If uniformity is known as a model restriction, the target $1/\omega$ is already known and no sample estimator is needed. Efficiency here is relative to unknown nearby smooth positive densities.

10 Scope and further questions

The sufficient bandwidth budget leaves a nonempty regime in every fixed dimension under the stated infinite smoothness. It is intentionally conservative. Sharper common-space resolvent estimates and more economical scalar perturbation bounds may reduce the number of bands and improve admissible bandwidths. Lower regularity and densities that vanish require new arguments: positivity, high-order bias and the relevant elliptic inverse are used essentially in this proof.

The results concern a finite collection of fixed scores at one density. They do not give a parameter-uniform process, curvature inference, growing-dimensional covariance, data-chosen scores or bandwidth selection. Exact empirical covariance is consistent, but a numerical implementation of its influence solves requires an additional verified error budget. The metric certificate gives existence of a finite certified computation, not useful finite-sample cost or coverage guarantees. Those distinctions also separate the theorem from fixed-support horizontal geometry and from fixed squared-cost transport inference.

AI assistance. AI systems assisted with mathematical development, literature discovery, proof checking and manuscript preparation. The author is responsible for the statements, attribution and presentation.

References

- [1] C. L. Wormell and S. Reich, *Spectral convergence of diffusion maps: improved error bounds and an alternative normalisation*, arXiv:2006.02037v3, 2021.
- [2] X. Cheng and B. Landa, *Bi-stochastically normalized graph Laplacian: convergence to manifold Laplacian and robustness to outlier noise*, arXiv:2206.11386v3, 2024.
- [3] N. Kessler, R. Magnet and J. Feydy, *Sinkhorn Normalization of Diffusion Kernels*, Proceedings of Machine Learning Research 306 (2026), 56764–56795; arXiv:2507.06161v2.
- [4] Z. Goldfeld, K. Kato, G. Rioux and R. Sadhu, *Limit Theorems for Entropic Optimal Transport Maps and the Sinkhorn Divergence*, Electronic Journal of Statistics (2024); arXiv:2207.08683.
- [5] C. Li, N. García Trillos, H. Li and L. Suchan, *Central limit theorems for the eigenvalues of graph Laplacians on data clouds*, arXiv:2507.18803v1, 2025.
- [6] T. Manole, S. Balakrishnan, J. Niles-Weed and L. Wasserman, *Plugin Estimation of Smooth Optimal Transport Maps*, arXiv:2107.12364v3, 2024.
- [7] H. Kono, *Central Limit Theorems for Optimal Transport Functionals*, arXiv:2610.27857v1, 2026.
- [8] H. Lavenant, J. Luckhardt, G. Mordant, B. Schmitzer and L. Tamanini, *The Riemannian geometry of Sinkhorn divergences*, Ann. Inst. H. Poincaré Anal. Non Linéaire 43 (2026), 1079–1144; arXiv:2405.04987v2.
- [9] G. Mordant and L. Xu, *Convergence of the Sinkhorn Riemannian metric for finitely supported measures*, arXiv:2609.14651v1, 2026.
- [10] S. Schubr, *Sharp convergence of heat-kernel Sinkhorn metrics*, public edition 1, 2026. Luminary Research Archive.
- [11] S. Schubr, *Leading bias and positive correction of heat-kernel Sinkhorn metrics*, public edition 1, 2026. Luminary Research Archive.
- [12] A. W. van der Vaart, *Asymptotic Statistics*, Cambridge University Press, 1998.